


Schoeneberger et al. 1998

Field Book for Describing and Sampling Soils

59365



Version 1.1

Revised, Refined, and Compiled by:
P.J. Schoeneberger, D.A. Wysocki,
E.C. Benham, and W.D. Broderson

National Soil Survey Center
Natural Resources Conservation Service
U.S. Department of Agriculture
Lincoln, Nebraska

Received by ER-RPF

SEP 18 1998

cmf



13531

FIELD SAMPLING

Compiled by: P.J. Schoeneberger and D.A. Wysocki, NRCS, Lincoln, NE.

INTRODUCTION

This section contains a variety of miscellaneous information pertinent to the sampling of soils in the field.

Additional details of soil sampling for the National Soil Survey Laboratory (NRCS, Lincoln, NE) are provided in Soil Survey Investigations Report No. 42 (Soil Survey Staff, 1996)

SOIL SAMPLING

The objective of the task determines the methodology and the location of the soil material collected for analysis. Sampling for Taxonomic Classification purposes involves different strategies than sampling for soil fertility, stratigraphy, hydric conditions, etc. There are several general types of samples and sampling strategies that are commonplace in soil survey


SOIL SAMPLE KINDS -

Reference Samples (also loosely referred to as "grab" samples) - This is applied to any samples that are collected for very specific, limited analyses; e.g., only pH. Commonly, reference samples are not collected for all soil layers in a profile; e.g., only the top 10 cm, only the most root restrictive layer, etc.

Characterization Samples - These samples include sufficient physical and chemical soil analyses, from virtually all layers, to fully characterize a soil profile for Soil Taxonomic and general interpretive purposes. The specific analyses required vary with the type of material; e.g., a Mollisol requires some different analyses than does an Andisol. Nonetheless, a wide complement of data (i.e., pH, particle size analysis, Cation Exchange Capacity, ECEC, Base Saturation, Organic Carbon content, etc.) are determined for all major soil layers.

SAMPLING STRATEGIES - [To be developed.]

Field Book for Describing and Sampling Soils



Version 1.1

Revised, Refined, and Compiled by:
P.J. Schoeneberger, D.A. Wysocki,
E.C. Benham, and W.D. Broderson

National Soil Survey Center
Natural Resources Conservation Service
U.S. Department of Agriculture
Lincoln, Nebraska

ACKNOWLEDGMENTS

The science and knowledge in this document are distilled from the collective experience of thousands of dedicated Soil Scientists during the nearly 100 years of the National Cooperative Soil Survey Program. A special thanks is due to these largely unknown stewards of the natural resources of this nation.

Special thanks and recognition are extended to those who contributed extensively to the preparation and production of this book: The 75 soil scientists from the NRCS along with NCSS cooperators who reviewed and improved it; Tammy Nepple for document preparation and graphics; Howard Camp for graphics; Jim Culver for sponsoring it; and the NRCS Soil Survey Division for funding it.

Proper citation for this document is:

Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., and Broderson, W.D. 1998. Field book for describing and sampling soils. Natural Resources Conservation Service, USDA, National Soil Survey Center, Lincoln, NE.

Cover Photo: Soil profile of a Segno fine sandy loam (*Plinthic Paleudalf*) showing reticulate masses or blocks of plinthite at 30 inches (profile tape is in feet). *Courtesy of Frankie F. Wheeler, NRCS, Temple TX; and Larry Ratliff (retired), National Soil Survey Center, Lincoln, NE.*

Use of trade or firm names is for reader information only, and does not constitute endorsement or recommended use by the U.S. Department of Agriculture of any commercial product or service.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal employment opportunity employer.

FOREWORD

Purpose: The following instructions, definitions, concepts, and codes are a field guide for making or reading soil descriptions and sampling soils as presently practiced in the USA.

Background: This document is an expanded and updated version of earlier guides and short-hand notation released by the Soil Conservation Service (e.g., Spartanburg, SC, 1961; Western Technical Center, Portland, OR, 1974). The knowledge base in those releases was developed by soil scientists during the formative years of the Soil Survey Program. Roy Simonson and others summarized this information in the 1950s (e.g., Soil Survey Staff, 1951; Soil Survey Staff, 1962). This document summarizes our present knowledge base.

Standards: This book summarizes the current National Cooperative Soil Survey conventions for describing soils. Where the content deviates from the initial sources (SSM, 1993; NSSH, 1996; PDP 3.6, 1996; etc.), excepting errors, this document updates them.

Regarding PEDON (PDP 3.5 / 3.6): This document is intended to be both current and useable by the entire soil science community; it is not a guide on "How to use the Pedon Description Program (PDP)". At this time, PDP is the most dated and therefore the least compatible NRCS document relative to the Soil Survey Manual, National Soil Survey Handbook, Keys to Soil Taxonomy, and NASIS. Differences and linkages between PDP 3.6 and NASIS are shown, where reasonable to do so, as an aid to conversions. Future releases of this book are unlikely to include PDP materials.

Standard procedures and terms for describing soils have changed and increased in recent years (e.g., redoximorphic features). Coincident with these changes has been the development and use of computer databases to store soil descriptions and information. The nature of databases, for better or worse, requires consistent and "correct" use of terms.

Sources: This Field Book draws from several primary sources: The Soil Survey Manual (Soil Survey Staff, 1993); the PEDON Description Program (PDP) Version 4 Design Documents (Soil Survey Staff, 1996d); and the National Soil Survey Handbook (NSSH) -- Parts 618 and 629 (Soil Survey Staff, 1996c). Other less pervasive sources are footnoted throughout the Field Book to encourage access to original information.

Brevity: In a field book, brevity is efficiency. Despite this book's apparent length, the criteria, definitions, and concepts presented here are condensed. We urge users to review the more comprehensive information in the original sources to avoid errors due to our brevity.

Units: It is critical to specify and consistently use units for describing a soil. Metric units are preferred. NASIS requires metric units. (In PDP, you can choose Metric or English units.)

Format: The "Site Description Section" and "Profile Description Section" in this book generally follow conventional profile description format and sequence (e.g., SCS-232, December 1984). Some data elements (descriptors) are rearranged in this document into a sequence that is more compatible with the description process in the field (e.g., **Horizon Boundary** is next to **Horizon Depth**, rather than at the very end). This sequence is somewhat different from and does not supersede the conventions followed in writing formal soil descriptions for Soil Survey Reports or Official Soil Series Descriptions (i.e., National Soil Survey Handbook, Part 614, p. 13-22; Soil Survey Staff, 1996).

Codes: Short-hand notation is listed in the *Code* column for each descriptor. Long-standing, conventional codes are retained because of their widespread recognition. Some codes of recent origin have been changed to make them more logical. Some data elements have different codes in various systems [e.g., conventional (Conv.) vs. NASIS vs. PEDON Description Program codes (PDP)] and several columns may be shown to facilitate conversions. The preferred, standard code is shown **bold**. If only 1 untitled code column is shown, it can be assumed that the conventional, NASIS, and PDP codes are all the same.

Standard Terms vs. Creativity: Describe and record what you observe. Choice lists in this document are a minimal set of descriptors. Use additional descriptors, notes, and sketches to record pertinent information and/or features for which no data element exists. Record such information as free-hand notes under **Miscellaneous Field Notes** (or **User Defined Entries** in PDP).

Changes: Soil Science is an evolving field. Changes to this Field Book should and will occur. Please send comments or suggestions to the authors at the National Soil Survey Center, USDA-NRCS; 100 Centennial Mall North, Rm. 152; Lincoln, NE 68508-3866.

TABLE OF CONTENTS

ACKNOWLEDGMENTS	i
FOREWORD	ii
SITE DESCRIPTION	1-1
Describer(s) Name	1-1
Date	1-1
Climate	1-1
Weather Conditions	1-1
Air Temperature	1-1
Soil Temperature	1-1
[Soil Temperature, Soil Temperature Depth]	
Location	1-2
[Latitude, Longitude, Datum Name]	
Topographic Quadrangle	1-2
Soil Survey Area Identification Number (SSID)	1-2
County FIPS Code	1-3
MLRA	1-3
Transects	1-3
[Transect ID, Stop Number, Interval]	
Series Name	1-4
Geomorphic Information	1-4
Physiographic Location	1-4
[Physiographic Division, Physiographic Province, Physiographic Section, State Physiographic Area, Local Physiographic/Geographic Name]	
Geomorphic Description	1-4
[Landscape, Landform, Microfeature, Anthropogenic Feature]	
Surface Morphometry	1-4
[Elevation, Slope Aspect, Slope Gradient, Slope Complexity, Slope Shape, Hillslope - Profile Position, Geomorphic Component (Hills, Terraces, Mountains, Flat Plains), Microrelief]	
Water Status	1-9
Drainage	1-9
Flooding	1-10
[Frequency, Duration, Months]	
Ponding	1-11
[Frequency, Depth, Duration]	
(Soil) Water State	1-12
Depth To Water Table	1-13
(Seasonal) High Water Table - Kind	1-13

Vegetation / Land Cover	1-14
[Earth Cover - Kind, Plant Symbol, Plant Common Name, Plant Scientific Name]	
Parent Material.....	1-16
Bedrock.....	1-18
[Kind, Fracture Interval Class, Hardness, Weathering Class, Depth]	
Erosion.....	1-20
[Kind, Degree Class]	
Runoff	1-21
Surface Runoff.....	1-21
The Index (Of) Surface Runoff Class.....	1-22
Surface Fragments (Formerly Surface Stoniness).....	1-22
Diagnostic Horizons Or Properties.....	1-23
[Kind, Depth]	
References.....	1-24
Profile / Pedon Description.....	2-1
Observation Method.....	2-1
[Kind, Relative Size]	
Taxonomic Classification	2-2
Horizon Nomenclature	2-2
Master, Transitional And Common Horizon Combinations.....	2-2
Horizon Suffixes.....	2-3
Other Horizon Modifiers.....	2-4
[Numerical Prefixes, Numerical Suffixes, The Prime]	
Diagnostic Horizons.....	2-4
Horizon Depth.....	2-4
Horizon Thickness	2-5
Horizon Boundary.....	2-5
[Distinctness, Topography]	
Soil Color	2-7
Decision Flowchart For Describing Soil Colors	2-7
(Soil) Matrix Color.....	2-7
[(Soil) Color, Moisture State, Location Or Condition]	
Mottles	2-9
[Quantity, Size, Contrast, Color, Moisture State, Shape]	
Redoximorphic Features (RMF) Discussion	2-13
Redoximorphic Features.....	2-14
[Kind, Quantity, Size, Contrast, Color, Moisture State, Shape, Location, Hardness, Boundary]	
Concentrations Discussion	2-17
Concentrations.....	2-18
[Kind, Quantity (Percent Of Area Covered), Size, Contrast, Color, Moisture State, Shape, Location, Hardness, Boundary]	
Ped & Void Surface Features	2-24
[Kind, Amount, Continuity, Distinctness, Location, Color]	

(Soil) Texture	2-28
Texture Class.....	2-28
Texture Triangle (Fine Earth).....	2-29
Texture Modifiers	2-29
Texture Modifiers.....	2-32
Compositional Texture Modifiers	2-33
Terms Used In Lieu Of Texture	2-34
Rock And Other Fragments	2-35
[Kind, Volume Percent, Roundness, Size Classes And Descriptive Terms]	
(Soil) Structure	2-38
[Type, Grade, Size]	
Consistence	2-46
Rupture Resistance	2-46
[Blocks, Peds, And Clods; Surface Crust And Plates]	
Cementing Agents	2-48
Manner Of Failure.....	2-49
Stickiness	2-50
Plasticity	2-50
Penetration Resistance.....	2-51
Excavation Difficulty	2-52
Roots.....	2-53
[Quantity, Size, Quantity (diagram), Location]	
Pores Discussion	2-56
Pores.....	2-56
[Quantity, Size, Shape, Vertical Continuity]	
Cracks.....	2-58
[Kind, Depth, Relative Frequency]	
Special Features	2-61
[Kind, Area (%) Occupied]	
Permeability / Saturated Hydraulic Conductivity (Discussion)	2-62
Permeability	2-63
Saturated Hydraulic Conductivity (K_{sa}).....	2-63
Chemical Response.....	2-64
Reaction (pH).....	2-64
Effervescence.....	2-65
[Class, Location, Chemical Agent]	
Reduced Conditions	2-66
Salinity	2-66
Sodium Adsorption Ratio (SAR)	2-66
Odor	2-67
Miscellaneous Field Notes.....	2-67
Minimum Data Set (For A Soil Description)	2-67
Profile Description Form	2-67
Profile Description Example.....	2-67

Profile Description Report Example (For Soil Survey Reports)	2-68
References.....	2-68
Geomorphic Description.....	3-1
Geomorphic Description System (Version 2.06 - 9/4/97).....	3-1
Part I: Physiographic Location.....	3-2
Part II: Geomorphic Description (Outline).....	3-10
Part II: Geomorphic Description.....	3-11
Part III: Surface Morphometry.....	3-22
References.....	3-26
Soil Taxonomy	4-1
Introduction	4-1
Horizon Nomenclature	4-1
Master And Transitional Horizons	4-1
Horizon Suffixes.....	4-3
Horizon Nomenclature Conversion Charts.....	4-5
Texture Triangle: Soil Texture Family Classes.....	4-7
Combined Texture Triangles: Soil Texture Family Classes And Fine Earth Texture Classes.....	4-8
References.....	4-9
Geology	5-1
Introduction	5-1
Bedrock - Kind	5-1
Rock Charts	5-3
Igneous Rocks Table.....	5-4
Metamorphic Rocks Table.....	5-5
Sedimentary And Volcaniclastic Rocks Table.....	5-6
North American Geologic Time Scale.....	5-7
Till Terms	5-8
Volcaniclastic Terms.....	5-9
References.....	5-10
Location.....	6-1
Public Land Survey.....	6-1
Townships And Ranges.....	6-1
Sections	6-2
Sub-Divisions.....	6-3
State Plane Coordinate System.....	6-4
Universal Transverse Mercator (UTM) Rectangular Coordinate System.....	6-4
References.....	6-5
Miscellaneous	7-1
Examples Of Percent Of Area Covered.....	7-1

Measurement Equivalents & Conversions.....	7-2
Metric To English.....	7-2
English To Metric.....	7-3
Common Conversion Factors.....	7-4
Guide To Map Scales And Minimum-Size Delineations.....	7-7
Common Soil Map Symbols (Traditional).....	7-8
Field Sampling.....	8-1
Introduction.....	8-1
Soil Sampling.....	8-1
Soil Sample Kinds.....	8-1
Reference Samples.....	8-1
Characterization Samples.....	8-1
Sampling Strategies.....	8-1
Field Equipment Checklist.....	8-2
Examples Of Common Field Sampling Equipment.....	8-3
References.....	8-4
Index.....	9-1

SITE DESCRIPTION

Compiled by: P.J. Schoeneberger, D.A. Wysocki, E.C. Benham, NRCS,
Lincoln, NE; W. D. Broderson, NRCS, Salt Lake City, UT.

DESCRIBER(S) NAME

NAME (or initials) - Record the observer(s) who make the description;
e.g., Erling E. Gamble or EEG.

DATE

MONTH / DAY / YEAR - Record the date of the observations. Use two digits
for each; e.g., 05/21/96 (for May 21, 1996).

CLIMATE

Document the prevailing, general weather conditions at the time of
observation. (Not a data element in PDP; a site-condition which affects
some field methods; e.g., K_{sa}). Record the dominant **Weather Conditions**
and **Air Temperature**; e.g., *Rain, 27°C*.

Weather Conditions	Code
Sunny / Clear	SU
Partly Cloudy	PC
Overcast	OV
Rain	RA
Sleet	SL
Snow	SN

AIR TEMPERATURE - The ambient air temperature at approximately chest
height (in degrees, Celsius or Fahrenheit); e.g., 27°C.

SOIL TEMPERATURE - Record the ambient **Soil Temperature** and the
Depth at which it is determined; e.g., 22°C, 50 cm. (**NOTE:** Soil Taxonomy
generally requires a depth of 50 cm.) Soil temperature should only be
determined from a freshly excavated surface that reflects the ambient soil
conditions. Avoid surfaces equilibrated with air temperatures.

Soil Temperature - Record the soil temperature (in °C or °F).

Soil Temperature Depth - Record the depth at which the ambient soil temperature is measured; e.g., *50 cm*.

LOCATION

Record the geographical location of the point / area of interest as precisely as possible. Latitude and Longitude are preferred. Record: degrees, minutes, seconds (decimal seconds), direction, and associated Datum.

LATITUDE - e.g., *46° 10' 19.38" N. Lat.*

LONGITUDE - e.g., *95° 23' 47.16" W. Long.*

NOTE: Latitude and Longitude are required in NASIS. For other location descriptors (e.g., *Public Land Survey, UTM, Metes and Bounds, State Plane Coordinates*, etc.), see the "Location Section".

DATUM NAME - Record the reference datum for the Latitude and Longitude from either topographic maps or GPS configuration used; e.g., *NAD 1983* (North American Datum, 1983) for most of the U.S.

TOPOGRAPHIC QUADRANGLE

Record the appropriate topographic map name (i.e., Quadrangle Name) covering the observation site (commonly a USGS topographic map). Include the scale (or map "series") and the year printed; e.g., *Pollard Creek - NW; TX; 1:24,000; 1972.*

SOIL SURVEY AREA IDENTIFICATION NUMBER (SSID)

An identification number must be assigned if samples are collected for analyses at the National Soil Survey Laboratory. This identifier consists of four required and one optional part. These are:

- 1) The letter *S* (for "soil characterization sample") and the two-digit calendar year; e.g., *S96* (for 1996).
- 2) The two-character state abbreviation; e.g., *OK* (for Oklahoma). For non-USA samples, use the abbreviation *FN*.
- 3) The three-digit county FIPS code; e.g., *061* (for Haskell County, OK). For non-USA samples, use the appropriate three-digit GSA world-wide geographical location code (Public Building Service, 1996).

- 4) A three-digit, sequential code to identify the individual pedons sampled within the county or other survey area during any given calendar year; e.g., 005. (**NOTE:** This sequential code starts over with 001 each January 1.)
- 5) (Optional) A one-character sub-sample code. This is generally used to indicate some relationship (such as satellite samples) between sampling sites; e.g., A.

A complete example is *S96OK061005A*. [Translation: A pedon sampled for soil characterization during 1996 (*S96*), from Oklahoma (*OK*), in Haskell County (*061*), it's the fifth pedon (*005*) sampled in that county during 1996, and it's a satellite sample (*A*) related to the primary pedon.]

COUNTY FIPS CODE

This is the three-digit FIPS code for the county (National Institute of Standards and Technology, 1990) in a U.S. state in which the pedon or site is located. It is usually an odd number; e.g., 061 (for Haskell County, OK). For non-USA samples, enter *FN* followed by the appropriate three-digit GSA world-wide geographical location code (Public Building Service, 1996); e.g., *FN260* (for Canada).

MLRA

This is the one- to three-digit (and one-character sub-unit, if applicable) Major Land Resource Area identifier (SCS, 1981); e.g., 58C (for Northern Rolling High Plains - Northeastern Part).

TRANSECTS

If the soil description is a point along a transect, record the appropriate transect information: **Transect ID**, **Stop Number**, and **Interval**. In NASIS, additional transect information can be recorded: **Transect Kind** (random point, fixed interval), **Transect Selection Method** (random, biased), **Delineation Size** (units), **Transect Direction** (compass heading in degrees).

TRANSECT ID - This is a four- to five-digit number that identifies the transect; e.g., 0029 (the 29th transect within the survey area).

STOP NUMBER - If the sample/pedon is part of a transect, enter the two-digit stop number along the transect; e.g., 07. (**NOTE:** NASIS allows up to 13 characters).

INTERVAL - Record distances between observation points, compass bearings, and GPS coordinates; or draw a route map in the **Field Notes** ("User Defined Section"). In PDP, if the observation is part of a transect, enter the distance (in feet or meters) between points; e.g., *30 m*.

SERIES NAME

This is the assumed Soil Series name at the time of the description; e.g., *Cecil*. (If unknown, enter *SND* for "Series Not Designated"). (**NOTE:** The field-assigned series name may ultimately change after additional data collection and lab analyses.)

GEOMORPHIC INFORMATION

See the "Geomorphology Section" for complete choice lists. Codes are shown following each example. Conventional "codes" traditionally consist of the entire name; e.g., *mountains*.

PART 1: PHYSIOGRAPHIC LOCATION

Physiographic Division - e.g., *Interior Plains* or *IN*

Physiographic Province - e.g., *Central Lowland* or *CL*

Physiographic Section - e.g., *Wisconsin Driftless Section* or *WDS*

State Physiographic Area (Opt.) - e.g., *Wisconsin Dells*

Local Physiographic/Geographic Name (Opt.) - e.g., *Bob's Ridge*

PART 2: GEOMORPHIC DESCRIPTION

Landscape - e.g., *Foothills* or *FH*

Landform - e.g., *Ridge* or *RI*

Microfeature - e.g., *Mound* or *M*


Anthropogenic Feature - e.g., *Midden* or *H*

PART 3: SURFACE MORPHOMETRY

Elevation - The height of a point on the earth's surface, relative to mean sea level (MSL). Use specific units; e.g., *106 m* or *348 ft*.


Recommended methods: Interpolation from topographic map contours; altimeter reading tied to a known datum. **NOTE:** At present, elevational determination by a sole Global Positioning System (GPS) unit is considered unacceptably inaccurate.

Slope Aspect - The compass direction (in degrees and accounting for declination) that a slope faces, looking downslope; e.g., 287°.




Slope Gradient - The angle of the ground surface (in percent) through the site and in the direction that overland water would flow. Commonly called "slope". Make observations facing downslope to avoid errors associated with some brands of clinometers; e.g., 18%.

Slope Complexity - Describe the relative uniformity (smooth linear or curvilinear = *simple* or *S*) or irregularity (*complex* or *C*) of the ground surface leading downslope through the point of interest; e.g., *simple* or *S*.




Slope Shape - Slope shape is described in two directions: up-and-down slope (perpendicular to the contour), and across slope (along the horizontal contour); e.g., *Linear, Convex* or *LV*.



(S&W, 1996)


Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., *backslope* or *BS*. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS




Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) *nose slope* or *NS*.

Hills	Code	
	PDP	NASIS
interfluve	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	---	BS




(PJS, 1996; adapted from Ruhe, 1975)

Terraces	Code
riser	RI
tread	TR



(S&W, 1996)

Mountains	Code
mountaintop	MT
mountainflank	MF
upper third mountainflank	UT
center third mountainflank	CT
lower third mountainflank	LT
mountainbase	MB



Flat Plains (proposed)	Code
talf	---
rise	---

Microrelief - Small, relative differences in elevation between adjacent areas on the earth's surface; e.g., *micro-high* or *MH*; or *micro-low* or *ML*.

WATER STATUS

DRAINAGE - An estimate of the natural drainage class (i.e., the prevailing wetness conditions) of a soil; e.g., *somewhat poorly drained* or *SP*.

Drainage Class	Code	
	PDP	NASIS
Very Poorly Drained	VP	VP
Poorly Drained	P	PD
Somewhat Poorly Drained	SP	SP
Moderately Well Drained	MW	MW
Well Drained	W	WD
Somewhat Excessively Drained	SE	SE
Excessively Drained	E	ED

The following definitions are the traditional, national criteria for Natural Soil Drainage Classes (Soil Survey Staff, 1993). More specific, regional definitions and criteria vary. (Contact an NRCS State Office for specific, local criteria).

Very Poorly Drained - Water is at or near the soil surface during much of the growing season. Internal free-water is *shallow* and *persistent* or *permanent*. Unless the soil is artificially drained, most mesophytic crops cannot be grown. Commonly, the soil occupies a depression or is level. If rainfall is persistent or high, the soil can be sloping.

Poorly Drained - The soil is wet at shallow depths periodically during the growing season or remains wet for long periods. Internal free-water is *shallow* or *very shallow* and *common* or *persistent*. Unless the soil is artificially drained, most mesophytic crops cannot be grown. The soil, however, is not continuously wet directly below plow depth. The water table is commonly the result of *low* or *very low* saturated hydraulic conductivity class or persistent rainfall, or a combination of both factors.

Somewhat Poorly Drained - The soil is wet at a shallow depth for significant periods during the growing season. Internal free-water is commonly *shallow to moderately deep* and *transitory to permanent*. Unless the soil is artificially drained, the growth of most mesophytic plants is markedly restricted. The soil commonly has a *low* or *very low* saturated hydraulic conductivity class, or a high water table, or receives water from lateral flow, or persistent rainfall, or some combination of these factors.

Moderately Well Drained - Water moves through the soil slowly during some periods of the year. Internal free water commonly is *moderately deep* and may be *transitory* or *permanent*. The soil is wet for only a short time within the rooting depth during the growing season. The soil commonly has a *moderately low*, or lower, saturated hydraulic conductivity class within 1 meter of the surface, or periodically receives high rainfall, or both.

Well Drained - Water moves through the soil readily, but not rapidly. Internal free-water commonly is *deep* or *very deep*; annual duration is not specified. Water is available to plants in humid regions during much of the growing season. Wetness does not inhibit growth of roots for significant periods during most growing seasons. The soil is deep to, or lacks redoximorphic features.

Somewhat Excessively Drained - Water moves through the soil rapidly. Internal free water commonly is *very rare* or *very deep*. The soils are commonly coarse-textured, have high saturated hydraulic conductivity, and lack redoximorphic features.

Excessively Drained - Water moves through the soil very rapidly. Internal free water commonly is *very rare* or *very deep*. The soils are commonly coarse-textured, have very high saturated hydraulic conductivity, and lack redoximorphic features.

FLOODING - Estimate the **Frequency**, **Duration**, and the **Months** flooding is expected; e.g., *rare, brief, Jan. - March*.

Frequency -

Frequency Class	Code		Criteria: estimated, average number of flood events per time span ¹
	PDP	NASIS	
None	NO ²	NO	No reasonable chance (e.g., < 1 time in 500 years)
Very Rare	---	VR	≥ 1 time in 500 years, but < 1 time in 100 years
Rare	RA	RA	1 to 5 times in 100 years
Occasional ³	OC	OC	> 5 to 50 times in 100 years
Frequent ^{3,4}	FR	FR	> 50 times in 100 years
Very Frequent ⁴	---	VF	> 50% of all months in year ²

¹ Flooding Frequency is an estimate of the natural, unmanaged conditions (ignore influence by dams, levees, etc.).

² In PDP, *None* class (< 1 time in 100 years) spans both *None* and *Very Rare* NASIS classes.

³ Historically, *Occasional* and *Frequent* classes could be combined and called *Common*; not recommended.

⁴ *Very Frequent* class takes precedence over *Frequent*, if applicable.

Duration -

Duration Class	Code	Criteria: estimated, average duration per flood event
Extremely Brief	EB	0.1 to < 4 hours
Very Brief	VB	4 to < 48 hours
Brief	BR	2 to < 7 days
Long	LO	7 to < 30 days
Very Long	VL	≥ 30 days

Months - Estimate the beginning and ending month(s) of the year that flooding generally occurs; e.g., *Dec. - Feb.*

PONDING - Estimate or monitor the **Frequency**, **Depth**, and **Duration** of standing water. In PDP, also note the months ponding generally occurs. A complete example is: *occasional, 50 cm, brief, Feb - Apr.*

Frequency -

Frequency Class	Code	Criteria: estimated average # of ponding events per time span
None	NO	< 1 time in 100 years
Rare	RA	1 to 5 times in 100 years
Occasional	OC	> 5 to 50 times in 100 years
Frequent	FR	> 50 times in 100 years

Depth - Estimate the average, representative depth of ponded water at the observation site and specify units; e.g., *1 ft*, or *30 cm*.

Duration -

Duration Class	Code	Criteria: estimated average time per ponding event
Very Brief	VB	< 2 days
Brief	BR	2 to < 7 days
Long	LO	7 to < 30 days
Very Long	VL	≥ 30 days

(SOIL) WATER STATE - (In NASIS and PDP this data element is called **Soil Moisture Status**.) Estimate the water state of the soil at the time of observation; e.g., *wet, satiated*. Soil temperature must be above 0°C. To record conditions with temperatures < 0°C (frozen water); for permanently frozen conditions, see **Texture Modifiers** or **Terms Used in Lieu of Texture** in the "Profile Description Section". **NOTE:** Criteria have changed.

Water State Class	Code	Criteria: tension	Traditional Criteria: tension and field
Dry ¹	D	> 1500 kPa	> 15 bars of tension ³
Moist ¹	M	≤ 1500 to > 1 or > 0.5 kPa ²	Former Usage: > 1/3 to 15 bars of tension (field capacity to wilting point)
Wet	W	< 1.0 or < 0.5 kPa ²	0 - 1/3 bars tension (field capacity or wetter)
Wet, Non-satiated ⁴	WN	> 0.01 and < 1.0 kPa or < 0.5 kPa ² No Free Water	Water films are visible; sand grains and peds glisten, but no free water is present
Wet, Satiated ⁴	WS	< 0.01 kPa, Free Water	Free water easily visible

¹ Additional subclasses of water state can be recognized for *Dry* and *Moist* classes, if desired (Soil Survey Staff, 1993; p. 91).

² Use the 1 kPa limit for all textures, except those coarser than loamy fine sand (Soil Survey Staff, 1993; p. 90).

³ Convention assumes 15 bars of tension as the wilting point for most annual, agricultural row-crops. Caution: Various perennials, shrubs, trees, and other native vegetation have wilting points up to 66 bars tension (= 6600 kPa) or more.

⁴ Satiation vs. Saturation: Satiation implies minor amounts of entrapped air in the smallest pores. True saturation excludes entrapped air. Satiation, for practical purposes, is ≈ saturation. Temporal monitoring of a water table by piezometer or other accepted methods may be needed to verify saturation. Related terms used for classifying soils (i.e., Soil Taxonomy) follow. *Endosaturation* is saturation in all layers to > 200 cm (80 inches). *Episaturation* requires saturated layers that overlie unsaturated layers within the upper 2 m (80 inches). *Anthric saturation*, a variant of episaturation, is saturation due to management-induced flooding (e.g., for rice or cranberry production).

DEPTH TO WATER TABLE - Measure or estimate the depth from the ground surface to the stabilized contact with free-standing water in an open bore-hole or well. Historically, record **Seasonal High Water Table - Kind**, and **Frequency** (duration, beginning month, and days); specify units (e.g., cm, ft). If seasonally variable water is absent at time of observation, it is common practice to estimate prevailing water table conditions based upon soil morphology (e.g., presence of Redoximorphic Features of chroma ≤ 2) in lieu of water-table monitoring data.

NOTE: Within NRCS's PDP and NASIS databases the traditional designation of **Seasonal High Water Table - Kind** and **Frequency** are replaced. In PDP (PEDON), all water table information is recorded in a temporal table. Record **Depth to Stabilized Free Water** and **Date of Observation**. In NASIS, all water table information is replaced by **(Soil) Water State** for each layer at the time of description; e.g., *layer A is moist, layer B is wet, layer C is dry*. For map unit component descriptions, **(Soil) Water State** is recorded (by layer) on a monthly basis, in NASIS.

(Seasonal) High Water Table - Kind - Traditional types of intermittent (e.g., seasonal) high water tables (Soil Survey Staff, 1983). Obsolete in NASIS.

Kind	Code PDP	Criteria:
Apparent	A	Level of stabilized water in a fresh, unlined borehole.
Artesian	---	The final water level within a cased borehole in which the water level rises above an impermeable layer due to a positive hydrostatic head.
Perched	P	A water table that lies above an unsaturated zone. The water table will fall if the borehole is extended.
Ponding ¹	---	Standing water in a closed depression on top of the soil.

¹ A kind of intermittent water table, but not a seasonal high water table (Soil Survey Staff, 1983).

VEGETATION / LAND COVER

EARTH COVER - KIND - Record the dominant land cover at the site; e.g., *intermixed hardwoods & conifers*. (Similar to **Landuse** in PDP.)

Kind †	Code	Kind †	Code
ARTIFICIAL COVER (A) - Nonvegetative cover; due to human activity.			
rural transportation - roads, railroads	ARU	urban and built-up - cities, farmsteads, industry	AUR
BARREN LAND (B) - < 5% vegetative cover naturally or from construction.			
culturally induced - saline seeps, mines, quarries, and oil-waste areas	BCI	other barren - salt flats, mud flats, slickspots, badlands	BOB
permanent snow or ice	BPS	rock	BRK
sand or gravel	BSG		
CROP COVER (C) - includes entire cropping cycle (land prep, crop, or crop residue) for annual or perennial herbaceous plants.			
close-grown crop - wheat, rice, oats, and rye; small grains	CCG	row crop - corn, cotton, soybeans, tomatoes, and other truck crops, tulips	CRC
GRASS / HERBACEOUS (G) - > 50% grass, grass-like (sedge/rushes), or forb cover, mosses, lichens, ferns; non-woody.			
hayland - alfalfa, fescue, brome-grass, timothy	GHL	rangeland, savanna - 10 to 20% tree cover	GRS
marshland - grasses and grass-like plants	GML	rangeland, shrubby - 20 to 50% shrub cover	GRH
pastureland, tame - fescues, brome-grass, timothy, and lespedeza	GPL	rangeland, tundra	GRT
rangeland, grassland; < 10% trees, < 20% shrubs; rangeland used for hayland	GRG	other grass & herbaceous cover	GOH
SHRUB COVER (S) - > 50% shrub or vine canopy cover.			
crop shrubs - filberts, blueberry, ornamental nursery stock	SCS	native shrubs - shrub live oak, mesquite, sage-brush, creosote bush; rangeland > 50% shrub cover	SNS
crop vines - grapes, blackberries, raspberries	SCV	other shrub cover	SOS

TREE COVER (T) - > 25% canopy cover by woody plants, natural or planted.			
conifers - spruce, pine, fir	TCD	swamp - trees, shrubs	TSW
crop, trees - nuts, fruit, nursery, Christmas trees	TCR	tropical - mangrove and royal palms	TTR
hardwoods - oak, hickory, elm, aspen	THW	other tree cover	TOC
intermixed hardwoods & conifers - oak-pine mix	TIM		
WATER (W) - water at the soil surface; includes seasonal frozen water.			

¹ Land Cover Kinds are presented at two levels of detail: Bolded table subheadings are the "NASIS - Level 1" choices (NSSH, Part 622, p. 8; Soil Survey Staff, 1996c). Individual choices under the subheadings are the "NASIS - Level 2" choices.

PLANT SYMBOL - Record the codes (scientific plant name abbreviations) for the major plant species found at the site (NRCS, 1996); e.g., *ANGE* (Andropogon gerardii or big bluestem). **NOTE:** This is the primary plant data element in NASIS.

PLANT COMMON NAME - Record the common names of the major plant species found at the site [NRCS, 1996 (electronic file); SCS, 1989 (hard copy)]; e.g., *cottonwood*, *big bluestem*. This item may be recorded as a secondary data element to augment the **Plant Symbol**. **CAUTION:** Multiple common names exist for some plants; not all common names for a given plant are in the National Plants database.

PLANT SCIENTIFIC NAME - Record the scientific plant name along with or in lieu of common names; e.g., *Acer rubrum* (Red Maple). **[NOTE:** Although used in the past, scientific names of plants (SCS, 1989) are not presently recorded by the NRCS; e.g., PDP has no data element for and does not recognize scientific plant names.] **(NOTE:** NASIS codes for common plant names are derived from the scientific names.)

PARENT MATERIAL

Record the **Kind(s)** of unconsolidated material (regolith) from which the soil is derived. If the soil is derived directly from the underlying bedrock (e.g., granite), identify the **Parent Material** as either *grus*, *saprolite*, or *residuum* and then record the appropriate **Bedrock - Kind** choice. Multiple parent materials, if present, should be denoted; e.g., *loess, over colluvium, over residuum*. Use numerical prefixes in the **Horizon** designations to denote different parent materials (lithologic discontinuities); e.g., *A, BE, 2Bt, 2BC, 3C*.

KIND - e.g., *saprolite, loess, colluvium*.

Kind ¹	Code		Kind ¹	Code	
	PDP	NASIS		PDP	NASIS
EOLIAN DEPOSITS (non-volcanic)					
eolian deposit	E	EOD	loess, calcareous	--	CLO
eolian sands	S	EOS	loess, noncalcareous	--	NLO
loess	W	LOE	parna	--	PAR
GLACIAL DEPOSITS					
drift	D	GDR	till, ablation	--	ATI
glaciofluvial deposit	--	GFD	till, basal	--	BTI
glaciolacustrine deposit	--	GLD	till, flow	--	FTI
glaciomarine deposit	--	GMD	till, lodgement	--	LTI
outwash	G	OTW	till, melt-out	--	MTI
supraglacial debris-flow	--	SGF	till, supraglacial	--	UTI
till	T	TIL	till, supraglacial melt-out	--	PTI
IN-PLACE DEPOSITS (non-transported)					
grus ²	--	GRU	saprolite ²	--	SAP
residuum ²	X	RES			
MASS MOVEMENT DEPOSITS ⁴					
mass movement deposit	--	MMD	mudflow deposit	--	MFD
block glide deposit	--	BGD	rockfall avalanche dep.	--	RAD
colluvium	V	COL	rockfall deposit	--	RFD
creep deposit	--	CRP	rotational landslide dep.	--	RLD
debris avalanche deposit	--	DAD	scree	--	SCR
debris flow deposit	--	DFD	soil fall deposit	--	SFD
debris slide deposit	--	DSD	talus	--	TAL
earthflow deposit	--	EFD	topple deposit	--	TOD
lateral spread deposit	--	LSD			

MISCELLANEOUS DEPOSITS					
crypturbate	--	CRY	mine spoil or earthy fill	F	MSE
diamicton	--	DIM			
ORGANIC DEPOSITS ⁵					
coprogenic materials	--	COM	organic, grassy materials	--	OGM
diatomaceous earth	--	DIE	organic, herbaceous mat.	--	OHM
marl	--	MAR	organic, mossy materials	--	OMM
organic materials	O	ORM	organic, woody materials	--	OWM
VOLCANIC DEPOSITS (unconsolidated; eolian and mass movement)					
ash (< 2 mm)	H	ASH	cinders (2-64 mm)	--	CIN
ash, acidic	--	ASA	lahar (volcaniclastic flow)	--	LAH
ash, andesitic	--	ASN	lapilli (2-64 mm, > 2.0 sg ³)	--	LAP
ash, basaltic	--	ASB	pumice (< 1.0 sg ³)	--	PUM
ash, basic	--	ASC	scoria (> 2.0 sg ³)	--	SCO
ash flow (pyroclastic)	--	ASF	tephra (all ejecta)	--	TEP
bombs (> 64 mm)	--	BOM			
WATER LAID or TRANSPORTED DEPOSITS					
alluvium	A	ALL	marine deposit	M	MAD
backswamp deposit	--	BSD	overbank deposit	--	OBD
beach sand	--	BES	pedisegment	--	PED
estuarine deposit	Z	ESD	slope alluvium	--	SAL
lacustrine deposit	L	LAD	valley side alluvium	--	VSA

¹ Parent material definitions are found in the "Glossary of Landforms and Geologic Terms", NSSH - Part 629 (Soil Survey Staff, 1996c), or the "Glossary of Geology" (Bates et al., 1987).

² Use the most precise term for the in situ material. Residuum is the most generic term.

³ sg = specific gravity = the ratio of a material's density to that of water [weight in air / (weight in air - weight in water)].

⁴ Cruden and Varnes, 1996.

⁵ These generic terms refer to the dominant, origin of the organic materials or deposits from which the organic soil has formed (i.e. parent material) (Soil Survey Staff, 1993). These terms partially overlap with those recognized in Soil Taxonomy (terms which refer primarily to what the organic material presently is); see the "Diagnostic Horizons Table" or "Properties Table".

BEDROCK

Describe the nature of the continuous hard rock underlying the soil. Specify the **Kind**, **Fracture Interval**, **Hardness**, and **Weathering Class**.

KIND - e.g., *limestone*.

Kind	Code ¹		Kind	Code ¹	
	PDP	NASIS		PDP	NASIS
IGNEOUS-INTRUSIVE					
diabase	--	DIA	monzonite	--	MON
diorite	--	DIO	peridotite	--	PER
gabbro	--	GAB	pyroxenite	--	PYX
granite	I4	GRA	syenite	--	SYE
granodiorite	--	GRD	syenodiorite	--	SYD
IGNEOUS-EXTRUSIVE					
aa (<i>lava</i>)	P8	AAL	pahoehoe (<i>lava</i>)	P9	PAH
andesite	I7	AND	pumice (<i>flow, coherent</i>)	E6	PUM
basalt	I6	BAS	rhyolite	--	RHY
dacite	--	DAC	scoria (<i>coherent, mass</i>)	E7	SCO
latite	--	LAT	trachyte	--	TRA
obsidian	--	OBS			
IGNEOUS-PYROCLASTIC					
ignimbrite	--	IGN	tuff breccia	P7	TBR
pyroclastics (<i>coherent</i>)	P0	PYR	volcanic breccia	P4	VBR
tuff	P1	TUF	volcanic breccia, acidic	P5	AVB
tuff, acidic	P2	ATU	volcanic breccia, basic	P6	BVB
tuff, basic	P3	BTU			
METAMORPHIC					
amphibolite	--	AMP	metavolcanics	--	MVO
gneiss	M1	GNE	migmatite	--	MIG
granofels	--	GRF	mylonite	--	MYL
granulite	--	GRL	phyllite	--	PHY
greenstone	--	GRE	schist	M5	SCH
hornfels	--	HOR	serpentinite	M4	SER
marble	L2	MAR	slate	M8	SLA
metaconglomerate	--	MCN	soapstone (<i>talc</i>)	--	SPS
metaquartzite	M9	MQT			

SEDIMENTARY-CLASTICS					
arenite	--	ARE	porcellanite	--	POR
argillite	--	ARG	sandstone	A0	SST
arkose	A2	ARK	sandstone, calcareous	A4	CSS
breccia, non-volcanic (angular fragments)	--	NBR	shale	H0	SHA
claystone	--	CST	shale, acid	--	ASH
conglomerate (rounded fragments)	C0	CON	shale, calcareous	H2	CSH
conglomerate, calcar.	C2	CCN	shale, clayey	H3	YSH
graywacke	--	GRY	siltstone	T0	SIS
mudstone	--	MUD	siltstone, calcareous	T2	CSI
orthoquartzite	--	OQT			
EVAPORITES, ORGANICS, AND PRECIPITATES					
chalk	L1	CHA	limestone, arenaceous	L5	ALS
chert	--	CHE	limestone, argillaceous	L6	RLS
coal	--	COA	limestone, cherty	L7	CLS
dolomite	L3	DOL	limestone, phosphatic	L4	PLS
gypsum	--	GYP	travertine	--	TRV
limestone	L0	LST	tufa	--	TUA
INTERBEDDED					
limestone-sandst.-shale	B1	LSS	sandstone-shale	B5	SSH
limestone-sandstone	B2	LSA	sandstone-siltstone	B6	SSI
limestone-shale	B3	LSH	shale-siltstone	B7	SHS
limestone-siltstone	B4	LSI			

¹ Definitions for bedrock are found in the "Glossary of Landforms and Geologic Terms", NSSH - Part 629 (Soil Survey Staff, 1996c), and in the "Glossary of Geology" (Bates, et al., 1987).

FRACTURE INTERVAL CLASS -

Average Distance Between Fractures	Code
< 10 cm	1
10 to < 45 cm	2
45 to < 100 cm	3
100 to < 200 cm	4
≥ 200 cm	5

HARDNESS (Obsolete -- used in PDP. NASIS uses **Dry Rupture**

Resistance classes (excluding *Loose*) and criteria.) -

Hardness Class	Code	Criteria:
Hard	H	Excavation Difficulty is VH or EH ¹
Soft	S	Paralithic contact criteria ²

¹ *Very Hard (VH)* and *Extremely Hard (EH)* classes from the "Consistence-Excavation Difficulty Table".

² See Keys to Soil Taxonomy (Soil Survey Staff, 1996b).

WEATHERING CLASS - The relative extent to which a bedrock has weathered as compared to a presumed, non-weathered state.

Class	Code	Criteria
Slight	SL	Not Available
Moderate	MO	
Strong	ST	

DEPTH (TO BEDROCK) - Record the depth (cm) from the ground surface to the contact with coherent (continuous) bedrock.

EROSION

Estimate the dominant kind and magnitude of accelerated erosion at the site. Specify the **Kind** and **Degree**.

KIND -

Kind	Code		Criteria ¹
	PDP	NASIS	
Wind	I	I	Deflation by wind
Water	W	---	Removal by running water
Sheet	---	S	Even soil loss, no channels
Rill	---	R	Small channels ²
Gully	---	G	Big channels ³
Tunnel	---	T	Subsurface voids within soil that enlarge by running water (i.e. piping)

¹ Soil Survey Staff, 1993, p82.

² Small, runoff channels that can be obliterated by conventional tillage.

³ Large, runoff channels that cannot be obliterated by conventional tillage.

DEGREE CLASS -

Class	Code	Criteria: Estimated % loss of the original A & E horizons or, the estimated loss of the upper 20 cm (if original combined A & E horizons were < 20 cm thick). ¹
None	0	0%
1	1	> 0 up to 25%
2	2	25 up to 75%
3	3	75 up to 100%
4	4	> 75% & total removal of A

¹ Soil Survey Staff; 1993, pp 86-89.

RUNOFF

SURFACE RUNOFF - Surface runoff (Hortonian flow, overland flow) is the flow of water from an area that occurs over the surface of the soil. Surface runoff differs from internal flow, or throughflow, that results when infiltrated water moves laterally or vertically within a soil, above the watertable. "The Index (of) Surface Runoff Classes" are relative estimates of surface runoff based on slope gradient and saturated hydraulic conductivity (K_{sat}). This index is specific to the following conditions (Soil Survey Staff, 1993).

- The soil surface is assumed to be bare.
- The soil is free of ice.
- Retention of water by ground-surface irregularities is negligible or low.
- Infiltration is assumed to be at the steady ponded infiltration stage.
- Water is added to the soil by precipitation or snowmelt that yields 50 mm in 24-hours with no more than 25 mm in any 1-hour period.
- Antecedent soil water state is assumed to be very moist or wet to: a) the base of the solum; b) a depth of 1/2 m; or c) through the horizon that has the minimum K_{sat} within the top 1 meter; whichever is the least depth.

Use the following table and the above conditions to estimate "The Index (of) Surface Runoff Class" for the site. If seasonal or permanent, internal free-water occurs a depth of ≤ 50 cm (very shallow and shallow internal free-water classes), use a K_{sat} of *Very Low*. If seasonal or permanent, internal free-water is deeper than 50 cm, use the appropriate K_{sat} from the table. In PDP, if estimating runoff from vegetated areas, define and record under **User Defined Property**.

Index (of) Surface Runoff Classes						
Slope Gradient Percent	Saturated Hydraulic Conductivity(K _{sat}) Class ¹					
	Very High	High	Mod. High	Mod. Low	Low	Very Low
	cm / hour					
	≥ 36	3.6	0.36	0.036	0.0036	< 0.0036
		to	to	to	to	
	< 36	< 3.6	< 0.36	< 0.036	< 0.0036	
Concave	N	N	N	N	N	N
< 1	N	N	N	L	M	H
1 to < 5	N	VL	L	M	H	VH
5 to < 10	VL	L	M	H	VH	VH
10 to < 20	VL	L	M	H	VH	VH
≥ 20	L	M	H	VH	VH	VH

¹ This table is based on the minimum K_{sat} occurring within 1/2 m of the soil surface. If the minimum K_{sat} for the soil occurs between 1/2 to 1 m, the runoff estimate should be reduced by one class (e.g., *medium to low*). If the minimum K_{sat} for the soil occurs below 1 meter, use the lowest K_{sat} class that occurs within 1 m of the surface.

Index (of) Surface Runoff Class Names	Code
Negligible	N
Very Low	VL
Low	L
Medium	M
High	H
Very High	VH

SURFACE FRAGMENTS (formerly Surface Stoniness)

Record the amount of surface fragment¹ cover (either as a class or as a numerical percent), as determined by either a "point count" or "line-intercept" method. In NASIS, additional details can be recorded: **Surface Fragment Kind**, (use "Rock Fragment - Kind Table"), **Mean Distance Between Fragments** (edge to edge), **Shape** [FL-flat or NF-nonflat], **Size**, **Roundness** (use classes and criteria found in "Rock Fragment - Roundness Table"), and **Rock Fragment - Rupture Resistance**.

Surface Fragment Class ¹	Code		Criteria: Percentage of surface covered
	Conv ²	NASIS	
Stony or Bouldery	1	%	0.01 to < 0.1
Very Stony or Very Bouldery	2	%	0.1 to < 3
Extremely Stony or Ext. Bouldery	3	%	3 to < 15
Rubbly	4	%	15 to < 50
Very Rubbly	5	%	≥ 50

¹ This data element is also used to record large wood fragments (e.g., tree trunks) on organic soils, if the fragments are a management concern and appear to be relatively permanent.

² Historically called *Surface Stoniness* classes (now *Surface Fragment* classes). Use as a map-unit phase modifier is restricted to stone-sized fragments, or larger (> 250 mm; Soil Survey Staff, 1953).

DIAGNOSTIC HORIZONS or PROPERTIES

Identify the **Kind** and **Upper** and **Lower Depths** of occurrence of Soil Taxonomic diagnostic horizons and properties; e.g., *mollic epipedon*; 0 - 45 cm. Multiple features per horizon can be recorded. (Called **Feature-Kind** in PDP.) In NASIS, **Thickness** (Representative Value (RV), High, Low) can also be recorded.

KIND - (See definitions in current Keys to Soil Taxonomy.)

Kind	Code		Kind	Code	
	PDP	NASIS		PDP	NASIS
EPIPEDONS (Diagnostic Surface Horizons)					
Anthropic	A	AN	Mollic	M	MO
Folistic	--	FO	Ochric	O	OC
Histic	H	HI	Plaggen	P	PL
Melanic	ME	ME	Umbric	U	UM
DIAGNOSTIC SUBSURFACE HORIZONS					
Agric	R	AG	Natric	N	NA
Albic	Q	AL	Ortstein	--	OR
Argillic	T	AR	Oxic	X	OX
Calcic	C	CA	Petrocalcic	E	PE
Cambic	B	CM	Petrogypsic	J	PG
Duripan	Z	DU	Placic	K	PA
Fragipan	F	FR	Salic	Y	SA
Glossic	TO	GL	Sombric	I	SO

Gypsic	G	GY	Spodic	S	SP
Kandic	KA	KA	Sulfuric	V	SU
DIAGNOSTIC PROPERTIES - MINERAL SOILS					
Abrupt textural change	AC	AC	Lamella / Lamellae	--	LA
Albic material	--	AM	Lithic contact	L	LC
Albic material, interfinger	IF	AI	Paralithic contact	W	PC
Andic soil properties	AN	AP	Paralithic material	--	PM
Aquic conditions	---	AQ	Permafrost	PF	PA
Carbonates, secondary ¹	LI	SC	Petroferric contact	PC	TC
Densic contact	--	DC	Plinthite	PL	PI
Densic material	--	DM	Slickensides	SL	SS
Durinodes	D	DN	Sulfidic material	SU	SM
Fragic soil properties	--	FP			
DIAGNOSTIC PROPERTIES - ORGANIC SOILS					
Fibric soil materials	FI	FM	Limnic materials	LM	LM
Hemic soil materials	HE	HM	Coprogeous earth	CO	CO
Humultuvic materials	HU	UM	Diatomaceous earth	DI	DI
Sapric soil materials	SA	RM	Marl	MA	MA

¹ Secondary carbonates, replaces "soft, powdery lime". **NOTE:** Gilgai (GI in PDP) is no longer a diagnostic feature in Soil Taxonomy.

DEPTH - Document the zone of occurrence for a diagnostic horizon or property, as observed, by recording the upper and lower depth and specify units; e.g., 22 - 39 cm. Record **Top Depth** and **Bottom Depth**.

REFERENCES

[References for this "Site Description Section" are combined with those at the end of the "Profile / Pedon Description Section" on page 2-68.]

PROFILE / PEDON DESCRIPTION

Compiled by: D.A. Wysocki, P.J. Schoeneberger, E.C. Benham, NRCS,
Lincoln, NE; W. D. Broderson, NRCS, Salt Lake City, UT.

OBSERVATION METHOD

For each layer, indicate the type and relative extent of the exposure upon which the primary observations are made. (Examples of common sampling devices are included in the "Field Sampling Section".) Describe **Kind** and **Relative Size**.

KIND -

Kind	Code	Criteria: Types (common size or ranges)
"Disturbed" Samples		
bucket auger	BA	e.g., open, closed, sand, mud buckets (5-12 cm diam.)
screw auger	SA	e.g., external thread hand augers, power (flight) auger (2-30 cm diam.)
"Undisturbed" Samples		
push tube	PT	e.g., hand held, hydraulic, hollow stem (2-10 cm diam.)
shovel "slice" ¹	SS	e.g., undisturbed block extracted with a shovel (sharpshooter: 20 x 40 cm)
WALL / FLOOR - "Undisturbed" Area or Exposure		
small pit	SP	e.g., hand dug (< 1 m x 2 m)
trench	TR	e.g., backhoe, pipeline (> 1 m x 2 m)
beveled cut	BC	e.g., roadcuts graded to < 60% slope
cut	CU	e.g., roadcut, streambank, medium-sized borrow pit wall > 60% slope (e.g., > 4 m, < 33 m)
large open pit or quarry	LP	large borrow pits, large or irregular banks (e.g., > 33 m)

¹ Field method used for hydric soil investigations.

RELATIVE SIZE - Record the approximate size of the exposure observed. Use cm for "Drill Cores" and m for "Wall/Floor" observations; e.g., *bucket auger, 3 cm, trench wall, 3 m.* (**NOTE:** Common size range for each method is indicated in the "Criteria" column of the "Observation Method - Kind Table". These dimensions are not intended to be restrictive or precise; merely approximate.)

TAXONOMIC CLASSIFICATION

After completely describing the soil, classify the pedon as completely as possible (to the lowest taxonomic level). See the most current version of Keys to Soil Taxonomy or NASIS for a complete choice list; e.g., *fine, mixed, active, mesic Typic Haplohumult*.

HORIZON NOMENCLATURE

Use capital letters to identify master horizons; e.g., *A, B*. Use suffixes (lowercase letters) to denote additional horizon characteristics or features; e.g., *Ap, Btk*. [For more detailed criteria, see the "Soil Taxonomy Section"; for complete definitions see Keys to Soil Taxonomy (Soil Survey Staff, 1996).] Label a horizon only after all morphology is recorded.

MASTER, TRANSITIONAL AND COMMON HORIZON COMBINATIONS ¹

Horizon	Criteria
O	Predominantly organic matter (litter & humus)
A	Mineral, organic matter (humus) accumulation, loss of Fe, Al, clay
AB (or AE)	Dominantly A horizon characteristics but also contains some characteristics of the B (or E) horizon
A/B (or A/E or A/C)	Discrete, intermingled bodies of A and B (or E or C) material; majority of horizon is A material
AC	Dominantly A horizon characteristics but also contains some characteristics of C horizon
E	Mineral, loss of Si, Fe, Al, clay, or organic matter
EA (or EB)	Dominantly E horizon characteristics but also contains some attributes of the A (or B) horizon
E/A	Discrete, intermingled bodies of E and A horizon material; majority of horizon is E material
E and Bt	Thin lamellae (Bt) within a dominantly E horizon
BA (or BE)	Dominantly B characteristics but also contains some attributes of A (or E) horizon
B/A (or B/E)	Discrete, intermingled bodies of B and A (E) material; majority of horizon is B material
B	Subsurface accumulation of clay, Fe, Al, Si, humus, CaCO ₃ , CaSO ₄ ; or loss of CaCO ₃ ; or accumulation of sesquioxides; or subsurface soil structure
BC	Dominantly B horizon characteristics but also contains some characteristics of the C horizon

B/C	Discrete, intermingled bodies of B and C material; majority of horizon is B material
CB (or CA)	Dominantly C horizon characteristics but also contains some characteristics of the B (or A) horizon
C/B (or C/A)	Discrete, intermingled bodies of C and B (or A) material; majority of horizon is C material
C	Little or no pedogenic alteration, unconsolidated earthy material, soft bedrock
R	Hard, continuous bedrock
W	A layer of liquid water (W) or permanently frozen water (Wf) within the soil (excludes water/ice above soil) ²

¹ Refer to the "Soil Taxonomy Section" for older horizon nomenclature.

² NRCS Soil Classification Staff, 1997; personal communication.

HORIZON SUFFIXES - Historically referred to as "Horizon Subscripts", and more recently as "Subordinate Distinctions" ¹. (Historical codes and conversions are shown in the "Soil Taxonomy Section".)

Horizon Suffix ²	Criteria
a	Highly decomposed organic matter
b	Buried genetic horizon (not used with C horizons)
c	Concretions or nodules
d	Densic layer (physically root restrictive)
e	Moderately decomposed organic matter
f	Permanently frozen soil or ice (permafrost); continuous, subsurface ice; not seasonal
ff	Permanently frozen soil ("Dry" permafrost); no continuous ice; not seasonal ³
g	Strong gley
h	Illuvial organic matter accumulation
i	Slightly decomposed organic matter
j	Jarosite accumulation ³
jj	Evidence of cryoturbation ³
k	Pedogenic carbonate accumulation
m	Strong cementation (pedogenic, massive)
n	Pedogenic, exchangeable sodium accumulation
o	Residual sesquioxide accumulation (pedogenic)

p	Plow layer or other artificial disturbance
q	Secondary (pedogenic) silica accumulation
r	Weathered or soft bedrock
s	Illuvial sesquioxide accumulation
ss	Slickensides
t	Illuvial accumulation of silicate clay
v	Plinthite
w	Weak color or structure within B (used only with B)
x	Fragipan characteristics
y	Pedogenic accumulation of gypsum
z	Pedogenic accumulation of salt more soluble than gypsum

¹ Keys to Soil Taxonomy, 6th Edition, 1994.

² Keys to Soil Taxonomy, 7th Edition, 1996.

³ NRCS Soil Classification Staff, 1997; personal communication.

OTHER HORIZON MODIFIERS -

Numerical Prefixes (2, 3, etc.) - Used to denote lithologic discontinuities. By convention, 1 is understood but is not shown; e.g., *A, E, B11, 2B12, 2BC, 3C1, 3C2*.

Numerical Suffixes - Used to denote subdivisions within a master horizon; e.g., *A1, A2, E, B11, B12, B13, Bs1, Bs2*.

The Prime (') -Used to indicate the second occurrence of an identical horizon descriptor(s) in a profile or pedon; e.g., *A, E, Bt, E' Btx, C*. The prime does not indicate either buried horizons (which are denoted by a lower case "b"; e.g., *Btb*), or lithologic discontinuities (denoted by numerical prefixes). Double and triple primes are used to denote subsequent occurrences of horizon descriptors in a pedon; e.g., *A, E, Bt, E', Btx, E'', Cd*.

DIAGNOSTIC HORIZONS - See the "Diagnostic Horizons Table" or "Properties Table", in the "Site Description Section".

HORIZON DEPTH - Record the depths of both the upper and lower boundary for each horizon; specify units (centimeters preferred); e.g., *15-24 cm*. Begin (zero datum) at the ground surface¹, which is not necessarily the mineral surface. (**NOTE:** Prior to 1993, the zero datum was at the top of the mineral surface, except for thick organic layers such as a peat or muck.

Organic horizons were recorded as above and mineral horizons recorded as below, relative to the mineral surface.) Example:

Zero Datum for the same horizons

At Present: Oe 0 - 5 cm, A 5 - 15 cm, E 15 - 24 cm
 Before 1993: Oe 5 - 0 cm, A 0 - 10 cm, E 10 - 19 cm

- ¹ Conventionally, the "soil surface" is considered to be the top boundary of the first layer that can support plant / root growth. This equates to:
- a) (for bare mineral soil) the air/fine earth interface;
 - b) (for vegetated mineral soil) the upper boundary of the first layer that can support root growth;
 - c) (for organic mantles) the same as b) but excludes freshly fallen plant litter, and includes litter that has compacted and begun to decompose; e.g., Oi horizon;
 - d) (for submerged soil) the same as b) but refers to the water/soil contact and extends out from shore to the limit of emergent, rooted plants;
 - e) (for rock mulches; e.g., desert pavement, scree) the same as a) unless the areal percentage of surface rock coverage is greater than 80%, the top of the soil is the mean height of the top of the rocks.

HORIZON THICKNESS - Record the average thickness and range in thickness of horizon; e.g., 15 cm (12 - 21 cm).


HORIZON BOUNDARY - Record **Distinctness** and **Topography** of horizon boundary. Distinctness is the distance through which one horizon grades into another. Topography is the lateral undulation and continuity of the boundary between horizons. A complete example is: *clear, wavy*, or *C,W*.

Distinctness -

Distinctness Class	Code		Criteria: thickness
	PDP	NASIS	
Very Abrupt	---	V	< 0.5 cm
Abrupt	A	A	0.5 to < 2 cm
Clear	C	C	2 to < 5 cm
Gradual	G	G	5 to < 15 cm
Diffuse	D	D	≥ 15 cm


Topography - Cross-sectional shape of the contact between horizons.

Topography	Code	Criteria
Smooth	S	Planar with few or no irregularities
Wavy	W	Width of undulation is > than depth
Irregular	I	Depth of undulation is > than width
Broken	B	Discontinuous horizons; discrete but intermingled, or irregular pockets



SOIL COLOR

DECISION FLOWCHART FOR DESCRIBING SOIL COLORS - Use the following chart to decide how and with which data elements the color patterns of a soil or soil feature should be described.



NOTE: *Reduced Matrix* color is described as a *Matrix Color* and in the associated "(Soil Color) - Location or Condition Described Table".

(SOIL) MATRIX COLOR - Record **Color(s)**, **(Soil Color) Moisture State**, **Location or Condition**. (In PDP, also record **Percent of Horizon**, if more than one matrix color is described.)

(Soil) Matrix Color - (Soil) Color - Use Munsell® notation (Hue, Value, Chroma); e.g., *10YR 3/2*. Neutral Gley colors are written as chroma of zero (0); e.g., *N 4/0*. Other gley colors use appropriate notation (see Munsell® Gley pages; e.g., *5GY 6/1*). For narrative descriptions (Soil Survey Reports, Official Series Descriptions) both the verbal name and the Munsell® notation are given; e.g., *dark brown, 10YR 3/3*.

(Soil) Matrix Color - Moisture State - Record the moisture condition of the soil described; e.g., *moist*. (Not to be confused with Soil Water State.)

Moisture State	Code
Dry	D
Moist	M

(Soil) Matrix Color - Location or Condition - Record pertinent circumstances of the color described.

Color Location or Condition	Code	
	PDP	NASIS
COLOR LOCATION		
Interior (<i>within ped</i>)	1	IN
Exterior (<i>ped surface</i>)	2	EX
COLOR, MECHANICAL CONDITION		
Broken Face	8	BF
Crushed	3	CR
Rubbed (<i>used only with Organic Matter</i>)	9	RU
COLOR, REDOXIMORPHIC CONDITION		
Oxidized ¹	5	OX
Reduced ²	---	RE
COLOR, INTRICATE MULTICOLORED PATTERN		
Variegated ³	---	VA

¹ Soil that is reduced in situ, but oxidizes (changes color) after extraction and exposure to air. A mineral example is vivianite.

NOTE: Not used for soil that's normally oxidized in-place. For indicators of reduction see **Redoximorphic Features**.


² Color immediately after extraction from a reduced environment, prior to oxidation; e.g., *FeS*. Also used to record **Reduced Matrix**.

³ Color pattern is too intricate (banded or patchy) with numerous, diverse colors to credibly identify dominant matrix colors.

MOTTLES - Describe mottles (areas of color that differ from the matrix color). These colors are commonly lithochromic or lithomorphic (attributes retained from the geologic source rather than from pedogenesis; e.g., gray shale). Mottles exclude: Redoximorphic Features (RMF) and Ped & Void Surface Features; e.g., clay films. Record **Quantity Class** (in NASIS/PDP, estimate a numerical value "Percent of Horizon Area Covered"), **Size**, **Contrast**, **Color**, and **Moisture State** (D or M). **Shape** is an optional descriptor (use the "Concentrations - Shape Table"). A complete example is: *few, medium, distinct, reddish yellow, moist, irregular mottles* or *f, 2, d, 7.5 YR 7/8, m, z, mottles*.


Mottles - Quantity (Percent of Area Covered) -

Quantity Class	Code		Criteria: range in percent
	Conv	NASIS	
Few	f	%	< 2% of surface area
Common	c	%	2 to < 20% of surface area
Many	m	%	≥ 20% of surface area



Mottles - Size - Record mottle size class. Use length if it's greater than 2 times the width; use width if the length is less than two times the width. Length is the greater of the two dimensions. (New size classes to be consistent with the new RMF size classes.)

Size Class	Code	Criteria
Fine	1	< 2 mm
Medium	2	2 to < 5 mm
Coarse	3	5 to < 20 mm
Very Coarse	4	20 to < 76 mm
Extremely Coarse	5	≥ 76 mm



Mottles - Contrast - Record the color difference between the mottle and the dominant matrix color. Use this table or the following chart to express the difference.

Contrast Class	Code	Difference in Color Between Matrix and Mottle		
		Hue ¹	Value	Chroma
Faint ²	F	same page	0 to ≤ 2	and ≤ 1
Distinct	D	same page	> 2 to < 4	and < 4
		1 page	< 4	or > 1 to < 4
Prominent	P	same page	≤ 2	and ≤ 1
		1 page	≥ 4	or ≥ 4
		≥ 2 pages	> 2	or > 1
			≥ 0	or ≥ 0

¹ One Munsell® Color Book page = 2.5 hue units. Table contents compiled from material in or intended by the Soil Survey Manual (Soil Survey Staff, 1993).

² *Faint* also includes mottles or RMFs that are similar in color to the matrix that have both low (e.g., ≤ 3) value and chroma, and differ by up to 2.5 units (one page) of hue.

Mottles - Color - Use standard Munsell® notation of hue, value, chroma; e.g., 5 YR 4/4 (for reddish brown).

Mottles - Moisture State - Record the moisture condition of the mottle (not to be confused with soil water state); e.g., *moist*.

Moisture State	Code
Dry	D
Moist	M

Mottles - Shape (optional) - Use "Concentrations - Shape Table"; e.g., *irregular*.

NOTE: In PDP, **Location (optional)**, and **Hardness (optional)** can be described. Use the choices in the appropriate "Redoximorphic Features Table".

Contrast of Soil Mottles (For Use with Munsell Color Book)




Chart Directions:

- Select the appropriate "Difference in Hue" line ("None" means "same page").
- Record greatest contrast of **value** or **chroma** at **hue** line intercept (faint, distinct, or prominent).

REDOXIMORPHIC FEATURES (RMF) DISCUSSION

Redoximorphic Features (RMF) are a color pattern in a soil due to loss (depletion) or gain (concentration) of pigment compared to the matrix color, formed by oxidation / reduction of Fe and/or Mn coupled with their removal, translocation, or accrual; or a soil matrix color controlled by the presence of Fe⁺². The composition and process of formation for a soil color or color pattern must be known or inferred before describing it as a RMF. Because of this inference, RMF are described separately from other mottles, concentrations; e.g., *salts*; or compositional features; e.g., *clay films*. RMF generally occur in one or more of these settings:

- a. In the soil matrix, unrelated to surfaces of peds or pores.
- b. On or beneath the surfaces of peds.
- c. As filled pores, linings of pores, or beneath the surfaces of pores.

RMFs include the following:

1. **Redox Concentrations** - Localized zones of enhanced pigmentation due to an accrual of, or a phase change in, the Fe-Mn minerals; or are physical accumulations of Fe-Mn minerals. **NOTE:** Iron concentrations may be either Fe⁺³ or Fe⁺². Types of redox concentrations are:
 - a. Masses - Noncemented bodies of enhanced pigmentation that have a redder or blacker color than the adjacent matrix.
 - b. Nodules or Concretions - Cemented bodies of Fe-Mn oxides.
2. **Redox Depletions** - Localized zones of "decreased" pigmentation that are grayer, lighter, or less red than the adjacent matrix. Redox depletions include, but are not limited to, what were previously called "low chroma mottles" (chroma ≤ 2). Depletions with chroma ≤ 2 are used to define aquic conditions in Soil Taxonomy and are used extensively in the field to infer occurrence and depth of saturation in soils. Types of redox depletions are:
 - a. Iron Depletions - Localized zones that have one or more of the following: a yellower, greener, or bluer hue; a higher value; or a lower chroma than the matrix color. Color value is normally ≥ 4 . Loss of pigmentation results from the loss of Fe and/or Mn. Clay content equals that in the matrix.
 - b. Clay Depletions - Localized zones that have either a yellower hue, a higher value, or a lower chroma than the matrix color. Color value is normally ≥ 4 . Loss of pigmentation results from a loss of Fe and/or Mn and clay. Silt coats or skeletons commonly form as depletions but can be non-redox concentrations, if deposited as flow material in pores or along faces of peds.

3. **Reduced Matrix** - A soil horizon that has an in situ matrix chroma ≤ 2 due to the presence of Fe^{+2} . Color of a sample becomes redder or brighter (oxidizes) when exposed to air. The color change usually occurs within 30 minutes. A 0.2% solution of α, α' -dipyridyl dissolved in 1N ammonium acetate (NH_4OAc) pH 7 can verify the presence of Fe^{+2} in the field (Childs, 1981).

NOTE: Use of RMF alters the traditional sequence for describing soil color (see the "Decision Flowchart for Describing Colors for Soil Matrix and Soil Features"). RMF are described separately from other color variations or concentrations. Mottles (color variations not due to loss or accrual of Fe-Mn oxides; e.g., variegated weathered rock) are still described under **Soil Color**. A Reduced Matrix is recorded as a RMF and as "reduced" in **Soil Color - Location or Condition Described**.

REDOXIMORPHIC FEATURES

Record **Kind, Quantity** (percent of area covered), **Size, Contrast, Color, Moisture State, Shape, Location, Hardness, and Boundary**. A complete example is: *common, medium, prominent, black Iron-Manganese nodules, moist, spherical in the matrix, hard, sharp or c, 2, p, 5 YR 2.5/1, FMM, M, o, h, s.*

REDOXIMORPHIC FEATURES - KIND -

Kind	Code		Kind	Code	
	PDP	NASIS		PDP	NASIS
REDUCED MATRIX (chroma ≤ 2 primarily from Fe^{+2})					
Reduced Matrix	---	RMX			
REDOX DEPLETIONS (loss of pigment or material)					
Clay Depletions	A3	CLD	Iron Depletions (includes depletion halo)	F5	FED
Chroma > 2	---	HCD	Chroma > 2	---	HFD
REDOX CONCENTRATIONS (accumulated pigment, material)					
Masses ¹ (noncemented)					
Iron (Fe^{+3}) ^{3,4,5}	F2	F3M	Iron-Manganese ^{3,4,5}	M2	FMM
Iron (Fe^{+2}) ²	---	F2M	Manganese ^{4,5}	M8	MNM
Nodules ¹ (cemented; no layers, crystals not visible at 10X)					
Ironstone	F4	FSN	Iron-Manganese ⁴	M5	FMN
Plinthite	F1	PLN			
Concretions ¹ (cemented; distinct layers, crystals not visible)					
Iron-Manganese ⁴				M3	FMC

Surface Coats / Films or Hypocoats		
Manganese (<i>mangans: black, very thin, exterior films</i>)	M ⁶	MNF
Ferriargillans (<i>Fe³ stained clay film</i>)	I ⁶	FEF


- ¹ See discussion under **Concentrations** for definitions.
- ² A concentration of reduced iron Fe²⁺; e.g., *FeS*.
- ³ A concentration of oxidized iron Fe³⁺; e.g., *hematite*, (formerly described as *reddish mottles*).
- ⁴ Iron and Mn commonly occur in combination and field identification of distinct phases is difficult. Use *Mn masses* only for those that are at least *Slightly Effervescent* with H₂O₂. Describe nodules and concretions as *Iron-Manganese* unless colors are unambiguous.
- ⁵ Suggested, color guidelines for field description of Fe vs. Mn Masses:

Color of Concentration Value		Dominant Composition
Value	Chroma	
≤ 2	≤ 2	Mn
> 2 & ≤ 4	> 2 & ≤ 4	Fe & Mn
> 4	> 4	Fe

- ⁶ In PDP, these features (codes) are recorded under **Coat - Kind**.

REDOXIMORPHIC FEATURES - QUANTITY (Percent of Area Covered) -

Class	Code		Criteria: Percent of Surface Area Covered
	Conv.	NASIS	
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20



REDOXIMORPHIC FEATURES - SIZE - See size class graphic under either **Mottles** or **Concentrations**.

Size Class	Code	Criteria
Fine	1	< 2 mm
Medium	2	2 to < 5 mm
Coarse	3	5 to < 20 mm
Very Coarse	4	20 to < 76 mm
Extremely Coarse	5	≥ 76 mm

REDOXIMORPHIC FEATURES - CONTRAST - Use "Mottle - Contrast Table" or "Mottle - Contrasts Chart"; e.g., *Prominent* or *p*.

REDOXIMORPHIC FEATURES - COLOR - Use standard Munsell® notation from the "Soil Color Section"; e.g., *light brownish gray* or *2.5Y 6/2*.

REDOXIMORPHIC FEATURES - MOISTURE STATE - Describe the moisture condition of the Redoximorphic Feature (use "Soil Color - Moisture State Table"); e.g., *Moist (M)* or *Dry (D)*.

REDOXIMORPHIC FEATURES - SHAPE - Describe the shape of the redoximorphic feature (use "Concentrations - Shape Table"); e.g., *Spherical (O)*.

REDOXIMORPHIC FEATURES - LOCATION - Describe the location(s) of the Redoximorphic Feature within the horizon (use "Concentrations - Location Table"); e.g., *In the matrix (R1)*.

REDOXIMORPHIC FEATURES - HARDNESS - Describe the hardness of the Redoximorphic Feature (use cementation classes from the "Consistence - Rupture Resistance for Blocks / Peds / Clods Table"); e.g., *Strongly Cemented (ST)*.

REDOXIMORPHIC FEATURES - BOUNDARY - The gradation between the Redoximorphic Feature and the adjacent matrix (use "Concentrations - Boundary Table"); e.g., *Sharp (S)*.


CONCENTRATIONS DISCUSSION

Concentrations are soil features that form by accumulation of material during pedogenesis. Dominant processes involved are chemical dissolution/precipitation; oxidation and reduction; and physical and/or biological removal, transport, and accrual. Types of concentrations (modified from Soil Survey Staff, 1993) include the following:

1. **Finely Disseminated Materials** are physically small precipitates (e.g., salts, carbonates) dispersed throughout the matrix of a horizon. The materials cannot be readily seen (10X lens), but can be detected by a chemical reaction (e.g., effervescence of CaCO_3 by HCl) or other proxy indicators.
2. **Masses** are noncemented ("Rupture Resistance Cementation Class" of *Extremely Weakly Cemented* or less) bodies of accumulation of various shapes that cannot be removed as discrete units, and do not have a crystal structure that is readily discernible in the field (10X hand lens). This includes finely crystalline salts and Redox Concentrations that do not qualify as nodules or concretions.
3. **Nodules** are cemented (*Very Weakly Cemented* or greater) bodies of various shapes (commonly spherical or tubular) that can be removed as discrete units from soil. Crystal structure is not discernible with a 10X hand lens.
4. **Concretions** are cemented bodies (*Very Weakly Cemented* or greater) similar to nodules, except for the presence of visible, concentric layers of material around a point, line, or plane. The terms "nodule" and "concretion" are not interchangeable.
5. **Crystals** are macro-crystalline forms of relatively soluble salts (e.g., halite, gypsum, carbonates) that form in situ by precipitation from soil solution. The crystalline shape and structure is readily discernible in the field with a 10X hand lens.
6. **Biological Concentrations** are discrete bodies accumulated by a biological process (e.g., fecal pellets), or pseudomorphs of biota or biological processes (e.g., insect casts).

General conventions for documenting various kinds of **Concentrations**:

Type of Distribution	Documentation	Examples
Finely Disseminated (not visible)	Horizon Suffix	Carbonates (Bk) Salts (Bz, Bn)
Masses, Nodules, Concretions, Crystals, Biological Features	Redoximorphic Features, or Concentrations	Mn nodules Fe concretions Insect casts
Continuous Cementation	Terms in Lieu of Texture	Duripan Petrocalcic



CONCENTRATIONS

Record **Kind, Quantity** (percent of area covered), **Size, Contrast, Color, Moisture State, Shape, Location, Hardness, and Boundary**. A complete example is: *many, fine, prominent, white, moist, cylindrical, carbonate nodules in the matrix, moderately cemented, clear or m, 1, p, 10YR 8/1, M, c, CAN, MAT, M, c.*

CONCENTRATIONS - KIND - Identify the composition and the physical state of the concentration in the soil. **NOTE:** Table sub-headings (e.g., *Masses*) are a guide to various physical states of materials. Materials with similar or identical chemical compositions may occur in multiple physical states (under several sub-headings); e.g., *salt masses* and *salt crystals*.

CONCENTRATIONS (NON-REDOX) (accumulations of material)					
Kind	Code		Kind	Code	
	PDP	NASIS		PDP	NASIS
MASSES (noncemented; crystals not visible with 10X hand lens)					
Barite (<i>BaSO₄</i>)	B2	BAM	Gypsum (<i>CaSO₄·2H₂O</i>)	G2	GYM
Carbonates (<i>Ca, Mg, NaCO₃</i>)	K2	CAM	Salt (<i>NaCl, Na-Mg Sulfates</i>)	H2	SAM
Clay Bodies	A2	CBM	Silica	S2	SIM
Gypsum (Nests)	G3	GNM			
NODULES (cemented; non-crystalline at 10X, no layers)					
Carbonates ¹	C4	CAN	Gibbsite (<i>Al₂O₃</i>)	E4	GBN
Durinodes (<i>SiO₂</i>)	S4	DNN	Opal	S1	OPN
CONCRETIONS (cemented; non-crystalline at 10X, distinct layers)					
Carbonates ¹	C3	CAC	Silica	S3	SIC
Gibbsite	E3	GBC	Titanium Oxide	---	TIC
CRYSTALS (crystals visible with 10X hand lens)					
Barite (<i>BaSO₄</i>)	B1	BAX	Gypsum (<i>CaSO₄·2H₂O</i>)	G1	GYX
Calcite (<i>CaCO₃</i>)	C1	CAX	Salt (<i>NaCl, Na-Mg Sulfates</i>)	H1	SAX
BIOLOGICAL CONCENTRATIONS (byproducts or pseudomorphs)					
Fecal Pellets	---	FPB	Shell Fragments (<i>terrestrial or aquatic</i>)	---	SFB
Insect Casts ²	T3	ICB	Sponge Spicules ³	---	SSB
Plant Phytoliths ³ (<i>plant opal</i>)	---	PPB	Worm Casts ²	T2	WCB
Root Sheaths	---	RSB			


¹ Also known as loess kinchen, loess puppies, etc.

² Worm casts are ovoid, fecal pellets excreted by earthworms. Insect casts are cemented (e.g., $CaCO_3$) molds of insect bodies or burrows.

³ May require magnification > 10X to be observed.

CONCENTRATIONS - QUANTITY (PERCENT OF AREA COVERED) -

Class	Code		Criteria: % of Surface Area Covered
	Conv.	NASIS	
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20




CONCENTRATIONS - SIZE - Use "RMF's" and "Mottle Size Classes". (See graphic on next page.)

Size Class	Code	Criteria
Fine	1	< 2 mm
Medium	2	2 to < 5 mm
Coarse	3	5 to < 20 mm
Very Coarse	4	20 to < 76 mm
Extremely Coarse	5	≥ 76 mm

CONCENTRATIONS - CONTRAST - Use "Mottle - Contrast Table" or "Mottle - Contrast Chart"; e.g., *distinct*.

CONCENTRATIONS - COLOR - Use standard Munsell® notation; e.g., *7.5 YR 8/1*.

CONCENTRATIONS - MOISTURE STATE - Use "Soil Color - Moisture State Table"; e.g., *Moist (M)* or *Dry (D)*.




CONCENTRATIONS - SHAPE -

Shape	Code		Criteria
	PDP	NASIS	
Cylindrical	C	C	tubular & elongated bodies; e.g., filled worm holes and insect burrows
Dendritic	D	D	tubular, elongated, branched bodies; e.g., pipestems (root pseudomorphs)
Irregular	Z	I	bodies of non-repeating spacing or shape
Platy	P	P	relatively thin, tabular sheets, lenses; e.g., lamellae
Reticulate	---	R	crudely interlocking bodies with similar spacing; e.g., plinthite
Spherical ¹	O	S	well-rounded to crudely spherical bodies; e.g., Fe / Mn "shot"
Threads	T	T	thin (e.g., < 1 mm diam.) elongated filaments; generally not dendritic; e.g., very fine CaCO ₃ stringers

¹ Called *Rounded* in PDP.

Examples of Mottles, Concentrations, and RMF Shapes



CONCENTRATIONS - LOCATION - (Also used for **Redoximorphic Features**.) Describe the location(s) of the concentration (or depletion for RMF's) within the horizon. Historically called **Concentrations - Distribution**.

Location	Code	
	PDP	NASIS
MATRIX (in soil matrix; not associated with peds or pores)		
In the matrix (not associated with peds/pores)	---	MAT
In matrix around depletions	---	MAD
In matrix around concentrations	---	MAC
Throughout (e.g., finely disseminated carbonates)	T	TOT
PEDS (on or associated with faces of peds)		
Between peds	P	BPF
Infused into the matrix along faces of peds (hypocoats) ¹	---	MPF
On faces of peds (all orientations)	---	APF
On horizontal faces of peds	---	HPF
On vertical faces of peds	---	VPF
PORES (in pores, or associated with surfaces along pores)		
On surfaces along pores	---	SPO
Infused into the matrix adjacent to pores (hypocoats) ¹	---	MPO
Lining pores ¹	---	LPO
OTHER		
In cracks	C	CRK
Top of horizon	M	TOH
Around rock fragments	S	ARF
On bottom of rock fragments (e.g., pendants)	---	BRF

¹ See illustration under **Ped and Void Surface Features - Kind**.

CONCENTRATIONS - HARDNESS - Describe the relative force required to crush the concentration body. Record the appropriate **Rupture Resistance - Cementation Class** (see "Rupture Resistance Table"); e.g., *Moderately Cemented* (exclude the *Non-Cemented* class). **NOTE:** PDP doesn't recognize the *Moderately Hard* class, dry nor moist (= *Very Weakly Cemented* class).

CONCENTRATIONS - BOUNDARY - The gradation between feature and matrix.

Class	Code	Criteria
Sharp	S	Color changes in < 0.1 mm; change is abrupt even under a 10X hand lens.
Clear	C	Color changes within 0.1 to < 2 mm; gradation is visible without 10X lens.
Diffuse	D	Color changes in ≥ 2 mm; gradation is easily visible without 10X hand lens.

PED & VOID SURFACE FEATURES

These features are coats/films, hypocoats, or stress features formed by translocation and deposition, or shrink-swell processes on or along surfaces. Describe **Kind**, **Amount Class** (percent in NASIS and PDP), **Distinctness**, **Location**, and **Color** (dry or moist). An example is: *many, faint, brown 10YR 4/6 (Moist), clay films on all faces of peds or m, f, 10YR 4/6 (M), CLF, PF.*

PED & VOID SURFACE FEATURES - KIND (non-redoximorphic) -



Kind	Code		Field Criteria
	PDP	NASIS	
COATS, FILMS (exterior, adhered to surface)			
Carbonate Coats	K	CAF	off-white, effervescent with HCl
Silica (silans, opal)	--	SIF	off-white, noneffervescent with HCl
Clay Films (Argillans)	T	CLF	waxy, exterior coats
Clay Bridging	D	BRF	"wax" between grains
Ferriargillans Described as RMF - Kind		see RMFs	Fe ⁺³ stained clay film.
Gibbsite Coats (sesquan)	G	GBF	Al(OH) ₃ , off-white, noneffervescent with HCl
Manganese (mangans) Described as RMF - Kind		see RMFs	black, thin films effervescent with H ₂ O ₂
Organic Stains	--	OSF	dark organic films
Organoargillans	O	OAF	dark, organic stained clay films
Sand Coats	Z	SNF	separate grains visible with 10X
Silt Coats ¹	R	SLF	separate grains not visible at 10X
Skeletans ² (sand or silt)	S	SKF	clean sand or silt grains as coats
Skeletans on argillans	A	SAF	clean sand or silt over clay coats
HYPOCOATS ³ (A stain infused beneath a surface.)			
STRESS FEATURES (exterior face)			
Pressure faces (i.e. stress cutans)	P	PRF	look like clay films; sand grains uncoated
Slickensides	K	SS	slip face; grooves, striations, glossy or shiny

¹ Individual silt grains are not discernible with a 10X lens. Silt coats occur as a fine, off-white, noneffervescent, "grainy" coat on surfaces.


² Skeletans are (pigment) stripped grains > 2 μm and < 2 mm (Brewer, 1976). Preferably describe either silt coats (grains not discernible with 10X lens), or sand coats (grains discernible with 10X lens).

³ Hypocoats, as used here, are field-scale features commonly expressed only as Redoximorphic Features. Micromorphological hypocoats include non-redox features (Bullock, et al., 1985).


**COATS/
FILMS**



HYPOCOATS




**STRESS
FEATURES**




PED & VOID SURFACE FEATURES - AMOUNT - Estimate the relative percent of the visible surface area that a ped-surface feature occupies in a horizon. (See graphic on next page.) In PDP & NASIS, record the estimate as a numeric percent; e.g., 20%.

Amount Class	Code		Criteria: percent of surface area
	Conv.	NASIS	
Very Few	vf	%	< 5 percent
Few	f	%	5 to < 25 percent
Common	c	%	25 to < 50 percent
Many	m	%	50 to < 90 percent
Very Many	vm	%	≥ 90 percent



5%

25%



50%

90%

PED & VOID SURFACE FEATURES - CONTINUITY (Obsolete in NRCS) - Replaced by **Ped & Void Surface Feature - Amount** in PDP.

Continuity Class	Code (Conv.)	Criteria: Features Occur As
Continuous	C	Entire Surface Cover
Discontinuous	D	Partial Surface Cover
Patchy	P	Isolated Surface Cover

PED & VOID SURFACE FEATURES - DISTINCTNESS - The relative extent to which a ped surface feature visually stands out from adjacent material.

Distinctness Class	Code	Criteria:
Faint	F	Visible with magnification only (10X hand lens); little contrast between materials.
Distinct	D	Visible without magnification; significant contrast between materials.
Prominent	P	Markedly visible without magnification; sharp visual contrast between materials.

PED & VOID SURFACE FEATURES - LOCATION - Specify where ped-surface features occur within a horizon; e.g., *Between sand grains*.

Location	Code	
	PDP	NASIS
PEDS		
On Bottom Faces of Peds	L ¹	BF
On Top Faces of Peds	U ¹	TF
On Vertical Faces of Peds	V	VF
On All Faces of Peds (<i>vertical & horizontal</i>)	P	PF
On Tops of Soil Columns	C	TC
OTHER (NON-PED)		
Between Sand Grains (<i>bridging</i>)	B	BG
On Surfaces Along Pores	I ¹	SP
On Surfaces Along Root Channels	I ¹	SC
On Concretions	O	CC
On Nodules	N	NO
On Rock Fragments	R	RF
On Top Surfaces of Rock Fragments	U ¹	TR
On Bottom Surfaces of Rock Fragments	L ¹	BR

¹ Codes are repeated because these choices are combined in PDP.

PED & VOID SURFACE FEATURES - COLOR - Use standard Munsell® notation (hue, value, chroma) to record feature color.

(SOIL) TEXTURE


This is the numerical proportion (percent by weight) of sand, silt, and clay in a soil. Sand, silt, and clay content is estimated in the field by hand (or quantitatively measured in the office/lab by hydrometer or pipette) and then placed within the texture triangle to determine **Texture Class**. Estimate the **Texture Class**; e.g., *sandy loam*; or **Subclass**; e.g., *fine sandy loam* of the fine earth (< 2 mm) fraction, or choose a **Term in Lieu of Texture**; e.g., *gravel*. If appropriate, use a **Textural Class Modifier**; e.g., *gravelly silt loam*.

NOTE: **Soil Texture** encompasses only the fine earth fraction (< 2 mm). **Particle Size Distribution (PSD)** encompasses the whole soil, including both the fine earth fraction (< 2 mm) and rock fragments (> 2 mm).

TEXTURE CLASS -

Texture Class	Code	
	Conv.	NASIS
Coarse Sand	cos	COS
Sand	s	S
Fine Sand	fs	FS
Very Fine Sand	vfs	VFS
Loamy Coarse Sand	lcos	LCOS
Loamy Sand	ls	LS
Loamy Fine Sand	lfs	LFS
Loamy Very Fine Sand	lvfs	LVFS
Coarse Sandy Loam	cosl	COSL
Sandy Loam	sl	SL
Fine Sandy Loam	fsl	FSL
Very Fine Sandy Loam	vfsl	VFSL
Loam	l	L
Silt Loam	sil	SIL
Silt	si	SI
Sandy Clay Loam	scl	SCL
Clay Loam	cl	CL
Silty Clay Loam	sicl	SICL
Sandy Clay	sc	SC
Silty Clay	sic	SIC
Clay	c	C

**Texture Triangle:
Fine Earth Texture Classes (—)**



TEXTURE MODIFIERS - Conventions for using "Rock Fragment Texture Modifiers" and for using textural adjectives that convey the "% volume" ranges for **Rock Fragments - Size & Quantity**.

Fragment Content % By Volume	Rock Fragment Modifier Usage
< 15	No texture adjective is used (noun only; e.g., <i>loam</i>).
15 to < 35	Use adjective for appropriate size; e.g., <i>gravelly</i> .
35 to < 60	Use "very" with the appropriate size adjective; e.g., <i>very gravelly</i> .
60 to < 90	Use "extremely" with the appropriate size adjective; e.g., <i>extremely gravelly</i> .
≥ 90	No adjective or modifier. If ≤ 10% fine earth, use the appropriate noun for the dominant size class; e.g., <i>gravel</i> . Use Terms in Lieu of Texture .

References for Table Comparing Particle Size Systems

- ¹ Soil Survey Staff. 1995. Soil survey laboratory information manual. USDA - Natural Resources Conservation Service, Soil Survey Investigations Report No. 45, Version 1.0, National Soil Survey Center, Lincoln, NE. 305 p.
- ² International Soil Science Society. 1993. *In: Soil Survey Manual*. Soil Survey Staff, USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C. 503 p.
- ³ ASTM. 1993. Standard classification of soils for engineering purposes (Unified Soil Classification System). ASTM designation D2487-92. *In: Soil and rock; dimension stone; geosynthetics*. Annual book of ASTM standards - Vol. 04.08.
- ⁴ ASSHTO. 1986a. Recommended practice for the classification of soils and soil-aggregate mixtures for highway construction purposes. ASSHTO designation M145-82. *In: Standard specifications for transportation materials and methods of sampling and testing; Part 1: Specifications (14th ed.)*. American Association of State Highway and Transportation Officials, Washington, D.C.
- ⁵ ASSHTO. 1986b. Standard definitions of terms relating to subgrade, soil-aggregate, and fill materials. ASSHTO designation M146-70 (1980). *In: Standard specifications for transportation materials and methods of sampling and testing; Part 1: Specifications (14th ed.)*. American Association of State Highway and Transportation Officials, Washington, D.C.
- ⁶ Ingram, R.L. 1982. Modified Wentworth scale. *In: Grain-size scales*. AGI Data Sheet 29.1. *In: Duto, J.T., Dietrich, R.V., and Foote, R.M.* 1989. AGI data sheets for geology in the field, laboratory, and office, 3rd edition. American Geological Institute, Washington, D.C.

TEXTURE MODIFIERS - (adjectives)

ROCK FRAGMENTS: Size & Quantity ¹	Code		Criteria: Percent (By Volume) of Total Rock Fragments and Dominated By (name size): ¹
	Conv.	PDP / NASIS	
HARD ROCK FRAGMENTS (> 2 mm)			
Gravelly	GR	GR	≥ 15% but < 35% gravel
Fine Gravelly	FGR	GRF	≥ 15% but < 35% fine gravel
Medium Gravelly	MGR	GRM	≥ 15% but < 35% med. gravel
Coarse Gravelly	CGR	GRC	≥ 15% but < 35% coarse gravel
Very Gravelly	VGR	GRV	≥ 35% but < 60% gravel
Extremely Gravelly	XGR	GRX	≥ 60% but < 90% gravel
Cobbly	CB	CB	≥ 15% but < 35% cobbles
Very Cobbly	VCB	CBV	≥ 35% but < 60% cobbles
Extremely Cobbly	XCB	CBX	≥ 60% but < 90% cobbles
Stony	ST	ST	≥ 15% but < 35% stones
Very Stony	VST	STV	≥ 35% but < 60% stones
Extremely Stony	XST	STX	≥ 60% but < 90% stones
Bouldery	BY	BY	≥ 15% but < 35% boulders
Very Bouldery	VBY	BYV	≥ 35% but < 60% boulders
Extremely Bouldery	XBY	BYX	≥ 60% but < 90% boulders
Channery	CN	CN	≥ 15% but < 35% channers
Very Channery	VCN	CNV	≥ 35% but < 60% channers
Extremely Channery	XCN	CNX	≥ 60% but < 90% channers
Flaggy	FL	FL	≥ 15% but < 35% flagstones
Very Flaggy	VFL	FLV	≥ 35% but < 60% flagstones
Extremely Flaggy	XFL	FLX	≥ 60% but < 90% flagstones
PARA (SOFT) ROCK FRAGMENTS (> 2 mm) ^{2,3}			
Parabouldery	PBY	PBY	(same criteria as bouldery)
Very Parabouldery	VPBY	PBYV	(same criteria as very bouldery)
Extr. Parabouldery	XPBY	PBYX	(same criteria as ext. bouldery)
etc.	etc.	etc.	(same criteria as non -para)

¹ The "Quantity" modifier (e.g., *very*) is based on the total rock fragment content. The "Size" modifier (e.g., *cobble*) is based on the largest, dominant fragment size. For a mixture of sizes (e.g., *gravel and stones*), the smaller size must exceed 2X the amount of the larger size to be named (e.g., 30% gravel and 14% stones = *very gravelly*; 20% gravel and 14% stones = *stony*).

² Use "Para" prefix if the rock fragments are soft (i.e., meet criteria for "para"). [Rupture Resistance - Cementation Class < Moderately Cemented, and do not slake (24 hrs in water).]

³ For "Para" codes, add "P" to "Size" and "Quantity" code terms. Precedes noun codes and follows quantity adjectives, e.g., paragravelly = *PGR*; very paragravelly = *VPGR*.

COMPOSITIONAL TEXTURE MODIFIERS ¹- (adjectives)

Types	Code		Criteria:
	PDP	NASIS	
VOLCANIC			
Ashy	--	ASHY	Neither hydrous nor medial and $\geq 30\%$ of the < 2 mm fraction is 0.02 to 2.00 mm in size of which $\geq 5\%$ is volcanic glass
Hydrous	--	HYDR	Andic properties and with field moist 15 bar water content $\geq 100\%$ of the dry weight
Medial	--	MEDL	Andic properties and with field moist 15 bar water content $\geq 30\%$ to < 100% of the dry weight or $\geq 12\%$ water content for air-dried samples
ORGANIC SOILS			
Grassy ²	--	GS	OM > 15% (vol.) grassy fibers
Herbaceous ²	--	HB	OM > 15% (vol.) herbaceous fibers
Mossy ²	--	MS	OM > 15% (vol.) moss fibers
Mucky	MK	MK	Mineral soil > 10% OM and < 17% fibers
Peaty	PT	PT	Mineral soil > 10% OM with > 17% fibers
Woody ²	--	WD	OM $\geq 15\%$ (vol.) wood pieces or fibers
LIMNIC MATERIALS			
Coprogenous	--	COP	
Diatomaceous	--	DIA	
Marly	--	MR	
OTHER			
Gypsiferous	--	GYP	$\geq 15\%$ (weight) gypsum
Permanently Frozen	PF	PF	e.g., Permafrost

¹ **Compositional Texture Modifiers** can be used with the **Soil Texture Name**; e.g., *gravelly ashly loam* or *mossy peat*. For definitions and usage of **Compositional Texture Modifiers**, see the National Soil Survey Handbook - Part 618 (Soil Survey Staff, 1996c).

² Used only with Histosols, histic epipedons, or mucky peats and peats.

TERMS USED IN LIEU OF TEXTURE - (nouns)

Terms Used in Lieu of Texture	Code	
	PDP	NASIS
SIZE (HARD ROCKS)		
Gravel	G	G
Cobbles	CB	CB
Stones	ST	ST
Boulders	B	BY
Channers	--	CN
Flagstones	--	FL
SIZE (SOFT ROCKS)		
Paragravel	--	PGR
Paracobbles	--	PCB
Parastones	--	PST
Paraboulders	--	PBY
Parachanners	--	PCN
Paraflagstones	--	PFL
COMPOSITION		
<i>Cemented / Consolidated:</i>		
Duripan (<i>silica cement</i>)	--	DUR
Ortstein (<i>organic with Fe and Al cement</i>)	--	OR
Petrocalcic (<i>carbonate cement</i>)	--	PC
Petroferric (<i>Fe cement</i>)	--	PF
Petrogypsic (<i>gypsum cement</i>)	--	PGP
Placic Horizon (<i>thin layer cemented by Fe & Mn</i>)	--	PL
Unweathered Bedrock (<i>unaltered</i>)	UWB	UB
Weathered Bedrock (<i>altered; e.g., some Cr horizons</i>)	WB	WB
<i>Organics:</i>		
Highly Decomposed Plant Material (<i>Oa</i>) ¹	---	HPM
Moderately Decomposed Plant Material (<i>Oe</i>) ¹	---	MPM
Slightly Decomposed Plant Material (<i>O</i>) ¹	---	SPM
Muck ² (<i>≈Oa</i>)	---	MUCK
Mucky Peat ² (<i>≈Oe</i>)	---	MPT
Peat ² (<i>≈O</i>)	---	PEAT
<i>Other:</i>		
Finely Stratified (<i>contrasting textures</i>)	--	FS
Ice (<i>permanently frozen</i>) ^{3,4}	--	IC
Water (<i>permanent</i>) ^{3,4}	--	W

¹ Use only with mineral soil layers.

² Use only with Histosols or histic epipedons.

³ Use only for layers found below the soil surface.

⁴ In NASIS, use permanently frozen water to convey permafrost.

ROCK and OTHER FRAGMENTS

These are discrete, water-stable particles > 2 mm. Hard Rock Fragments have a Rupture Resistance - Cementation Class \geq Strongly Cemented. Other Fragments (e.g., soft rock, wood) are less strongly cemented. Describe **Kind**, **Volume Percent** (classes given below), **Roundness or Shape**, and **Size** (mm).

ROCK AND OTHER FRAGMENTS - KIND - Use the choice list given for **Bedrock - Kind** and the additional choices in the table below. **NOTE:** Interbedded rocks from the "Bedrock - Kind Table" are not appropriate choices or terminology for rock fragments.

Kind	Code		Kind	Code	
	PDP	NASIS		PDP	NASIS
Includes all choices in Bedrock - Kind (except <i>Interbedded</i>), plus:					
Calcrete (<i>caliche</i>) ¹	---	CA	Scoria	---	SC
Charcoal	---	CH	Volcanic bombs	---	VB
Cinders	E5	CI	Wood	---	WO
Lapilli	---	LA			

¹ Fragments strongly cemented by carbonate, may include fragments derived from petrocalcic horizons.


ROCK AND OTHER FRAGMENTS - VOLUME PERCENT - Estimate the quantity on a volume percent basis. **NOTE:** For proper use of **Texture Modifiers**, refer to the "Percent Volume Table" found under **Texture**.

ROCK AND OTHER FRAGMENTS - ROUNDNESS - Estimate the relative roundness of rock fragments; use the following classes. (Called **Fragment Roundness** in PDP.)

Roundness Class	Code		Criteria: visual estimate ¹
	PDP	NASIS	
Very Angular	---	VA	[Use Roundness graphic on next page]
Angular	1	AN	
Subangular	2	SA	
Subrounded	3	SR	
Rounded	4	RO	
Well Rounded	5	WR	

¹ The criteria consist of a visual estimation; use the following graphic.

Estimate the relative rounding of rock fragments. (Ideally, use the average roundness based on 50 or more fragments.) The conventional geologic and engineering approach is presented in the following graphic. (**NOTE:** NRCS does not quantify **Sphericity**. It is included here for completeness and to show the range in **Fragment Roundness**.)



¹ After Powers, 1953.

² Numerical values below *roundness* and *sphericity* columns are class midpoints (median rho values) (Folk, 1955) used in statistical analysis.

ROCK AND OTHER FRAGMENTS - SIZE CLASSES AND DESCRIPTIVE TERMS -

Size ¹	Noun	Adjective ²
SHAPE - SPHERICAL or CUBELIKE (discoidal, subdiscoidal, or spherical)		
> 2 - 75 mm diameter	gravel	gravelly
> 2 - 5 mm diameter	fine gravel	fine gravelly
> 5 - 20 mm diameter	medium gravel	medium gravelly
> 20 - 75 mm diameter	coarse gravel	coarse gravelly
> 75 - 250 mm diameter	cobbles	cobbly
> 250 - 600 mm diameter	stones	stony
> 600 mm diameter	boulders	bouldery
SHAPE - FLAT (prismoidal or subprismoidal)		
> 2 - 150 mm long	channers	channery
> 150 - 380 mm long	flagstones	flaggy
> 380 - 600 mm long	stones	stony
> 600 mm long	boulders	bouldery

¹ Fragment size is measured by sieves; class limits have a > lower limit.

² For a mixture of sizes (e.g., gravels and stones present), the smaller size must exceed 2X the quantity of the larger size (e.g., 30% gravel and 14% stones = *gravelly*; but 20% gravel and 14% stones = *stony*).

(SOIL) STRUCTURE


(Soil) Structure is the naturally occurring arrangement of soil particles into aggregates that results from pedogenic processes. Record **Grade, Size, and Type**. For compound structure, list each **Size and Type**; e.g., *medium and coarse SBK parting to fine GR*. Up to ten entries (per horizon) are permitted in PDP. (For PDP only, estimate the percent of each type.) Lack of structure (structureless) has two end members: *massive (MA)* or *single grain (SG)*. A complete example is: *weak, fine, subangular blocky or 1, f, sbk*.

(SOIL) STRUCTURE - TYPE (formerly **Shape**) -


Type	Code Conv NASIS	Criteria: (definition)
NATURAL SOIL STRUCTURAL UNITS (pedogenic structure)		
Granular	gr GR	Small polyhedrals, with curved or very irregular faces.
Angular Blocky	abk ABK	Polyhedrals with faces that intersect at sharp angles (planes).
Subangular Blocky	sbk SBK	Polyhedrals with sub-rounded and planar faces, lack sharp angles.
Platy	pl PL	Flat and tabular-like units.
Wedge	--- WEG	Elliptical, interlocking lenses that terminate in acute angles, bounded by slickensides; not limited to vertic materials.
Prismatic	pr PR	Vertically elongated units with flat tops.
Columnar	cpr COL	Vertically elongated units with rounded tops which commonly are "bleached".
STRUCTURELESS		
Single Grain	sg SGR	No structural units; entirely noncoherent; e.g., loose sand.
Massive	m MA	No structural units; material is a coherent mass (not necessarily cemented).
ARTIFICIAL EARTHY FRAGMENTS OR CLOUDS¹ (non-pedogenic structure)		
Cloddy ¹	--- CDY	Irregular blocks created by artificial disturbance; e.g., tillage or compaction.

¹ Used only to describe oversized, "artificial" earthy units that are not pedogenically derived soil structural units; e.g., the direct result of mechanical alteration; use **Blocky Structure Size** criteria.

Examples of Soil Structure Types



Example of Wedge Structure, Gilgai Microfeatures, & Microrelief



Modified from: Lynn and Williams, Soil Survey Horizons, 1992.

(SOIL) STRUCTURE - GRADE -

Grade	Code	Criteria
Structureless	0	No discrete units observable in place or in hand sample.
Weak	1	Units are barely observable in place or in a hand sample.
Moderate	2	Units well-formed and evident in place or in a hand sample.
Strong	3	Units are distinct in place (undisturbed soil), and separate cleanly when disturbed.


(SOIL) STRUCTURE - SIZE -

Size Class	Code		Criteria: structural unit size ¹ (mm)		
	Conv	NASIS	Granular Platy ² Thickness	Columnar, Prismatic, Wedge ³	Angular & Subangular Blocky
Very Fine (Very Thin ²)	vf (vn ¹)	VF (VN ¹)	< 1	< 10	< 5
Fine (Thin ¹)	f (tn ¹)	F (TN ¹)	1 to < 2	10 to < 20	5 to < 10
Medium	m	M	2 to < 5	20 to < 50	10 to < 20
Coarse (Thick ²)	co (tk ²)	CO (TK ²)	5 to < 10	50 to < 100	20 to < 50
Very Coarse (Very Thick)	vc (vk ²)	VC (VK ²)	≥ 10	100 to < 500	≥ 50
Extr. Coarse	ec	EC	---	≥ 500	---

- ¹ Size limits always denote the smallest dimension of the structural units.
- ² For platy structure only, substitute *thin* for *fine* and *thick* for *coarse* in the size class names.
- ³ Wedge structure is generally associated with Vertisols (for which it is a requirement) or related soils with high amounts of smectitic clays.

Granular

Codes



Platy

Codes


Very Thin
(<1 mm thick)

VN



Thin
(1 to <2 mm thick)

TN




Medium
(2 to <5 mm thick)

M



Thick
(5 to <10 mm thick)

TK




Very Thick
(≥ 10 mm thick)

VK


Angular & Subangular Blocky

Codes




Prismatic & Columnar

Codes



Wedge

Codes




CONSISTENCE


Consistence is the degree and kind of cohesion and adhesion that soil exhibits, and/or the resistance of soil to deformation or rupture under an applied stress. Soil-water state strongly influences consistence. Field evaluations of consistence include: **Rupture Resistance** (Blocks, Peds, and Clods; or Surface Crusts and Plates), **Resistance to Penetration**, **Plasticity**, **Stickiness**, and **Manner of Failure**. Historically, consistence applied to dry, moist, or wet soil as observed in the field. Wet consistence evaluated Stickiness and Plasticity. **Rupture Resistance** now applies to dry soils and to soils in a water state from moist through wet. **Stickiness** and **Plasticity** of soil are independent evaluations.

RUPTURE RESISTANCE - A measure of the strength of soil to withstand an applied stress. Separate estimates of **Rupture Resistance** are made for **Blocks/Peds/Clods** and for **Surface Crusts and Plates** of soil. Block-shaped specimens should be approximately 2.8 cm across. If 2.8 cm cubes (e.g., $\approx 2.5 - 3.1$ cm) are not obtainable, use the following equation and the table below to calculate the stress at failure: $[(2.8 \text{ cm} / \text{cube length cm})^2 \times \text{estimated stress (N) at failure}]$; e.g., for a 5.6 cm cube $[(2.8/5.6)^2 \times 20 \text{ N} = 5 \text{ N} \Rightarrow \text{Soft Class}$. Plate-shaped specimens (surface crusts or platy structure) should be approximately 1.0 - 1.5 cm long by 0.5 cm thick (or the thickness of occurrence, if < 0.5 cm thick).

Blocks/Peds



Crusts/Plates



RUPTURE RESISTANCE for:

Blocks, Peds, and Clods - Estimate the class by the force required to rupture (break) a soil unit. Select the column for the appropriate soil moisture condition (*dry vs. moist*) and / or the *Cementation* column, if applicable.

Dry		Moist		Cementation ¹		Specimen Fails Under
Class	Code ²	Class	Code ²	Class	Code ²	
Loose	L	Loose	L	<i>Not Applicable</i>		<i>Intact specimen not obtainable</i>
	d(lo)		m(lo)			
Soft	S	Very Friable	VFR	Non-Cemented	NC	Very slight force between fingers. <8 N
	d(so)		m(vfr)			
Slightly Hard	SH	Friable	FR	Extremely Weakly Cemented	EW	Slight force between fingers. 8 to < 20 N
	d(sh)		m(fr)			
Mod. Hard	MH	Firm	FI	Very Weakly Cemented	VW	Moderate force between fingers. 20 to < 40 N
	d(h)		m(fi)			
Hard	HA	Very Firm	VFI	Weakly Cemented	W c(w)	Strong force between fingers. 40 to < 80 N
	d(h)		m(vfi)			
Very Hard	VH	Extr. Firm	EF	Moderately Cemented	M	Moderate force between hands. 80 to < 160 N
	d(vh)		m(efi)			
Extremely Hard	EH	Slightly Rigid	SR	Strongly Cemented	ST c(s)	Foot pressure by full body weight. 160 to < 800 N
	d(eh)		m(efi)			
Rigid	R	Rigid	R	Very Strongly Cemented	VS	Blow of < 3 J but not body weight. 800 N to < 3 J
	d(eh)		m(efi)			
Very Rigid	VR	Very Rigid	VR	Indurated	I c(l)	Blow of ≥ 3 J. (3 J = 2 kg weight dropped 15 cm).
	d(eh)		m(efi)			

¹ This is not a field test; specimen must be air dried overnight and then submerged in water for a minimum of 1 hour prior to test.

² Codes in parentheses are obsolete criteria (Soil Survey Staff, 1951).

Soil Moisture Status (Consistence) (OBSOLETE) - Historical classes (Soil Survey Staff, 1953).

(d) ¹ Dry Soil Class ²		(m) ¹ Moist Soil Class		Cementation Class	
Code		Code		Code	
Loose	(d) lo	Loose	(m) lo	Weakly Cemented	(c) w
Soft	(d) so	Very Friable	(m) vfr	Strongly Cemented	(c) s
Slightly Hard	(d) sh	Friable	(m) fr		
Hard ²	(d) h	Firm	(m) fi	Indurated	(c) I
Very Hard	(d) vh	Very Firm	(m) vfi		
Extr. Hard	(d) eh	Extr. Firm	(m) efi		

¹ Historically, consistence prefixes (*d* for dry, *m* for moist) were commonly omitted, leaving only the root code; e.g., *vfr* for *mvfr*.

² *Hard Class (Dry)* was split into *Moderately Hard* and *Hard* (Soil Survey Staff, 1993).

Surface Crust and Plates -

Class (air dried)	Code	Force ¹ (Newtons)
Extremely Weak	EW	<i>Not Obtainable</i>
Very Weak	VW	Removable, < 1N
Weak	W	1 to < 3N
Moderate	M	3 to < 8N
Moderately Strong	MS	8 to < 20N
Strong	S	20 to < 40N
Very Strong	VS	40 to < 80N
Extremely Strong	ES	≥ 80N

¹ For operational criteria [field estimates of force (N)] use the *Fails Under* column, in the "Rupture Resistance for Blocks, Peds, Clods Table".

CEMENTING AGENTS - Record kind of cementing agent, if present.

Kind	Code ¹
carbonates	K
gypsum	G
humus	H
iron	I
silica (SiO ₂)	S

¹ Conventional codes traditionally consist of the entire material name or its chemical symbols; e.g., *silica* or *SiO₂*. Consequently, the *Conv.* code column would be redundant and is not shown in this table.

MANNER OF FAILURE - The rate of change and the physical condition soil attains when subjected to compression. Samples are moist or wetter.

Failure Class	Code		Criteria: Related Field Operation
	PDP	NASIS	
BRITTLENESS			
<i>Use a 3 cm block. (Press between thumb & forefinger.)</i>			
Brittle	B	BR	Ruptures abruptly ("pops" or shatters).
Semi-Deformable	SD	SD	Rupture occurs before compression to < 1/2 original thickness.
Deformable	D	DF	Rupture occurs after compression to \geq 1/2 original thickness.
FLUIDITY			
<i>Use a palmful of soil. (Squeeze in hand.)</i>			
Nonfluid	NF	NF	No soil flows through fingers with full compression.
Slightly Fluid	SF	SF	Some soil flows through fingers, most remains in the palm, after full pressure.
Moderately Fluid	MF	MF	Most soil flows through fingers, some remains in palm, after full pressure.
Very Fluid	VF	VF	Most soil flows through fingers, very little remains in palm, after gentle pressure.
SMEARINESS			
<i>Use a 3 cm block. (Press between thumb & forefinger.)</i>			
Non-Smeary ¹	NS	NS	At failure, the sample does not change abruptly to fluid, fingers do not skid, no smearing occurs.
Weakly Smeary ¹	WS	WS	At failure, the sample changes abruptly to fluid, fingers skid, soil smears, little or no water remains on fingers.
Moderately Smeary ¹	MS	MS	At failure, the sample changes abruptly to fluid, fingers skid, soil smears, some water remains on fingers.
Strongly Smeary ¹	SM	SM	At failure, the sample abruptly changes to fluid, fingers skid, soil smears and is slippery, water easily seen on fingers.

¹ *Smeary* failure classes are used dominantly with Andic materials, but may also be used with some spodic materials.

STICKINESS - The capacity of soil to adhere to other objects. Stickiness is estimated at the moisture content that displays the greatest adherence when pressed between thumb and forefinger.


Stickiness Class	Code			Criteria: Work moistened soil between thumb and forefinger
	Conv	PDP	NASIS	
Non-Sticky	(w) so	SO	SO	Little or no soil adheres to fingers, after release of pressure.
Slightly Sticky	(w) ss	SS	SS	Soil adheres to both fingers, after release of pressure. Soil stretches little on separation of fingers.
Moderately Sticky ¹	(w) s	S	MS	Soil adheres to both fingers, after release of pressure. Soil stretches some on separation of fingers.
Very Sticky	(w) vs	VS	VS	Soil adheres firmly to both fingers, after pressure release. Soil stretches greatly upon separation of fingers.

¹ Historically, the *moderately sticky* class was simply called *sticky*.

PLASTICITY - The degree to which "puddled" or reworked soil can be permanently deformed without rupturing. The evaluation is made by forming a roll (wire) of soil at a water content where the maximum plasticity is expressed.

Plasticity Class	Code			Criteria: Make a roll of soil 4 cm long
	Conv	PDP	NASIS	
Non-Plastic	(w) po	PO	PO	Will not form a 6 mm diameter roll, or if formed, can't support itself if held on end.
Slightly Plastic	(w) ps	SP	SP	6 mm diameter roll supports itself; 4 mm diameter roll does not.
Moderately Plastic ¹	(w) p	P	MP	4 mm diameter roll supports itself, 2 mm diameter roll does not.
Very Plastic	(w) vp	VP	VP	2 mm diameter roll supports its weight.

¹ Historically, the *moderately plastic* class was simply called *plastic*.



PENETRATION RESISTANCE - The ability of soil in a confined (field) state to resist penetration by a rigid object of specified size. A pocket penetrometer (Soil-Test Model CL-700) with a rod diameter of 6.4 mm (area 20.10 mm²) and insertion distance of 6.4 mm (note line on rod) is used for the determination. An average of five or more measurements should be used to obtain a value for penetration resistance. In PDP, record the **Penetration Resistance** value in mega-pascals (MPa), **Orientation** of the rod (vertical (V) or horizontal (H)), and **Water State** of the soil.

NOTE: The pocket penetrometer has a scale of 0.25 to 4.5 tons/ft² (tons/ft² ≈ kg/cm²). The penetrometer does not directly measure penetration resistance. The penetrometer scale is correlated to, and gives a field estimate of unconfined compressive strength of soil as measured with a Tri-Axial Shear device. The table below converts the scale reading on the pocket penetrometer to penetration resistance in MPa. Penetrometer readings are dependent on the spring type used. Springs of varying strength are needed to span the range of penetration resistance found in soil.

Penetrometer Scale Reading tons / ft ²	Spring Type ^{1, 2, 3}			
	Original MPa	Lee MPa	Jones 11 MPa	Jones 323 MPa
0.25	0.32 L	0.06 VL	1.00 M	3.15 H
0.75	0.60	0.13 L	1.76	4.20
1.00	0.74	0.17	2.14 H	4.73
1.50	1.02 M	0.24	2.90	5.78
2.75	1.72	0.42	4.80	8.40 EH
3.50	2.14 H	0.53	---	---

- ¹ On wet or "soft" soils, a larger "foot" may be used (Soil Survey Staff, 1993).
- ² Each bolded value highlights the force associated with a rounded value on the penetrometer scale that is closest to a *Penetration Resistance Class* boundary. The bolded letter; e.g., **M**, represents the moderate *Penetration Resistance Class* from the following table.
- ³ Each spring type spans only a part of the range of penetration resistance possible in soils; various springs are needed to span all *Penetration Resistance Classes*.

Penetration Resistance Class	Code	Criteria: Penetration Resistance (MPa)
Extremely Low	EL	< 0.01
Very Low	VL	0.01 to < 0.1
Low	L	0.1 to < 1
Moderate	M	1 to < 2
High	H	2 to < 4
Very High	VH	4 to < 8
Extremely High	EH	≥ 8

EXCAVATION DIFFICULTY - The relative force or energy required to dig soil out of place. Describe the **Excavation Difficulty Class** and the moisture condition (*moist* or *dry*, but not *wet*); use the "(Soil) Water State Table"; e.g., *moderate, moist* or *M, M*. Estimates can be made for either the most limiting layer or for each horizon.

Class	Code	Criteria
Low	L	Excavation by tile spade requires arm pressure only; impact energy or foot pressure is not needed.
Moderate	M	Excavation by tile spade requires impact energy or foot pressure; arm pressure is insufficient.
High	H	Excavation by tile spade is difficult, but easily done by pick using over-the-head swing.
Very High	VH	Excavation by pick with over-the-head swing is moderately to markedly difficult. Backhoe excavation by a 50-80 hp tractor can be made in a moderate time.
Extremely High	EH	Excavation via pick is nearly impossible. Backhoe excavation by a 50-80 hp tractor cannot be made in a reasonable time.

ROOTS

Record the **Quantity**, **Size**, and **Location** of roots in each horizon. **NOTE:** Describe **Pores** using the same **Quantity** and **Size** classes and criteria as **Roots** (use the combined tables). A complete example for roots is: *Many, fine, roots In Mat at Top of Horizon or 3, f (roots), M.*

ROOTS - QUANTITY (Roots and Pores) - Describe the quantity (number) of roots for each size class in a horizontal plane. (**NOTE:** Typically, this is done across a vertical plane, such as a pit face.) Record the average quantity from 3 to 5 representative unit areas. **CAUTION:** The unit area that is evaluated varies with the *Size Class* of the roots being considered. Use the appropriate unit area stated in the *Soil Area Observed* column of the "Size (Roots and Pores) Table". In NASIS and PDP, record the actual number of roots/unit area (which outputs the appropriate class). Use class names in narrative description.

Quantity Class ¹	Code		Average Count ² (per unit area)
	Conv	NASIS	
Few	1	#	< 1 per area
Very Few ¹	---	#	< 0.2 per area
Moderately Few ¹	---	#	0.2 to < 1 per area
Common	2	#	1 to < 5 per area
Many	3	#	≥ 5 per area

¹ The *Very Few* and *Moderately Few* sub-classes can be described for roots (optional) but do not apply to pores.



² The applicable area for appraisal varies with the size of roots or pores. Use the appropriate area stated in the *Soil Area Assessed* column of the "Size (Roots and Pores) Table" or use the following graphic.

ROOTS - SIZE (Roots and Pores) - See the following graphic for size.


Size Class	Code		Diameter	Soil Area ¹ Assessed
	Conv	NASIS		
Very Fine	vf	VF	< 1 mm	1 cm ²
Fine	f	F	1 to < 2 mm	1 cm ²
Medium	m	M	2 to < 5 mm	1 dm ²
Coarse	co	C	5 to < 10 mm	1 dm ²
Very Coarse	vc	VC	≥ 10 mm	1 m ²

¹ One dm² = a square that is 10 cm on a side, or 100 cm².

ROOTS - QUANTITY (Roots and Pores) - Soil area to be assessed.



Root and Pore Size Classes



ROOTS - LOCATION (Roots) -

Location	Code
Between Peds	P
In Cracks	C
Throughout	T
In Mat at Top of Horizon ¹	M
Matted Around Rock Fragments	R

¹ Describing a root mat at the top of a horizon rather than at the bottom or within the horizon, flags the horizon that restricts root growth.

PORES DISCUSSION

Pores are the air or water filled voids in soil. Historically, description of soil pores, called "nonmatrix" pores in the Soil Survey Manual (Soil Survey Staff, 1993), excluded inter-structural voids, cracks, and in some schemes, interstitial pores. *Inter-structural voids* (i.e., the sub-planar fractures between peds; also called interpedal or structural faces/planes), which can be inferred from soil structure descriptions, are not recorded directly. *Cracks* can be assessed independently (Soil Survey Staff, 1993). *Interstitial pores* (i.e. visible, primary packing voids) may be visually estimated, especially for fragmental soils, or can be inferred from soil porosity, bulk density, and particle size distribution. Clearly, one cannot assess the smallest interstitial pores (e.g., < 0.05 mm) in the field. Field observations are limited to those that can be seen through a 10X hands lens, or larger. Field estimates of interstitial pores are considered to be somewhat tenuous, but useful.

PORES

Record **Quantity** and **Size** of pores for each horizon. Description of soil pore **Shape** and **Vertical Continuity** is optional. A complete example for pores is: *common, medium, tubular pores, throughout* or *c, m, TU (pores), T*.

PORES - QUANTITY - See and use **Quantity (Roots and Pores)**.

PORES - SIZE - See and use **Size (Roots and Pores)**.

PORES - SHAPE (or Type) - Record the dominant form (also called "type") of pores discernible with a 10X hand lens and by the unaided eye. (See following graphic.)


Description	Code		Criteria
	PDP	NASIS	
SOIL PORES¹			
Dendritic Tubular	TE	DT	Cylindrical, elongated, branching voids; e.g., <i>empty root channels</i> .
Irregular	---	IG	Non-connected cavities, chambers; e.g., <i>vughs</i> ; various shapes.
Tubular	TU	TU	Cylindrical and elongated voids; e.g., <i>worm tunnels</i> .
Vesicular	VS	VE	Ovoid to spherical voids; e.g., <i>solidified pseudomorphs of entrapped, gas bubbles concentrated below a crust</i> ; most common in arid to semi-arid environments.
PRIMARY PACKING VOIDS²			
Interstitial	IR	IR	Voids between sand grains or rock fragments.

¹ Called "Nonmatrix Pores" in the Soil Survey Manual (Soil Survey Staff, 1993).

² *Primary Packing Voids* include a continuum of sizes. As used here, they have a minimum size that is defined as pores that are visible with a 10X hand lens. *Primary Packing Voids* are called "Matrix Pores" in the Soil Survey Manual (Soil Survey Staff, 1993).


Tubular

(e.g. small worm tunnels)




Vesicular

(e.g. isolated, spherical-ovoid cavities)




Dendritic Tubular

(e.g. abandoned root channels)




Irregular

(e.g. vughs)



Interstitial

(e.g. primary packing voids)




PORES - VERTICAL CONTINUITY - The average vertical distance through which the minimum pore diameter exceeds 0.5 mm. Soil must be moist or wetter.

Class	Code		Criteria: vertical distance
	Conv.	NASIS	
Low	---	L	< 1 cm
Moderate	---	M	1 to < 10 cm
High	---	H	≥ 10 cm

CRACKS

Also called "Extra-Structural Cracks" (Soil Survey Staff, 1993) are fissures other than those attributed to soil structure. Cracks are commonly vertical, sub-planar, polygonal, and are the result of desiccation, dewatering, or consolidation of earthy material. Cracks are much longer and can be much wider than planes that surround soil structural units such as prisms, columns, etc. Cracks are key to preferential flow, also called "bypass flow" (Bouma, et al., 1982) and are a primary cause of temporal (transient) changes in ponded infiltration and hydraulic conductivity in soils (Soil Survey Staff, 1993). Cracks are primarily associated with, but not restricted to, clayey soils and are most pronounced in high shrink-swell soils (high COLE value). Record the **Kind**, **Depth**, and **Relative Frequency** (Areal Percentage). A complete example is: *3, 25 cm deep, reversible trans-horizon cracks*.



CRACKS - KIND - Identify the dominant types of fissures.

Kind	Code ¹	General Description
CRUST-RELATED CRACKS ² (shallow, vertical cracks related to crusts; derived from raindrop-splash and soil puddling, followed by dewatering / consolidation and desiccation)		
Reversible Crust-Related Cracks ³	RCR	Very shallow (e.g., 0.1 - 0.5 cm); very transient (generally persist less than a few weeks); formed by drying from surface down; minimal, seasonal influence on ponded infiltration (e.g., <i>rain-drop crust cracks</i>).
Irreversible Crust-Related Cracks ⁴	ICR	Shallow (e.g., 0.5 - 2 cm); seasonally transient (not present year-round nor every year); minor influence on ponded infiltration (e.g., <i>freeze-thaw crust & associated cracks</i>).
TRANS-HORIZON CRACKS ⁵ (deep, vertical cracks that commonly extend across more than one horizon and may extend to the surface; derived from wetting and drying or original dewatering and consolidation of parent material)		
Reversible Trans-Horizon Cracks ⁶	RTH	Transient (commonly seasonal; close when rewetted); large influence on ponded infiltration and Ksat; formed by wetting and drying of soil; (e.g., <i>Vertisols, vertic subgroups</i>).
Irreversible Trans-Horizon Cracks ⁷	ITH	Permanent (persist year-round; see Soil Taxonomy), large influence on ponded infiltration and Ksat (e.g., <i>extremely coarse subsurface fissures within glacial till; drained polder cracks</i>).

¹ No conventional codes, use entire term; NASIS codes are shown.

² Called "Surface-Initiated Cracks" (Soil Survey Staff, 1993).

³ Called "Surface-Initiated Reversible Cracks" (Soil Survey Staff, 1993).


⁴ Called "Surface-Initiated Irreversible Cracks" (Soil Survey Staff, 1993).

⁵ Also called "Subsurface-Initiated Cracks" (Soil Survey Staff, 1993).

⁶ Called "Subsurface-Initiated Reversible Cracks" (Soil Survey Staff, 1993).

⁷ Called "Subsurface-Initiated Irreversible Cracks" (Soil Survey Staff, 1993).

Crust-related Cracks Trans-horizon Cracks



CRACKS - DEPTH - Record the **Average, Apparent Depth** (also called a "depth index value" in the Soil Survey Manual), measured from the surface, as determined by the wire-insertion method (\cong 2 mm diameter wire). **NOTE:** This method commonly gives a standard but conservative measure of the actual fracture depth. Do not record this data element for cracks that are not open to the surface. Depth (and apparent vertical length) of subsurface cracks can be inferred from the *Horizon Depth* column of layers exhibiting subsurface cracks.

CRACKS - RELATIVE FREQUENCY - Record the **Average Number of Cracks**, per meter, across the surface or **Lateral Frequency** across a soil profile as determined with a line-intercept method. This data element cannot be assessed from cores or push tube samples.

SPECIAL FEATURES

Record **Kind** and **Area (%) Occupied**. Describe the special soil feature by kind, and estimate the cross sectional area (%) of the horizon that the feature occupies. In PDP, three items are grouped in this data element: 1) **Special Features** - both Kind (e.g., *krotovinas* and *tongues*) and the Percent (%) of Area Covered (the area a feature occupies within a horizon); 2) **Percent of Profile** - estimate the area of the profile an individual horizon comprises; and 3) **Percent (Volume) of Pedon** occupied.

SPECIAL FEATURES - KIND - Identify the kind of special soil feature.

Kind	Code ¹	Criteria
desert pavement	DP	A natural, concentration of closely packed and polished stones at the soil surface in a desert (may or may not be an erosional lag).
hydrophobic layer	HL	Either a surface or subsurface layer that repels water (e.g. dry organic materials; scorch layers in chaparral, etc.).
ice wedge cast	IC	A vertical, often trans-horizon, wedge-shaped or irregular form caused by infilling of a cavity as an ice wedge melts, commonly stratified.
krotovinas	KR	Filled faunal burrows.
lamellae ²	---	Thin (e.g., > 1 cm), pedogenically formed plates or intermittent layers.
lamina	LN	Thin (e.g., < 1 cm), geogenically deposited strata or layers of alternating texture (e.g., silt and fine sand or silt and clay).
microbiotic crust	MC	Thin, biotically dominated ground or surface crusts; e.g., cryptogamic crust (algae, lichen, mosses, or cyanobacteria).
stone line	SL	A natural concentration of rock fragments caused by water erosion or transport erosional lag (i.e. carpedolith).
Tongues of Albic Material	E	
Tongues of Argillic Material	B	

¹ Conventional codes consist of the entire name; e.g., *Tongues of Albic Material*. Consequently, no *Conv. code* is shown.

² In NASIS, described under **Diagnostic Horizon or Property - Kind**.

SPECIAL FEATURES - AREA (%) OCCUPIED - Estimate the cross sectional area (%) of the horizon that the feature occupies.

PERMEABILITY / SATURATED HYDRAULIC CONDUCTIVITY (DISCUSSION)

The traditional SCS (now NRCS) concept of soil permeability and permeability classes are becoming obsolete. The concept of permeability was originally derived from the "permeability coefficient" as used by engineers (Soil Survey Staff, 1951). Specifically, the permeability coefficient represents the ability of a porous medium to transmit fluids or gases. It is a unitless coefficient totally independent of the working fluid; e.g., water, air, hydrocarbons, molasses.

Permeability (as traditionally used by NRCS) considers only water, at field saturation, as the working fluid. This results in units of length / time; (e.g., inches / hour, cm / hr, etc.) and values that can't be extrapolated to other fluids (e.g., hydrocarbons). Furthermore, permeability (as used by NRCS) has changed through time. The original work (O'Neil, 1952) measured falling head, vertical K_{sat} for a limited number of soil cores and referred to the permeability coefficient. Over time, the term "coefficient" was dropped. Extrapolation and inference from the original, modest K_{sat} data set resulted in widespread estimations of the ability of other soils to internally transmit water. Hence, permeability is now a qualitative estimate who's "values" (i.e., classes) are inferred from soil texture or other proxies instead of actual measurements (Exhibit 618-9, NSSH; Soil Survey Staff, 1996c). It is a soil quality, as is soil tilth, which cannot be directly quantified.

A much preferred parameter (and concept) has largely replaced permeability. **Hydraulic Conductivity (K)** is the current standard for measuring a soil's ability to transmit water. Hydraulic conductivity quantifies a material's ability to transmit water. Hydraulic conductivity is a numerical variable in an equation that can be either measured or estimated. It is one of the terms in Darcy's law: $Q = K A i$, [where "Q" is outflow (volume), "K" is the hydraulic conductivity of the material, "A" is the area through which the fluid moves per unit time, and "i" is the pressure gradient ($\Delta Head / \Delta Length$); (Amoozegar and Warrick, 1986; Bouma, et al., 1982)].

Hydraulic conductivity under saturated conditions is called **Saturated Hydraulic Conductivity (K_{sat})** and is the easiest condition to assess. It is also the most common reference datum used to compare water movement in different soils, layers, or materials.

Permeability is a qualitative estimate of the relative ease with which soil transmits water. Hydraulic conductivity is a specific mathematical coefficient (quantitative) that relates the rate of water movement to the hydraulic gradient.

Direct measurement of saturated hydraulic conductivity (K_{sat}) is strongly recommended rather than an estimation of permeability inferred from other soil properties. **NOTE:** It's highly recommended to determine the K_{sat} of a soil layer by averaging at least three determinations (\approx replications); more reps (e.g., ≥ 5) are preferred. K_{sat} is notoriously variable due to unequal distribution of soil pores and temporal changes in some soil voids (e.g., cracks, bio-pores, etc.). Replications help to capture the natural variation of K_{sat} within soils and to reduce the influence of data population outliers.

NOTE: As with the virtuous child and the non-virtuous look-alike, superficial similarities are deceptive. Permeability and K_{sat} are not synonyms and should not be treated as such.

PERMEABILITY

Estimate the **Permeability Class** for each horizon. Guidelines for estimating permeability are found in Exhibit 618-9, NSSH (Soil Survey Staff, 1996c).

Permeability Class	Code		Criteria: estimated in / hr ¹
	PDP	NASIS	
Impermeable	IM	IM	< 0.0015
Very Slow	VS	VS	0.0015 to < 0.06
Slow	S	SL	0.06 to < 0.2
Moderately Slow	MS	MS	0.2 to < 0.6
Moderate	M	MO	0.6 to < 2.0
Moderately Rapid	MR	MR	2.0 to < 6.0
Rapid	RA	RA	6.0 to < 20
Very Rapid	VR	VR	≥ 20

¹ These class breaks were originally defined in English units and are retained here, as no convenient metric equivalents are available.

SATURATED HYDRAULIC CONDUCTIVITY (K_{SAT})

Saturated Hydraulic Conductivity is used to convey the rate of water movement through soil under (field) saturated conditions. Record the **Average K_{sat} (X)**, **Standard Deviation (s)**, and **Number of Replications (n)** of each major layer/horizon as measured with a constant-head method (e.g., Amoozemeter, Guelph Permeameter, etc.). **NOTE:** This data element should be measured rather than estimated and subsequently placed into classes. Estimates of water movement based on texture or other proxies must use the preceding "Permeability Class Table".

K _{sat} Class	Code ¹		Criteria ² :	
	PDP	NASIS	cm / hr	in / hr
Very Low	1	#	< 0.0036	< 0.001417
Low	2	#	0.00360 to < 0.036	0.001417 to < 0.01417
Mod. Low	3	#	0.0360 to < 0.360	0.01417 to < 0.1417
Mod. High	4	#	0.360 to < 3.60	0.1417 to < 1.417
High	5	#	3.60 to < 36.0	1.417 to < 14.17
Very High	6	#	≥ 36.0	≥ 14.17

¹ There are no "codes" for K_{sat}; record the average of measured K_{sat} values (#) which can then be assigned to the appropriate class.

² For alternative units commonly used for these class boundaries [e.g., Standard International Units (Kg s / m³)], see the Soil Survey Manual (Soil Survey Staff, 1993; p 107).

CHEMICAL RESPONSE

Chemical response is the response of a soil sample to an applied chemical solution or a measured chemical value. Responses are used to identify the presence or absence of certain materials; to obtain a rough assessment of the amount present; to measure the intensity of a chemical parameter (e.g., pH.); or to gauge the "reducing" status of the soil.

REACTION (pH) - (Called **Field pH** in NASIS.) Record the pH value to the nearest tenth, as measured by pH meter for 1:1 (water:soil), or estimated by the Hellige-Truog® field kit. In PDP, record **pH** by other techniques (e.g., CaCl₂ or Lamotte pH) as a **User Defined Property**.

Descriptive Term	Code ¹	Criteria: pH range
Ultra Acid	#	< 3.5
Extremely Acid	#	3.5 to 4.4
Very Strongly Acid	#	4.5 to 5.0
Strongly Acid	#	5.1 to 5.5
Moderately Acid	#	5.6 to 6.0
Slightly Acid	#	6.1 to 6.5
Neutral	#	6.6 to 7.3
Slightly Alkaline	#	7.4 to 7.8
Moderately Alkaline	#	7.9 to 8.4
Strongly Alkaline	#	8.5 to 9.0
Very Strongly Alkaline	#	> 9.0

¹ No "codes"; enter the measured value; class is assigned by PDP.

EFFERVESCENCE - The gaseous response (seen as bubbles) of soil to applied HCl (carbonate test), H₂O₂ (MnO₂ test), or other chemicals. Commonly, ≈1 N HCL is used. Apply the chemical to the soil matrix (for HCL, effervescence refers only to the matrix; do not include carbonate masses, which are described as "concentrations"). Record **Effervescence Class** and **Chemical Agent**. A complete example is: *strongly effervescent with 1N-HCL or 2, 1*. In PDP, record percent of carbonate (measured with a carbonate field kit) as a **User Defined Property**.

Effervescence - Class -

Effervescence Class	Code		Criteria
	PDP	NASIS	
Noneffervescent	4	NE	No bubbles form.
Very Slightly Effervescent	0	VS	Few bubbles form.
Slightly Effervescent	1	SL	Numerous bubbles form.
Strongly Effervescent	2	ST	Bubbles form a low foam.
Violently Effervescent	3	VE	Bubbles form a thick foam.

Effervescence - Location - Use locations (and codes) from **(Ped & Void) Surface Features - Location**. **NOTE:** Application of chemicals (e.g., HCL acid) to soil matrix makes many location choices invalid.

Effervescence - Chemical Agent -

Effervescence Agent	Code		Criteria
	PDP	NASIS	
HCl (unspecified) ¹	H	H1	Hydrochloric Acid: Concentration Unknown
HCl (1N) ^{1,2}	I	H2	Hydrochloric Acid: Concentration = 1 Normal
HCl (3N) ^{1,3}	J	H3	Hydrochloric Acid: Concentration = 3 Normal
HCl (6N) ^{1,4}	---	H4	Hydrochloric Acid: Concentration = 6 Normal
H ₂ O ₂ (unspecified) ^{5,6}	P	P1	Hydrogen Peroxide: Concentration Unknown
H ₂ O ₂ ^{5,6}	O	P2	Hydrogen Peroxide: Concentration 3-4%

¹ Positive reaction indicates presence of carbonates (e.g., CaCO₃).
² Concentration of acid preferred for the effervescence field test.
NOTE: A (1N HCl) solution is made by combining 1 part concentrated (37%) HCl (which is widely available) with 11 parts distilled H₂O.

- ³ This concentration is not used for determining **Effervescence Class**, but is required for the calcium carbonate equivalent test (CO₂ evolution, not effervescence). An approximately 3N HCl solution (actually 10% HCl or 2.87N) is made by combining 6 parts concentrated (37%) HCl (which is widely available) with 19 parts distilled H₂O.
- ⁴ This concentration is not used for determining **Effervescence Class**, but is preferred for the dolomite test (effervescence by dolomitic carbonates). A 6N HCl solution is made by combining 2 parts concentrated (37%) HCl (which is widely available) with 11 parts distilled H₂O. Soil sample should be saturated in a spot plate and allowed to react for 1-2 minutes; froth = positive response. Reaction is slower and less robust than CaCO₃ effervescence.
- ⁵ Positive reaction indicates presence of manganese oxides (e.g., MnO₂).
- ⁶ Some forms of organic matter will react slowly with (3-4%) H₂O₂, whereas Mn reacts rapidly.

REDUCED CONDITIONS -

Chemical Agent	Code	Criteria
α, α' -dipyridyl ⁷	P (= <i>positive</i>) N (= <i>negative</i>)	α, α' -dipyridyl conc.= 0.2%, (Childs, 1981)

⁷ Positive reaction indicates presence of Fe⁺² (i.e., reduced conditions).

SALINITY - The concentration of dissolved salts (more soluble than gypsum; e.g., NaCl) in a water extract. Estimate the **Salinity Class**. If the electrical conductivity is measured, record the actual value and the method used.

Salinity Class	Code	Criteria: (Electrical Conductivity) dS/m (mmhos/cm)
Non-Saline	0	< 2
Very Slightly Saline	1	2 to < 4
Slightly Saline	2	4 to < 8
Moderately Saline	3	8 to < 16
Strongly Saline	4	≥ 16

SODIUM ADSORPTION RATIO (SAR) - An indirect estimate of the equilibrium between soluble sodium (Na) in a salt solution and the exchangeable Na adsorbed by the soil (Soil Survey Staff, 1995). It is presented in the form of a ratio. It is used for soil solution extracts and

irrigation waters to express the relative activity of sodium (Na) ions in exchange reactions with the soil. It is calculated from: $SAR = [Na^+] / [(Ca^{+2} + Mg^{+2}) / 2]^{0.5}$, where "x" is the cation concentration in millimoles per liter. As a field method, it is commonly determined with soil paste and an electronic wand.

ODOR

Record the presence of any strong smell, by horizon. No entry implies no odor. (Proposed for addition to NASIS.)

Odor - Kind	Code	Criteria
Sulphurous	S	Presence of H ₂ S (hydrogen sulfide); "rotten eggs"; commonly associated with strongly reduced soil containing sulfur compounds.
Petrochemical	P	Presence of gaseous or liquid gasoline, oil, creosote, etc.

MISCELLANEOUS FIELD NOTES

Use additional adjectives, descriptors, and sketches to capture and convey pertinent information and any features for which there is no pre-existing data element or code. Record such additional information as free-hand notes under **Field Notes** ("User Defined Entries" in PDP).

MINIMUM DATA SET (for a soil description)

Purpose, field logistics, habits, and soil materials all influence the specific properties necessary to "adequately" describe a given soil. However, some soil properties or features are so universally essential for interpretations or behavior prediction that they should always be recorded. These include: **Location, Horizon, Horizon Depth, Horizon Boundary, Color, Redoximorphic Features, Texture, Structure, and Consistence.**

PROFILE DESCRIPTION FORM

[To be developed.]

PROFILE DESCRIPTION EXAMPLE

[To be developed.]

PROFILE DESCRIPTION REPORT EXAMPLE (for Soil Survey Reports)

[To be developed.]

REFERENCES

- Amoozegar, A. and A.W. Warrick. 1986. Hydraulic conductivity of saturated soils: field methods. *In*: Klute, A. (ed). 1986. Methods of soil analysis: Part 1, Physical and mineralogical methods, 2nd ed. American Society of Agronomy, Agronomy Monograph No. 9, Madison, WI.
- ASSHTO. 1986a. Recommended practice for the classification of soils and soil-aggregate mixtures for highway construction purposes. ASSHTO Designation: M145-82. *In*: Standard specifications for transportation materials and methods of sampling and testing; Part 1 - Specifications (14th ed.). American Association of State Highway and Transportation Officials, Washington, D.C.
- ASSHTO. 1986b. Standard definitions of terms relating to subgrade, soil-aggregate, and fill materials. ASSHTO Designation: M146-70 (1980). *In*: Standard specifications for transportation materials and methods of sampling and testing; Part 1 - Specifications (14th ed.). American Association of State Highway and Transportation Officials, Washington, D.C.
- ASTM. 1993. Standard classification of soils for engineering purposes (Unified Soil Classification System). ASTM designation: D2487-92. *In*: Soil and rock; dimension stone; geosynthetics. Annual book of ASTM standards - Vol. 04.08.
- Bates, R.L., and Jackson, J.A. (eds). 1987. Glossary of Geology, 3rd Ed. American Geological Institute, Alexandria, VA. 788 pp.
- Bouma, J., Paetzold, R.F., and Grossman, R.B. 1982. Measuring hydraulic conductivity for use in soil survey. Soil Survey Investigations Report No. 38. USDA - Soil Conservation Service, U.S. Gov. Print. Office, Washington, D.C. 14 pp.
- Brewer, R. 1976. Fabric and mineral analysis of soils. Krieger Publishing Co., Huntington, NY. 482 pp.

- Bullock, P., Fedoroff, N., Jongerius, A., Stoops, G., Tursina, T. 1985. Handbook for soil thin section description. Waine Research Publications, Wolverhampton, England. 152 pp.
- Childs, C.W. 1981. Field tests for ferrous iron and ferric-organic complexes (on exchange sites or in water-soluble forms) in soils. *Australian Journal of Soil Research*. 19:175-180.
- Cruden, D.M., and Varnes, D.J. 1996. Landslide types and processes. *In*: Turner, A.K., and Schuster, R.L., eds. Landslides investigation and mitigation. Special Report 247, Transportation Research Board, National Research Council, National Academy Press, Washington, D.C. 675 pp.
- Guthrie, R.L. and Witty, J.E. 1982. New designations for soil horizons and layers and the new Soil Survey Manual. *Soil Science Society America Journal*. 46:443-444.
- Folk, R.L. 1955. Student operator error in determination of roundness, sphericity and grain size. *Journal of Sedimentary Petrology*. 25:297-301.
- Ingram, R.L. 1982. Modified Wentworth scale. *In*: Grain-size scales. AGI Data Sheet 29.1. *In*: Dutro, J.T., Dietrich, R.V., and Foose, R.M. 1989. AGI data sheets for geology in the field, laboratory, and office, 3rd edition. American Geological Institute, Washington, D.C.
- International Soil Science Society. 1993. *In*: Soil Survey Manual. Soil Survey Staff, USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C. 503 pp.
- Lynn, W., and D. Williams. 1992. The making of a Vertisol. *Soil Survey Horizons*. 33:23-52.
- National Institute of Standards and Technology. 1990. Counties and equivalent entities of the United States, it's possessions and associated areas. U.S. Dept. Commerce, Federal Information Processing Standards Publication (FIPS PUB 6-4).
- Natural Resources Conservation Service. 1996b. The national PLANTS database. USDA - National Plant Data Center, Baton Rouge, LA. (electronic database).

- O'Neil, A.M. 1952. A key for evaluating soil permeability by means of certain field clues. *Soil Science Society America Proceedings*. 16:312-315.
- Peterson, F.F. 1981. Landforms of the basin and range province: Defined for soil survey. *Nevada Agricultural Experiment Station Technical Bulletin 28*, University of Nevada - Reno, Reno, NV. 52 pp.
- Powers, M.C. 1953. A new roundness scale for sedimentary particles. *Journal of Sedimentary Petrology*. 23:117-119.
- Public Building Service. Sept. 1996. Worldwide geographic location codes. U.S. General Services Administration, Washington, D.C.
- Ruhe, R.V. 1975. *Geomorphology: geomorphic processes and surficial geology*. Houghton-Mifflin Co., Boston, MA. 246 pp.
- Schoeneberger, P.J., Wysocki, D.A. 1996. Geomorphic descriptors for landforms and geomorphic components: effective models, weaknesses and gaps. [Abstract]. *American Society of Agronomy, Annual Meetings, Indianapolis, IN*.
- Soil Conservation Service. 1981. *Land Resource Regions and Major Land Resource Areas of the United States*. USDA Agricultural Handbook 296. U.S. Gov. Print. Office, Washington, D.C.
- Soil Survey Staff. 1951. *Soil Survey Manual*. USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C. 437 pp.
- Soil Survey Staff. 1962. Supplement to Agricultural Handbook No.18, *Soil Survey Manual* (replacing pages 173-188). USDA - Soil Conservation Service, U.S. Gov. Print. Office, Washington, D.C.
- Soil Survey Staff. 1983. *National Soil Survey Handbook, Part 603*, p.45. USDA - Soil Conservation Service, U.S. Gov. Print. Office, Washington, D.C.
- Soil Survey Staff. 1993. *Soil Survey Manual*. USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C. 503 pp.
- Soil Survey Staff. 1995. *Soil survey laboratory information manual*. USDA - Natural Resources Conservation Service, Soil Survey Investigations Report No. 45, Version 1.0, National Soil Survey Center, Lincoln, NE. 305 pp.

Soil Survey Staff. 1996a. Data Dictionary. *In*: National Soils Information System (NASIS), Release 3.0. USDA - Natural Resource Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 1996b. Keys to Soil Taxonomy, 7th ed. USDA - Soil Conservation Service, U.S. Gov. Print. Office, Washington, D.C. 644 pp.

Soil Survey Staff. 1996c. National Soil Survey Handbook. USDA - Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

Soil Survey Staff. 1996d. Pedon Description Program, version 4 design documents. National Soil Survey Center, Lincoln, NE. (unpublished)

Vepraskas, M.J. 1992. Redoximorphic features for identifying aquic conditions. North Carolina Agricultural Research Service Technical Bulletin 301, North Carolina State University, Raleigh, NC. 33 pp.

GEOMORPHIC DESCRIPTION

GEOMORPHIC DESCRIPTION SYSTEM

(Version 2.06 - 9/4/97)

P.J. Schoeneberger, D.A. Wysocki, NRCS, Lincoln, NE

PART I: PHYSIOGRAPHIC LOCATION

- A) Physiographic Division
- B) Physiographic Province
- C) Physiographic Section
- D) State Physiographic Area
- E) Local Physiographic / Geographic Name

PART II: GEOMORPHIC DESCRIPTION

- A) Landscape
- B) Landform
- C) Microfeature
- D) Anthropogenic Features

PART III: SURFACE MORPHOMETRY

- A) Elevation
- B) Slope Aspect
- C) Slope Gradient
- D) Slope Complexity
- E) Slope Shape
- F) Hillslope - Profile Position
- G) Geomorphic Component
 - 1. Hills
 - 2. Terraces
 - 3. Mountains
 - 4. Flat Plains (Proposed)
- H) Microrelief

NOTE: Italicized NASIS short-codes, if available, follow each choice.

PART I: PHYSIOGRAPHIC LOCATION

Reference: **A, B, & C** see Fenneman's 1946 map (reprinted 1957), and Wahrhaftig, 1965.

Physiographic Divisions (A)	Physiographic Provinces (B)	Physiographic Sections (C)
Laurentian Upland	<i>LU</i>	1. Superior Upland <i>SU</i>
Atlantic Lowland	<i>AL</i>	2. Continental Shelf <i>CS</i>
		3. Coastal Plain <i>CP</i>
		a. Embayed section <i>EMS</i>
		b. Sea Island section <i>SIS</i>
		c. Floridian section <i>FLS</i>
		d. East Gulf Coastal plain <i>EGC</i>
		e. Mississippi alluvial valley <i>MAV</i>
		f. West Gulf Coastal plain <i>WGC</i>
Appalachian Highlands	<i>AH</i>	4. Piedmont Province <i>PP</i>
		a. Piedmont upland <i>PIU</i>
		b. Piedmont lowlands <i>PIL</i>
		5. Blue Ridge Province <i>BR</i>
		a. Northern section <i>NOS</i>
		b. Southern section <i>SOS</i>
		6. Valley and Ridge Province <i>VR</i>
		a. Tennessee section <i>TNS</i>
		b. Middle section <i>MIS</i>
		c. Hudson Valley <i>HUV</i>
		7. St. Lawrence Valley <i>SL</i>
		a. Champlain section <i>CHS</i>
		b. St. Lawrence Valley, - northern section <i>NRS</i>
		8. Appalachian Plateau <i>AP</i>
		a. Mohawk section <i>MOS</i>
		b. Catskill section <i>CAS</i>
		c. Southern New York sect. <i>SNY</i>
		d. Allegheny Mountain sect. <i>AMS</i>
		e. Kanawaha section <i>KAS</i>

		f. Cumberland Plateau sect.	<i>CPS</i>
		g. Cumberland Mountain sect.	<i>CMS</i>
		9. New England Province	<i>NE</i>
		a. Seaboard lowland sect.	<i>SLS</i>
		b. New England upland sect.	<i>NEU</i>
		c. White Mountain section	<i>WMS</i>
		d. Green Mountain section	<i>GMS</i>
		e. Taconic section	<i>TAS</i>
		10. Adirondack Province	<i>AD</i>
Interior Plains	<i>IN</i>	11. Interior Low Plateaus	<i>IL</i>
		a. Highland rim section	<i>HRS</i>
		b. Lexington lowland	<i>LEL</i>
		c. Nashville basin	<i>NAB</i>
		d. Possible western section (not delimited on map)	<i>WES</i>
		12. Central Lowland Province	<i>CL</i>
		a. Eastern lake section	<i>ELS</i>
		b. Western lake section	<i>WLS</i>
		c. Wisconsin driftless section	<i>WDS</i>
		d. Till plains	<i>TIP</i>
		e. Dissected till plains	<i>DTP</i>
		f. Osage plain	<i>OSP</i>
		13. Great Plains Province	<i>GP</i>
		a. Missouri plateau, glaciated	<i>MPG</i>
		b. Missouri plateau, unglaciated	<i>MPU</i>
		c. Black Hills	<i>BLH</i>
		d. High Plains	<i>HIP</i>
		e. Plains Border	<i>PLB</i>
		f. Colorado Piedmont	<i>COP</i>
		g. Raton section	<i>RAS</i>
		h. Pecos valley	<i>PEV</i>
		i. Edwards Plateau	<i>EDP</i>
		k. Central Texas section	<i>CTS</i>

This division includes portions of Alaska
(see "Alaskan Physiographic Areas")

Interior Highlands	<i>IH</i>	14. Ozark Plateau	<i>OP</i>
		a. Springfield-Salem plateaus	<i>SSP</i>
		b. Boston "Mountains"	<i>BOM</i>

		15. Ouachita Province	<i>OU</i>
		a. Arkansas Valley	<i>ARV</i>
		b. Ouachita Mountains	<i>OUM</i>
Rocky Mountain System	<i>RM</i>	16. Southern Rocky Mountains	<i>SR</i>
		17. Wyoming Basin	<i>WB</i>
		18. Middle Rocky Mountains	<i>MR</i>
		19. Northern Rocky Mountains	<i>NR</i>

This division includes portions of Alaska
(see "Alaskan Physiographic Areas")

Intermontane Plateaus	<i>IP</i>	20. Columbia Plateau	<i>CR</i>
		a. Walla Walla Plateau	<i>WWP</i>
		b. Blue Mountain section	<i>BMS</i>
		c. Payette section	<i>PAS</i>
		d. Snake River Plain	<i>SRP</i>
		e. Harney section	<i>HAS</i>
		21. Colorado Plateau	<i>CO</i>
		a. High Plateaus of Utah	<i>HPU</i>
		b. Uinta Basin	<i>UIB</i>
		c. Canyon Lands	<i>CAL</i>
		d. Navajo section	<i>NAS</i>
		e. Grand Canyon section	<i>GCS</i>
		f. Datil section	<i>DAS</i>
		22. Basin and Range Province	<i>BP</i>
		a. Great Basin	<i>GRB</i>
		b. Sonoran Desert	<i>SOD</i>
		c. Salton Trough	<i>SAT</i>
		d. Mexican Highland	<i>MEH</i>
		e. Sacramento section	<i>SAS</i>

This division includes portions of Alaska
(see "Alaskan Physiographic Areas")

Pacific Mountain	<i>PM</i>	23. Cascade-Sierra Mountains	<i>CM</i>
		a. Northern Cascade Mtns.	<i>NCM</i>
		b. Middle Cascade Mtns.	<i>MCM</i>
		c. Southern Cascade Mtns.	<i>SCM</i>
		d. Sierra Nevada	<i>SIN</i>

- 24. Pacific Border Province *PB*
 - a. Puget Trough *PUT*
 - b. Olympic Mountains *OLM*
 - c. Oregon Coast Range *OCR*
 - d. Klamath Mountains *KLM*
 - e. California Trough *CAT*
 - f. California Coast Ranges *CCR*
 - g. Los Angeles Ranges *LAR*

- 25. Lower California Province *LC*

This division includes portions of Alaska
(see "Alaskan Physiographic Areas")

Alaskan Physiographic Areas (Wahrhaftig, 1965)

The following Alaskan-Peninsula physiographic areas are extensions of the previous North American Physiographic Divisions (e.g., Rocky Mountain System). These extensions are presented separately, rather than intermingled with the previous Division / Province lists because they: a) constitute a geographically coherent package (Wahrhaftig, 1965); b) these extensions were not contained within Fenneman's original work which dealt only with the conterminous U.S. (Fenneman, 1931; 1938; & 1946); and c) Wahrhaftig's map-unit numbers are independent of, and inconsistent with Fenneman's. Wahrhaftig's original map unit scheme and numbers are retained here for simplicity in using his map of the Alaskan peninsula. **CAUTION:** Wahrhaftig's map unit numbers should not be confused with similar map unit numbers from Fenneman's map for the conterminous U.S.

- | | | | |
|------------------------|-----------|---|-----------|
| Interior Plains | <i>IN</i> | 1. Arctic Coastal Plain Province | -- |
| | | a. Teshekpuk Hills section | -- |
| | | b. White Hills section | -- |
| | | 2. Arctic Foothills Province | <i>AF</i> |
| | | a. Northern Section | -- |
| | | b. Southern Section | -- |
| Rocky Mountains System | <i>RM</i> | Arctic Mountains Province | <i>AM</i> |
| | | 3. Delong Mountains section | -- |
| | | 4. Noatak Lowlands section | -- |
| | | 5. Baird Mountains section | -- |
| | | 6. Central & E. Brooks Range sect. | -- |
| | | 7. Ambler-Chandalar Ridge & Lowland sect. | -- |

NOTE: The map-unit numbering sequence shown here is from Wahrhaftig (1965), and is independent of, and not consistent with, that of Fenneman.

Intermontane Plateaus	Northern Plateaus Province	--
<i>IP</i>	8. Porcupine Plateau section	--
	a. Thazzik Mountain	
	9. Old Crow Plain section	--
	(noted but not described)	
	10. Olgivie Mountains section	--
	11. Tintina Valley (Eagle Trough) sect.	--
	12. Yukon-Tanana Upland section	--
	a. Western Part	
	b. Eastern Part	
	13. Northway - Tanacross Lowland sect.	--
	14. Yukon Flats section	--
	15. Rampart Trough section	--
	16. Kokrine - Hodzana Highlands sect.	--
	a. Ray Mountains	
	b. Kokrine Mountains	
	Western Alaska Province	--
	17. Kanuti Flats section	--
	18. Tozitna - Melozitna Lowland sect.	--
	19. Indian River Upland section	--
	20. Pah River Section	--
	a. Lockwood Hills	
	b. Pah River Flats	
	c. Zane Hills	
	d. Purcell Mountains	
	21. Koyukuk Flats section	--
	22. Kobuk-Selawik Lowland section	--
	a. Waring Mountains	
	23. Selawik Hills section	--
	24. Buckland River Lowland section	--
	25. Nulato Hills section	--
	26. Tanana - Kuskowin Lowland sect.	--
	27. Nowitna Lowland section	--
	28. Kuskokwim Mountains section	--
	29. Innoko Lowlands section	--
	30. Nushagak - Big River Hills section	--
	31. Holitna Lowland section	--
	32. Nushagak-Bristol Bay Lowland sect.	--
	33. Seward Peninsula Province	<i>SEP</i>
	a. Bendeleben Mountains	
	b. Kigluaik Mountains	
	c. York Mountains	

Bering Shelf Province		<i>BES</i>
34. Yukon- Kuskokwim Coastal		
Lowland sect.		--
a. Norton Bay Lowland		
35. Bering Platform section		--
a. St. Lawrence Island		
b. Pribilof Island		
c. St. Matthew Island		
d. Nunivak Island		
36. Ahklun Mountains Province		---

NOTE: The map-unit numbering sequence shown here is from Wahrhaftig (1965), and is independent of, and not consistent with, that of Fenneman.

Pacific Mountain		Alaska - Aleutian Province	<i>AAC</i>
System	<i>PM</i>	37. Aleutian Islands section	--
		38. Aleutian Range section	--
		39. Alaska Range (Southern Part) sect.	--
		40. Alaska Range (Central &	
		Eastern Parts) section	--
		a. Mentasta - Nutzotin Mtn. segment	
		41. Northern Foothills of the Alaska	
		Range section	--
		Coastal Trough Province	--
		42. Cook Inlet - Susitna Lowland sect.	--
		43. Broad Pass Depression section	--
		44. Talkeetna Mountains section	--
		a. Chulitna Mountains	
		b. Fog Lakes Upland	
		c. Central Talkeetna Mountains	
		d. Clarence Lake Upland	
		e. Southeastern Talkeetna Mountains	
		45. Upper Matanuska Valley section	--
		46. Clearwater Mountains section	--
		47. Gulkana Upland section	--
		48. Copper River Lowland section	--
		a. Eastern Part	
		b. Western Part: Lake Louis Plateau	
		49. Wrangell Mountains section	--
		50. Duke Depression (not described)	
		51. Chatham Trough section	--
		52. Kupreanof Lowland section	--
		Pacific Border Ranges Province	<i>PBS</i>
		53. Kodiak Mountains section	--

- 54. Kenai - Chugach Mountains sect. --
- 55. St Elias Mountains section --
 - a. Fairweather Range subsection
- 56. Gulf of Alaska Coastal section --
- 57. Chilkat - Baranof Mountains sect. --
 - a. Alesek Ranges subsection
 - b. Glacier Bay subsection
 - c. Chichagof Highland subsection
 - d. Baranof Mountains subsection
- 58. Prince of Whales Mountains sect. --

- Coast Mountains Province *COM*
- 59. Boundary Pass section --
- 60. Coastal Foothills section --

Other Physiographic Areas

(not addressed by Fenneman, 1946; or Wahrhaftig, 1965)

- Pacific Rim *PR* Pacific Islands Province *PI*
 - a. Hawaiian Islands *HAI*
 - b. Guam *GUM*
 - c. Trust territories * *TRT*
 - d. other (?)

* Most of the former U.S. Trust Territories of the Pacific are now independent nations. This designation is used here solely for brevity and to aid in accessing archived, historical data.

- Caribbean Basin *CB* Caribbean Islands Province *CI*
 - a. Greater Antilles (Puerto Rico) *GRA*
 - b. Lesser Antilles (U.S. Virgin Is.) *LEA*
 - c. other (?)

- Undesignated *UN* Other *OT*
(reserved for temporary, or international designations)

State Physiographic Area (E)

(OPTIONAL) (Entries presently undefined; to be developed in conjunction with each State Geological Survey; target scale is approximately 1:100,000.)

Local Physiographic / Geographic Name (F)

(OPTIONAL) (Entries presently undefined; to be developed in conjunction with each State

Geological Survey; may include area names found on USGS 7.5 & 15 minute topographic maps; target scale is approximately 1:24,000.)

Sources:

Fenneman, N.M. 1931. Physiography of the western United States. McGraw-Hill Co., New York, NY. 534 p.

Fenneman, N.M. 1938. Physiography of the eastern United States. McGraw-Hill Co., New York, NY. 714 p.

Fenneman, N.M. 1946 (reprinted 1957). Physical divisions of the United States. U.S. Geological Survey, U.S. Gov. Print. Office, Washington, D.C. 1 sheet; 1:7,000,000.

Wahrhaftig, C. 1965. Physiographic divisions of Alaska. U.S. Geological Survey Professional Paper 482. 52p.

PART II: GEOMORPHIC DESCRIPTION (OUTLINE)

A) Landscape Terms

B) Landform Terms

- i) Alphabetical List (comprehensive, master list)
- ii) Landform Subset Lists (landform terms grouped by "process" or common setting)
 - 1. Beach, Coastal, Marine, and Lacustrine Landforms
 - 2. Depressional Landforms
 - 3. Eolian Landforms
 - 4. Erosional Landforms
 - 5. Fluvial Landforms
 - 6. Glacial Landforms
 - 7. Mass Movement Landforms
 - 8. Periglacial Landforms
 - 9. Solution Landforms
 - 10. Slope Landforms
 - 11. Tectonic, Structural, and Volcanic Landforms
 - 12. Wetland Terms and Landforms
 - 13. Water "Landforms" and Related Terms

C) Microfeature Terms

D) Anthropogenic Terms

NOTE: Italicized NASIS short-codes, if available, follow each choice.

PART II: GEOMORPHIC DESCRIPTION

A) Landscape
(LF = Landform)

badlands	<i>BA</i>	marine terrace (also LF)	--
bajada (also LF)	<i>BJ</i>	meander belt	<i>MB</i>
basin	<i>BS</i>	mountains (singular = LF)	<i>MO</i>
bolson	<i>BO</i>	piedmont	<i>PI</i>
breaks	<i>BK</i>	plains (also LF)	<i>PL</i>
canyonlands	--	plateau (also LF)	<i>PT</i>
coastal plain (also LF)	<i>CP</i>	river valley	<i>RV</i>
delta plain (also LF)	--	sandhills	<i>SH</i>
drumlin field	--	sand plain	--
dune field	--	scabland	<i>SC</i>
fan piedmont (also LF)	<i>FP</i>	semi-bolson	<i>SB</i>
foothills	<i>FH</i>	shore complex	--
hills (singular = LF)	<i>HI</i>	tableland	<i>TB</i>
intermontane basin (also LF)	<i>IB</i>	thermokarst	<i>TK</i>
island (also LF)	--	till plain (also LF)	<i>TP</i>
karstland	<i>KP</i>	upland	<i>UP</i>
lava plateau (also LF)	<i>LL</i>	valley (also LF)	<i>VA</i>

B) Landform

(LS = Landscape; micro = microfeature; w = water body. Italicized NASIS code follows each term.)

i) Alphabetical Landform List

a'a lava flow	--	ballon	<i>BV</i>
alas	<i>AA</i>	bar	<i>BR</i>
alluvial fan	<i>AF</i>	barchan dune	<i>BQ</i>
alluvial flat	<i>AP</i>	barrier beach	<i>BB</i>
anticline	<i>AN</i>	barrier flat	<i>BP</i>
arete	<i>AR</i>	barrier island	<i>BI</i>
arroyo	<i>AY</i>	basin floor	<i>BC</i>
ash flow (also material)	<i>AS</i>	basin-floor remnant	<i>BD</i>
atoll	<i>AT</i>	bay (w)	<i>WB</i>
avalanche chute	<i>AL</i>	bayou (w)	<i>WC</i>
backshore	<i>AZ</i>	beach	<i>BE</i>
backswamp	<i>BS</i>	beach plain	<i>BP</i>
bajada (also LS)	<i>BJ</i>	beach ridge	<i>BG</i>
ballena	<i>BL</i>	beach terrace	<i>BT</i>

berm	<i>BM</i>	drainageway	<i>DQ</i>
blind valley	<i>VB</i>	draw	<i>DW</i>
block field (also material)	<i>BW</i>	drumlin	<i>DR</i>
block glide (also material)	--	dune	<i>DU</i>
block stream (also material)	<i>BX</i>	earth flow (also material)	<i>EF</i>
blowout	<i>BY</i>	end moraine	<i>EM</i>
bluff	<i>BN</i>	ephemeral stream	
bog (also wetland)	<i>BO</i>	(also micro)	--
braided stream	<i>BZ</i>	erosion remnant	<i>ER</i>
butte	<i>BU</i>	escarpment	<i>ES</i>
caldera	<i>CD</i>	esker	<i>EK</i>
canyon	<i>CA</i>	estuary (w)	<i>WD</i>
Carolina Bay	<i>CB</i>	faceted spur	<i>FS</i>
channel (also micro)	<i>CC</i>	fall (also material)	<i>FB</i>
chenier	<i>CG</i>	fan	<i>FC</i>
chenier plain	<i>CH</i>	fan apron	<i>FA</i>
cinder cone	<i>CI</i>	fanhead trench	<i>FF</i>
cirque	<i>CQ</i>	fan piedmont (also LS)	<i>FG</i>
cliff	<i>CJ</i>	fan remnant	<i>FH</i>
coastal plain (also LS)	<i>CP</i>	fan skirt	<i>FI</i>
col	<i>CL</i>	fault-line scarp	<i>FK</i>
collapsed ice-floored lakebed	<i>CK</i>	fen	<i>FN</i>
collapsed ice-walled lakebed	<i>CN</i>	fjord (w)	<i>FJ</i>
collapsed lake plain	<i>CS</i>	flat	<i>FL</i>
collapsed outwash plain	<i>CT</i>	flood plain	<i>FP</i>
complex landslide	--	flood-plain playa	<i>FY</i>
coulee	<i>CE</i>	flood-plain splay	<i>FM</i>
cove	<i>CO</i>	flood-plain step	<i>FO</i>
crater (volcanic)	<i>CR</i>	flute	<i>FU</i>
crevasse filling	<i>CF</i>	fold (also structure)	<i>FQ</i>
cuesta	<i>CU</i>	foredune	<i>FD</i>
cutoff	<i>CV</i>	fosse	<i>FV</i>
debris avalanche		free face	<i>FW</i>
(also material)	<i>DA</i>	gap	<i>GA</i>
debris flow (also material)	<i>DF</i>	giant ripple	<i>GC</i>
debris slide (also material)	--	glacial drainage channel	<i>GD</i>
deflation basin	<i>DB</i>	glacial lake (w)	<i>WE</i>
delta	<i>DE</i>	glacial lake (relict)	<i>GL</i>
delta plain (also LS)	<i>DC</i>	gorge	<i>GO</i>
depression	<i>DP</i>	graben	<i>GR</i>
diapir	<i>DD</i>	ground moraine	<i>GM</i>
dike	<i>DK</i>	gulch	<i>GT</i>
dipslope	<i>DL</i>	gut (channel); (w)	<i>WH</i>
disintegration moraine	<i>DM</i>	gut (valley)	<i>GV</i>
divide	<i>DN</i>	hanging valley	<i>HV</i>
dome	<i>DO</i>	headland	<i>HE</i>

headwall	<i>HW</i>	meander	<i>MB</i>
highmoor bog	<i>HB</i>	meandering channel	<i>MC</i>
hill	<i>HI</i>	meander scar	<i>MS</i>
hogback	<i>HO</i>	meander scroll	<i>MG</i>
horn	<i>HR</i>	medial moraine	<i>MH</i>
horst	<i>HT</i>	mesa	<i>ME</i>
inselberg	<i>IN</i>	monadnock	<i>MD</i>
inset fan	<i>IF</i>	monocline (also structure)	<i>MJ</i>
interdune	<i>ID</i>	moraine	<i>MU</i>
interfluve (also Geom. Component - Hills)	<i>IV</i>	mountain (also LS)	<i>MM</i>
intermittent stream		mountain slope	<i>MN</i>
(also micro)	--	mountain valley	<i>MV</i>
intermontane basin (also LS)	<i>IB</i>	mud flat	<i>MF</i>
island (also LS)	--	mudflow (also material)	<i>MW</i>
kame	<i>KA</i>	muskeg	<i>MX</i>
kame moraine	<i>KM</i>	natural levee	<i>NL</i>
kame terrace	<i>KT</i>	notch	<i>NO</i>
kettle	<i>KE</i>	nunatak	<i>NU</i>
kipuka	--	outwash fan	<i>OF</i>
knob	<i>KN</i>	outwash plain	<i>OP</i>
knoll	<i>KL</i>	outwash terrace	<i>OT</i>
lagoon (w)	<i>WI</i>	overflow stream (channel)	--
lahar (also material)	<i>LA</i>	oxbow	<i>OX</i>
lake (w)	<i>WJ</i>	oxbow lake (w)	<i>WK</i>
lakebed (relict)	<i>LB</i>	oxbow lake (ephemeral)	<i>OL</i>
lake plain	<i>LP</i>	paha	<i>PA</i>
lakeshore	<i>LF</i>	pahoehoe lava flow	--
lake terrace	<i>LT</i>	paleoterrace (or relict terrace)	--
landslide (also material)	<i>LK</i>	parabolic dune	<i>PB</i>
lateral moraine	<i>LM</i>	parna dune	<i>PD</i>
lateral spread (also material)	--	partial ballena	<i>PF</i>
lava flow	<i>LC</i>	patterned ground	<i>PG</i>
lava plain	<i>LN</i>	peak	<i>PK</i>
lava plateau (also LS)	<i>LL</i>	peat plateau	<i>PJ</i>
lava tube	--	pediment	<i>PE</i>
ledge	<i>LE</i>	perennial stream (w)	--
levee (stream)	<i>LV</i>	pingo	<i>PI</i>
loess bluff	<i>LO</i>	pitted outwash plain	<i>PM</i>
loess hill	<i>LQ</i>	pitted outwash terrace	--
longshore bar [relict]	<i>LR</i>	plain (also LS)	<i>PN</i>
louderback (also structure)	<i>LU</i>	plateau (also LS)	<i>PT</i>
lowmoor bog	<i>LX</i>	playa	<i>PL</i>
marine terrace	<i>MT</i>	playa lake (w)	<i>WL</i>
marsh	<i>MA</i>	plug dome	<i>PP</i>
mawae	--	pluvial lake (w)	<i>WM</i>
		pluvial lake (relict)	<i>PQ</i>

pocosin	<i>PO</i>	soil fall	--
point bar	<i>PR</i>	spit	<i>SP</i>
pothole (also micro)	<i>PH</i>	spur	<i>SQ</i>
pothole lake (w)	<i>WN</i>	stack	<i>SR</i>
pressure ridge (volc)	<i>PU</i>	steptoe	<i>ST</i>
proglacial lake (w)	<i>WO</i>	strand plain	<i>SS</i>
proglacial lake (relict)	--	strath terrace	<i>SU</i>
raised beach	<i>RA</i>	stratovolcano	<i>SV</i>
raised bog	<i>RB</i>	stream (w)	--
ravine	<i>RV</i>	stream terrace	<i>SX</i>
recessional moraine	<i>RM</i>	string bog	<i>SY</i>
reef	<i>RF</i>	structural bench	<i>SB</i>
ribbed fen	<i>RG</i>	swale (also micro)	<i>SC</i>
ridge	<i>RI</i>	swallow hole	<i>TB</i>
rim	<i>RJ</i>	swamp	<i>SW</i>
rise	--	syncline (also structure)	<i>SZ</i>
river (w)	--	talus slope	--
roche moutonnee	<i>RN</i>	terminal moraine	<i>TA</i>
rock fall (also micro)	--	terrace	<i>TE</i>
rock avalanche (also material)	--	thermokarst depression	<i>TK</i>
rock glacier	<i>RO</i>	thermokarst lake (w)	<i>WV</i>
rotational landslide		tidal flat	<i>TF</i>
(also material)	<i>RP</i>	till plain (also LS)	<i>TP</i>
saddle	<i>SA</i>	tombolo	<i>TO</i>
salt marsh	<i>SM</i>	topple	--
salt pond (w)	<i>WQ</i>	tor	<i>TQ</i>
sand flow (also material)	<i>RW</i>	translational slide	<i>TS</i>
sand sheet	<i>RX</i>	transverse dune	<i>TD</i>
scarp	<i>RY</i>	trough	<i>TR</i>
scarp slope	<i>RS</i>	tunnel valley	<i>TV</i>
scree slope	--	U-shaped valley	<i>UV</i>
sea cliff	<i>RZ</i>	valley	<i>VA</i>
seif dune	<i>SD</i>	valley flat	<i>VF</i>
shield volcano	--	valley floor	<i>VL</i>
shoal (w)	<i>WR</i>	valley side	<i>VS</i>
shoal (relict)	<i>SE</i>	valley train	<i>VT</i>
shore	--	volcanic cone	<i>VC</i>
shrub-coppice dune (micro)	<i>SG</i>	volcanic dome	<i>VD</i>
sill	<i>RT</i>	volcano	<i>VO</i>
sinkhole	<i>SH</i>	V-shaped valley	<i>VV</i>
slackwater (w)	<i>WS</i>	wash	<i>WA</i>
slide (also material)	<i>SJ</i>	washover fan	<i>WF</i>
slough (ephemeral water)	<i>SL</i>	wave-built terrace	<i>WT</i>
slough (permanent water)	<i>WU</i>	wave-cut platform	<i>WP</i>
slump	<i>SK</i>	wind gap	<i>WG</i>
slump block	<i>SN</i>	yardang (also micro)	--

yardang trough (also micro) --

ii) Landform Subset Lists (Landform terms grouped by "process" or common setting)

1. Beach, Coastal, Marine, and Lacustrine Landforms

atoll	AT	lakebed (relict)	LB
backshore	AZ	lake plain	LP
bar	BR	lake terrace	LT
barrier beach	BB	longshore bar [relict]	LR
barrier flat	BF	marine terrace (also LS)	MT
barrier island	BI	mud flat	MF
beach	BE	playa	PL
beach plain	BP	pluvial lake (relict)	PQ
beach ridge	BG	raised beach	RA
beach terrace	BT	reef	RF
berm	BM	salt marsh	SM
bluff	BN	sea cliff	RZ
chenier	CG	shoal (relict)	SE
chenier plain	CH	shore	--
coastal plain	CP	spit	SP
delta	DE	stack	SR
delta plain (also LS)	DC	strand plain	SS
flat	FL	tidal flat	TF
foredune	FD	tombolo	TO
headland	HE	washover fan	WF
island (also LS)	--	wave-built terrace	WT
lagoon	WI	wave-cut platform	WP

2. Depressional Landforms

alluvial flat	AP	gulch	GT
basin floor	BC	gut (valley)	GV
basin floor remnant	BD	interdune	ID
canyon	CA	intermontane basin	IB
Carolina Bay	CB	kettle	KE
col	CL	mountain valley	MV
coulee	CE	playa	PL
cove	CO	pothole (also micro)	PH
depression	DP	ravine	RV
drainageway	DQ	saddle	SA
gap	GA	slough (ephemeral)	SL
gorge	GO	swale (also micro)	SC

trough	<i>TR</i>	valley floor	<i>VL</i>
U-shaped valley	<i>UV</i>	V-shaped valley	<i>VV</i>
valley	<i>VA</i>		

3. Eolian Landforms

barchan dune	<i>BQ</i>	loess hill	<i>LQ</i>
blowout	<i>BY</i>	paha	<i>PA</i>
deflation basin	<i>DB</i>	parabolic dune	<i>PB</i>
dune	<i>DU</i>	parna dune	<i>PD</i>
foredune	<i>FD</i>	sand sheet	<i>RX</i>
interdune	<i>ID</i>	seif dune	<i>SD</i>
loess bluff	<i>LO</i>	transverse dune	<i>TD</i>

4. Erosional Landforms - Water erosion (overland flow) related and excluding fluvial, glaciofluvial, and eolian erosion.

arete	<i>AR</i>	monadnock	<i>MD</i>
ballena	<i>BL</i>	notch	<i>NO</i>
ballon	<i>BV</i>	paha	<i>PA</i>
basin floor remnant	<i>BD</i>	partial ballena	<i>PF</i>
col	<i>CL</i>	peak	<i>PK</i>
cuesta	<i>CU</i>	pediment	<i>PE</i>
erosion remnant	<i>ER</i>	saddle	<i>SA</i>
free face	<i>FW</i>	scarp slope	<i>RS</i>
gap	<i>GA</i>	strath terrace	<i>SU</i>
hogback	<i>HO</i>	structural bench	<i>SB</i>
horn	<i>HR</i>	tor	<i>TQ</i>
inselberg	<i>IN</i>	wind gap	<i>WG</i>
meander scar	<i>MS</i>		

5. Fluvial Landforms - Dominantly related to concentrated water (channel flow), both erosional and depositional processes, and excluding glaciofluvial landforms.

alluvial fan	<i>AF</i>	braided stream	<i>BZ</i>
alluvial flat	<i>AP</i>	canyon	<i>CA</i>
arroyo	<i>AY</i>	channel	<i>CC</i>
backswamp	<i>BS</i>	coulee	<i>CE</i>
bajada	<i>BJ</i>	cutoff	<i>CV</i>
bar	<i>BR</i>	delta	<i>DE</i>
basin-floor remnant	<i>BD</i>	delta plain (also LS)	<i>DC</i>
block stream	<i>BX</i>	drainageway	<i>DQ</i>

draw	<i>DW</i>	meander scar	<i>MS</i>
fanhead trench	<i>FF</i>	meander scroll	<i>MG</i>
fan skirt	<i>FJ</i>	natural levee	<i>NL</i>
flood plain	<i>FP</i>	overflow stream (channel)	--
flood-plain playa	<i>FY</i>	oxbow	<i>OX</i>
flood-plain splay	<i>FM</i>	oxbow lake (ephemeral)	<i>OL</i>
flood-plain step	<i>FO</i>	pediment	<i>PE</i>
giant ripple	<i>GC</i>	point bar	<i>PR</i>
gorge	<i>GO</i>	ravine	<i>RV</i>
gulch	<i>GT</i>	strath terrace	<i>SU</i>
gut (valley)	<i>GV</i>	stream terrace	<i>SX</i>
inset fan	<i>IF</i>	wash	<i>WA</i>
levee (stream)	<i>LV</i>	wind gap	<i>WG</i>

6. Glacial Landforms (including glaciofluvial forms)

arete	<i>AR</i>	kame moraine	<i>KM</i>
cirque	<i>CQ</i>	kame terrace	<i>KT</i>
col	<i>CL</i>	kettle	<i>KE</i>
collapsed ice-floored lakebed	<i>CK</i>	lateral moraine	<i>LM</i>
collapsed ice-walled lakebed	<i>CN</i>	medial moraine	<i>MH</i>
collapsed lake plain	<i>CS</i>	moraine	<i>MU</i>
collapsed outwash plain	<i>CT</i>	nunatak	<i>NU</i>
crevasse filling	<i>CF</i>	outwash fan	<i>OF</i>
disintegration moraine	<i>DM</i>	outwash plain	<i>OP</i>
drumlin	<i>DR</i>	outwash terrace	<i>OT</i>
end moraine	<i>EM</i>	paha	<i>PA</i>
esker	<i>EK</i>	pitted outwash plain	<i>PM</i>
fjord (w)	<i>FJ</i>	pitted outwash terrace	--
flute	<i>FU</i>	pressure ridge (ice)	--
fosse	<i>FV</i>	proglacial lake (relict)	--
giant ripple	<i>GC</i>	recessional moraine	<i>RM</i>
glacial drainage channel	<i>GD</i>	roche moutonnee	<i>RN</i>
glacial lake (relict)	<i>GL</i>	rock glacier	<i>RO</i>
ground moraine	<i>GM</i>	terminal moraine	<i>TA</i>
hanging valley	<i>HV</i>	till plain (also LS)	<i>TP</i>
kame	<i>KA</i>	tunnel valley	<i>TV</i>
		U - shaped valley	<i>UV</i>

7. Mass Movement Landforms (including creep forms)

ash flow	AS	rock fall (also micro)	--
avalanche chute	AL	rockfall avalanche	--
block glide	--	rock glacier	RO
complex slide	--	rotational landslide	RP
debris avalanche	--	sand flow	RW
debris flow	DF	slide	SJ
debris slide	--	slump	SK
earth flow	EF	slump block	SN
fall	FB	soil fall	--
lahar	LA	talus	--
landslide	LK	topple	--
lateral spread	--	translational slide	TS
mudflow	MW		

8. Periglacial Landforms (modern, relict, and patterned ground)

alas	AA	peat plateau	PJ
block field	BW	pingo	PI
muskeg	MX	rock glacier	RO
patterned ground		string bog	SY
(see Microfeatures)	PG	thermokarst depression	TK

9. Solution Landforms

blind valley	VB	swallow hole	TB
sinkhole	SH	thermokarst depression	TK

10. Slope Landforms - Terms that tend to be generic and that emphasize their form rather than any particular genesis or process.

bluff	BN	headwall	HW
butte	BU	hill (plural = LS)	HI
cliff	CJ	hogback	HO
cuesta	CU	horn	HR
dome	DO	horst	HT
escarpment	ES	inselberg	IN
faceted spur	FS	interfluvium (also Geom.	
fault-line scarp	FK	Component - Hills)	IV
free face	FW	knob	KN
gap	GA	knoll	KL

lahar	<i>LA</i>	plain (also LS)	<i>PN</i>
ledge	<i>LE</i>	plateau (also LS)	<i>PT</i>
meander scar	<i>MS</i>	ridge	<i>RJ</i>
mesa	<i>ME</i>	rim	<i>RJ</i>
monadnock	<i>MD</i>	scarp	<i>RY</i>
mountain	<i>MM</i>	spur	<i>SQ</i>
mountain slope	<i>MN</i>	structural bench	<i>SB</i>
mountain valley	<i>MV</i>	tor	<i>TQ</i>
notch	<i>NO</i>	U - shaped valley	<i>UV</i>
paha	<i>PA</i>	V - shaped valley	<i>VV</i>
peak	<i>PK</i>	wind gap	<i>WG</i>

11. Tectonic, Structural, and Volcanic Landforms

a'a lava flow	--	lava plain	<i>LN</i>
anticline	<i>AN</i>	lava plateau (also LS)	<i>LL</i>
caldera	<i>CD</i>	lava tube	--
cinder cone	<i>CI</i>	louderback	<i>LU</i>
crater (volcanic)	<i>CR</i>	mawae	--
cuesta	<i>CU</i>	monocline	<i>MJ</i>
diapir	<i>DD</i>	pahoehoe lava flow	--
dike	<i>DK</i>	plug dome	<i>PP</i>
dipslope	<i>DL</i>	pressure ridge (volcanic)	<i>PU</i>
dome	<i>DO</i>	scarp slope	<i>RS</i>
fault-line scarp	<i>FK</i>	shield volcano	--
graben	<i>GR</i>	steptoe	<i>ST</i>
hogback	<i>HO</i>	stratovolcano	<i>SV</i>
horst	<i>HT</i>	structural bench	<i>SB</i>
kipuka	--	syncline (also structure)	<i>SZ</i>
lahar	<i>LA</i>	volcanic cone	<i>VC</i>
lava flow	<i>LC</i>	volcanic dome	<i>VD</i>

12. Wetland Terms and Landforms (provisional list: conventional, geologic definitions; not legalistic or regulatory usage)

alas	<i>AA</i>	marsh	<i>MA</i>
backswamp	<i>BS</i>	mud flat	<i>MF</i>
bog	<i>BO</i>	muskeg	<i>MX</i>
Carolina Bay	<i>CB</i>	oxbow lake (ephemeral)	<i>OL</i>
estuary	<i>WD</i>	peat plateau	<i>PJ</i>
fen	<i>FN</i>	playa (intermittent water)	<i>PL</i>
highmoor bog	<i>HB</i>	pocosin	<i>PO</i>
lowmoor bog	<i>LX</i>	pothole (intermittent water)	<i>PH</i>

raised bog	RB	string bog	SY
ribbed fen	RG	swamp	SW
salt marsh	SM	tidal flat	TF
slough (intermittent water)	SL		

13. Water "Landforms" and Related Terms - Discrete landform terms but treated generically as "water" in soil survey.

bay	WB	playa lake	WL
bayou	WC	pluvial lake	WM
ephemeral stream (also micro)	--	pond (micro)	--
estuary	WD	pool (micro)	--
fjord	FJ	pothole (lake)	WN
glacial lake	WE	proglacial lake	WO
gut (channel)	WH	river (w)	--
intermittent stream (also micro)	--	salt pond	WQ
lagoon	WI	shoal	WR
lake	WJ	slackwater	WS
oxbow lake	WK	slough (permanent water)	WU
perennial stream (w; also micro)	--	stream (w)	--
		tank (micro)	--
		thermokarst lake	WV

C) Microfeature Terms

bar	--	hoodoo	--
channel (also LF)	--	mound	M:
earth pillar	--	<i>patterned ground</i> microfeatures	
frost boil	--	(see below; used in association	
groove	--	with the landform " <i>patterned</i>	
gully	--	<i>ground</i> " (PG))	
hillock	--		

a) Periglacial *patterned ground* microfeatures:

circle	--	polygons	--
earth hummocks	--	sorted circles	--
high-center polygons	--	stripes	--
ice wedge polygons	--	turf hummocks	--
low-center polygons	--		
non-sorted circles	--		
palsa, palsen (= peat hummocks)	--		

b) Other *patterned ground* microfeatures:

bar and channel	--	mima mounds	--
gilgai	<i>G</i>	pimple mounds	--
hummocks	--		
pinnacle	--	solifluction sheet	--
pond (also water list)	--	solifluction terrace	--
pool (also water list)	--	swale (also LF)	--
pothole (also LF)	--	tank (also water list)	--
rib	--	terraces	<i>T</i>
rill	--	tree-tip mound	--
ripple mark	--	tree-tip pit	--
sand boil	--	yardang (also LF)	--
scour (mark)	--	yardang trough (also LF)	--
solifluction lobe	--		

D) Anthropogenic Features

artificial collapsed depression (e.g., arising from subsurface mining subsidence)	<i>G</i>	pond (human-made)	--
artificial levees	<i>A</i>	quarry	--
borrow pit	--	railroad bed	<i>D</i>
burial mound	<i>B</i>	rice paddy	<i>E</i>
cut (road, railroad)	--	road bed	<i>I</i>
ditch	--	sand pit	--
fill	--	sanitary landfill	--
gravel pit	--	sewage lagoon	--
landfill	--	spoil bank	--
levelled land	--	surface mine	--
midden	<i>H</i>	<i>tillage / management features</i>	<i>F</i>
		(see below for common, more specific types)	


a) *Tillage / management features* (common types):

conservation terrace (modern)	--	drainage ditch	--
double-bedding mound (i.e., bedding mound used for timber; lower Coastal Plain)	--	furrow	--
		hillslope terrace (e.g., archeological features; China, Peru)	--
		inter-furrow	--
urban land	--		

NOTE: Italicized NASIS short-codes, if available, follow each choice.


PART III: SURFACE MORPHOMETRY

- A) **Elevation:** The height of a point on the earth's surface, relative to mean sea level (msl); indicate units; e.g., *106 m* or *348 ft*.
- B) **Slope Aspect:** The compass bearing (in degrees, corrected for declination) that a slope faces, looking downslope; e.g., *287°*.
- C) **Slope Gradient:** The angle of the ground surface (in percent) through the site and in the direction that overland water would flow; e.g., *18%*. (Commonly referred to as slope.)
- D) **Slope Complexity:** Describe the relative uniformity (smooth linear or curvilinear = *simple* or *S*) or irregularity (*complex* or *C*) of the ground surface leading downslope through the point of interest; e.g., *simple* or *S*.




- E) **Slope Shape:** Slope shape is described in two directions: 1) up and down slope (perpendicular (normal) to the contour); and 2) across slope (along the horizontal contour). In PDP, this data element is split into two sequential parts (Slope Across and Slope Up & Down); e.g., *Linear*, *Convex* or *LV*.

Down Slope (Vertical)	Across Slope (Horizontal)	Code	
		PDP3.5	NASIS
Concave	Concave	CC, CC	CC
Concave	Convex	CC, CV	CV
Concave	Linear	CC, LL	CL
Convex	Concave	CV, CC	VC
Convex	Convex	CV, CV	VV
Convex	Linear	CV, LL	VL
Linear	Concave	LL, CC	LC
Linear	Convex	LL, CV	LV
Linear	Linear	LL, LL	LL




- (F) **Hillslope - Profile Position** (Hillslope Position in PDP): Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., *backslope* or *BS*. This is best applied to transects or points, not areas.

Position	Code PDP & NASIS
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS




G) **Geomorphic Component** (Geomorphic Position in PDP): Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) *nose slope* or *NS*.


1) Hills	Code	
	PDP	NASIS
interfluve	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	---	BS



2) Terraces	Code
riser	RI
tread	TR



3) Mountains	Code
mountaintop	MT
mountainflank	MF
upper third mountainflank	UT
center third mountainflank	CT
lower third mountainflank	LT
mountainbase	MB



4) Flat Plains (proposed)	Code
---------------------------	------

H) **Microrelief**: Small, relative differences in elevation between adjacent areas on the earth's surface; e.g., *micro-high* or *MH*; or *micro-low* or *ML*.

NOTE: Microfeature **Kind** and **Pattern** have been deleted from PDP; these phenomena and terms are now captured within the data element "Microfeature".

REFERENCES

Ruhe, R.V. 1975. *Geomorphology: Geomorphic processes and surficial geology*. Houghton-Mifflin, Boston, MA. 246 p.

Schoeneberger, P.J. and Wysocki, D.A. 1996. Geomorphic descriptors for landforms and geomorphic components: effective models and weaknesses. *Agronomy abstracts, American Society of Agronomy Annual Meetings, 1996, Indianapolis, IN.*

Soil Survey Staff. 1998. *Glossary of landforms and geologic materials. Part 629, National Soil Survey Handbook, USDA, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.*

SOIL TAXONOMY

Compiled by: P.J. Schoeneberger, D.A. Wysocki, and E.C. Benham, NRCS,
Lincoln, NE.

INTRODUCTION

The purpose of this section is to expand upon and augment the abbreviated soil taxonomic contents of the "Soil Profile Description Section".

HORIZON NOMENCLATURE

MASTER AND TRANSITIONAL HORIZONS -

Horizon	Criteria ¹
O	Dominated by organic matter (OM); mineral material is a small percent by volume (< 5% by weight), excludes illuvial OM; generally darker than underlying horizons.
A	Mineral soil, formed at surface or below O, no remnant rock structure, and both or either: 1) accumulation of humified organic matter but dominated by mineral matter, and not E or B; or 2) cultivation properties. Excludes recent eolian or alluvial deposits (< 75 cm thick) that retain stratification.
AB (or AE)	Dominantly A horizon characteristics but also has some recognizable characteristics of B (or E) horizon.
A/B (or A/E or A/C)	Discrete, intermingled bodies of two horizons: A and B (or E or C) material; majority of layer is A material.
AC	Dominantly A horizon characteristics but also has some recognizable characteristics of C horizon.
E	Mineral soil, loss of silica, iron, aluminum, or clay leaving a net concentration of sand and silt; no remnant rock structure; typically lighter color (higher value, chroma) and coarser texture than A.
EA (or EB)	Dominantly E horizon characteristics, but also has some recognizable characteristics of A (or B) horizon.
E/A	Discrete, intermingled bodies of two horizons: E and A material; majority of layer is E horizon material.
E and Bt	Presence of thin, heavier textured lamellae (Bt) within a predominantly E horizon with less clay.
BA (or BE)	Dominantly B characteristics but also has some recognizable attributes of A (or E) horizon.

B/A (or B/E)	Discrete, intermingled bodies of two horizons, majority of horizon is B (or E) material.
B	Mineral soil, formed below O, A, or E; little or no rock structure; and one or more of the following: 1) illuvial accumulation of silicate clay, Fe, Al, humus, carbonates, gypsum, or silica (one or more); 2) removal of carbonates; 3) residual accumulation of sesquioxides; 4) sesquioxide coatings; 5) alterations which form silicate clays or liberates oxides and forms pedogenic structure; 6) brittleness (includes any illuvial horizon, cemented or not; and excludes horizons of clay films coating rock fragments or covering finely stratified, unconsolidated sediments; discontinuous carbonate accumulation not contiguous to overlying horizon; and gleyed layers lacking additional pedogenic features).
BC	Dominantly B horizon characteristics but also has some recognizable characteristics of the C horizon.
B/C	Discrete, intermingled bodies of two horizons; majority of horizon is B material.
CB (or CA)	Dominantly C horizon characteristics but also has some recognizable characteristics of the B (or A) horizon.
C/B (or C/A)	Discrete, intermingled bodies of two horizons; majority of horizon is C material.
C	Mineral soil, excludes hard bedrock; layers little affected by pedogenesis and lacks properties of O, A, E, or B horizons. May or may not be related to parent material of the solum.
W	A layer of liquid water (W) or permanently frozen ice (Wf) within the soil (excludes water / ice above soil). ²
R	Hard bedrock.

¹ Soil Survey Staff, 1996.

² NRCS Soil Classification Staff, 1997; personal communication.

HORIZON SUFFIXES -

Horizon Suffixes	Criteria ¹
a	Highly decomposed organic matter (OM); rubbed fiber content < 17% (by vol.); see <i>e, i</i> .
b	Buried genetic horizon (not used with in organic materials or to separate organic from mineral materials).
c	Concretions or nodules; significant accumulation of cemented bodies, enriched with Fe, Al, Mn, Ti [cement not specified except <u>excludes</u> silica (see <i>q</i>)]; not used for calcite, dolomite, or soluble salts (see <i>z</i>).
d	Physical root restriction due to high bulk density (natural or human-made materials / conditions; e.g., lodgement till, plow pans etc.
e	Moderately (intermediately) decomposed organic matter; rubbed fiber content 17-40% (by vol.); see <i>a, i</i> .
f	Permafrost (permanently frozen soil or ice); excludes seasonally frozen ice; continuous subsurface ice.
ff	Dry permafrost [permanently frozen soil; not used for seasonally frozen; no continuous ice bodies (see <i>f</i>)]. ²
g	Strong gley (Fe reduced and pedogenically removed); typically ≤ 2 chroma; may have other redoximorphic (RMF) features; not used for geogenic gray colors.
h	Illuvial organic matter (OM) accumulation (with B: accumulation of illuvial, amorphous OM sesquioxide complexes); coats sand and silt particles or more; use <i>Bhs</i> if significant accumulation of sesquioxides <u>and</u> moist chroma value ≤ 3.
i	Slightly decomposed organic matter; rubbed fiber content > 40% (by vol.); see <i>a, e</i> .
j	Jarosite accumulation ² ; e.g., acid sulfate soils.
jj	Evidence of cryoturbation ² ; e.g., irregular or broken boundaries, sorted rock fragments (patterned ground), or O.M. in lower boundary between active layer and permafrost layer.
k	Pedogenic accumulation of carbonates; e.g. CaCO ₃ .
m	Strong pedogenic cementation or induration (> 90% cemented, even if fractured); physically root restrictive; you can indicate cement type by using letter combinations; e.g., <i>km</i> - carbonates, <i>qm</i> - silica, <i>kqm</i> - carbonates and silica; <i>sm</i> - iron, <i>ym</i> - gypsum; <i>zm</i> - salts more soluble than gypsum.
n	Pedogenic, exchangeable sodium accumulation.
o	Residual accumulation of sesquioxides.

p	Tillage or other disturbance of surface layer (pasture, plow, etc.). Designate <i>Op</i> for disturbed organic surface; <i>Ap</i> for mineral surface even if the layer clearly was originally an E, B, C, etc.
q	Accumulation of secondary (pedogenic) silica.
r	Used with C to indicate weathered or soft bedrock (root restrictive saprolite or soft bedrock; partially consolidated sandstone, siltstone or shale; Excavation Difficulty classes are <i>low to high</i>).
s	Illuvial accumulation of amorphous, dispersible, sesquioxides and organic matter complexes and color value or chroma ≥ 4 . Used with B horizon; used with h as <i>Bhs</i> if color value or chroma is ≤ 3 .
ss	Slickensides; e.g., oblique shear faces 20 - 60° off horizontal; due to shrink-swell clay action; wedge-shaped peds and seasonal surface cracks are also commonly present.
t	Illuvial accumulation of silicate clays (clayskins, lamellae, or clay bridging in some part of the horizon).
v	Plinthite (high Fe, low OM, reddish contents; firm to very firm moist consistence; irreversible hardening with repeated wetting and drying).
w	Incipient color or pedogenic structure development; minimal illuvial accumulations (excluded from use with transition horizons).
x	Fragipan characteristics (brittleness, firmness, bleached prisms).
y	Pedogenic accumulation of gypsum.
z	Pedogenic accumulation of salts more soluble than gypsum; e.g., NaCl, etc.

¹ Soil Survey Staff, 1996

² NRCS Soil Classification Staff, 1998; personal communication.

HORIZON NOMENCLATURE CONVERSION CHARTS -

Master Horizons and Combinations			
1951 ¹	1962 ²	1981 ³	1997 ⁴
---	O	O	O
Aoo	---	(see <i>Oi</i>)	(see <i>Oi</i>)
Ao	O1	Oi and/or Oe	Oi and/or Oe
---	O2	Oe and/or Oa	Oe and/or Oa
A	A	A	A
A1	A1	A	A
A2	A2	E	E
A3	A3	AB or EB	AB or EB
AB	AB	----	----
A&B	A&B	A/B or E/B	A/B or E/B
AC	AC	AC	AC
B	B	B	B
B1	B1	BA or BE	BA or BE
B&A	B&A	B/A or B/E	B/A or B/E
B2	B2	B or Bw	B or Bw
B3	B3	BC or CB	BC or CB
G	---	---	---
Cca	---	---	---
Ccs	---	---	---
---	C	C	C
D	---	---	---
Dr	R	R	R
---	---	---	W

¹ Soil Survey Staff, 1951.

² Soil Survey Staff, 1962.

³ Guthrie and Witty, 1982. Additional changes to lithologic discontinuities.

⁴ NRCS Soil Classification Staff, 1997; personal communication.

Horizon Suffixes (also called "Horizon Subscripts", and "Subordinate Distinctions")			
1951 ¹	1962 ²	1981 ³	1997 ⁴
---	---	a	a
b	b	b	b
ca	ca	(see <i>k</i>)	(see <i>k</i>)
cn	cn	c	c
cs	cs	(see <i>y</i>)	(see <i>y</i>)
---	---	e	e
f	f	f	f
---	---	---	ff
g	g	g	g
h	h	h	h
ir	ir	(see <i>s</i>)	(see <i>s</i>)
---	---	i	i
---	---	---	j
---	---	---	jj
(see <i>ca</i>)	(see <i>ca</i>)	k	k
m	m	m	m
sa	sa	n	n
---	---	o	o
p	p	p	p
(see <i>si</i>)	(see <i>si</i>)	q	q
---	r	r	r
(see <i>ir</i>)	(see <i>ir</i>)	s	s
si	si	(see <i>q</i>)	(see <i>q</i>)
---	---	---	ss (1991)
t	t	t	t
---	---	v	v
---	---	w	w
x	x	x	x
(see <i>cs</i>)	(see <i>cs</i>)	y	y
---	sa	z	z


¹ Soil Survey Staff, 1951.

² Soil Survey Staff, 1962

³ Guthrie and Witty, 1982.


⁴ NRCS Soil Classification Staff, 1997; personal communication.

**Texture Triangle:
Soil Textural Family Classes ()**



¹ Very fine sand (0.05 - 0.1) is treated as silt for family groupings; coarse fragments are considered the equivalent of coarse sand in the boundary between the silty and loamy classes.

**Combined Texture Triangles:
 Fine Earth Texture Classes (—) &
 Soil Textural Family Classes (—)**



¹ Very fine sand (0.05 - 0.1) is treated as silt for family groupings; coarse fragments are considered the equivalent of coarse sand in the boundary between the silty and loamy classes.

REFERENCES

- Guthrie, R.L. and Witty, J.E. 1982. New designations for soil horizons and layers and the new Soil Survey Manual. Soil Science Society America Journal, vol. 46. p.443-444.
- Soil Survey Staff. 1975. Soil Taxonomy: A basic system of soil classification for making and interpreting soil surveys. USDA - Soil Conservation Service, Agricultural Handbook No. 436, U.S. Gov. Print. Office, Washington, D.C. 754 pp.
- Soil Survey Staff. 1962. Identification and nomenclature of soil horizons. USDA - Soil Conservation Service, Supplement to to Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C.
- Soil Survey Staff. 1951. Soil Survey Manual. USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C. 437 pp.
- Soil Survey Staff. 1962. Supplement to Agricultural Handbook No.18, Soil Survey Manual (replacing pages 173-188). USDA - Soil Conservation Service, U.S. Gov. Print. Office, Washington, D.C.
- Soil Survey Staff. 1993. Soil Survey Manual. USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington, D.C. 503 pp.
- Soil Survey Staff. 1996. Keys to Soil Taxonomy, 7th ed. USDA - Soil Conservation Service, U.S. Gov. Print. Office, Washington, D.C. 644 pp.

GEOLOGY

Compiled by: P.J. Schoeneberger, D.A. Wysocki, and E.C. Benham, NRCS,
Lincoln, NE.

INTRODUCTION

The purpose of this section is to expand and augment the geologic information found or needed in the "Site Description Section" and "Soil Profile Description Section".

BEDROCK - KIND

(This table is repeated here from the "Site Selection Section" for convenience in using the following rock charts.)

Kind	Code ¹		Kind	Code ¹	
	PDP	NASIS		PDP	NASIS
IGNEOUS-INTRUSIVE					
diabase	--	DIA	monzonite	--	MON
diorite	--	DIO	peridotite	--	PER
gabbro	--	GAB	pyroxenite	--	PYX
granite	I4	GRA	syenite	--	SYE
granodiorite	--	GRD	syenodiorite	--	SYD
IGNEOUS-EXTRUSIVE					
aa (lava)	P8	AAL	pahoehoe (lava)	P9	PAH
andesite	I7	AND	pumice (flow, coherent)	E6	PUM
basalt	I6	BAS	rhyolite	--	RHY
dacite	--	DAC	scoria (coherent, mass)	E7	SCO
latite	--	LAT	trachyte	--	TRA
obsidian	--	OBS			
IGNEOUS-PYROCLASTIC					
ignimbrite	--	IGN	tuff breccia	P7	TBR
pyroclastics (coherent)	P0	PYR	volcanic breccia	P4	VBR
tuff	P1	TUF	volcanic breccia, acidic	P5	AVB
tuff, acidic	P2	ATU	volcanic breccia, basic	P6	BVB
tuff, basic	P3	BTU			

METAMORPHIC					
amphibolite	--	AMP	metavolcanics	--	MVO
gneiss	M1	GNE	migmatite	--	MIG
granofels	--	GRF	mylonite	--	MYL
granulite	--	GRL	phyllite	--	PHY
greenstone	--	GRE	schist	M5	SCH
hornfels	--	HOR	serpentine	M4	SER
marble	L2	MAR	slate	M8	SLA
metaconglomerate	--	MCN	soapstone (talc)	--	SPS
metaquartzite	M9	MQT			
SEDIMENTARY-CLASTICS					
arenite	--	ARE	porcellanite	--	POR
argillite	--	ARG	sandstone	A0	SST
arkose	A2	ARK	sandstone, calcareous	A4	CSS
breccia, non-volcanic (angular fragments)	--	NBR	shale	H0	SHA
claystone	--	CST	shale, acid	--	ASH
conglomerate (rounded fragments)	C0	CON	shale, calcareous	H2	CSH
conglomerate, calcar.	C2	CCN	shale, clayey	H3	YSH
graywacke	--	GRY	siltstone	T0	SIS
mudstone	--	MUD	siltstone, calcareous	T2	CSI
orthoquartzite	--	OQT			
EVAPORITES, ORGANICS, AND PRECIPITATES					
chalk	L1	CHA	limestone, arenaceous	L5	ALS
chert	--	CHE	limestone, argillaceous	L6	RLS
coal	--	COA	limestone, cherty	L7	CLS
dolostone	L3	DOL	limestone, phosphatic	L4	PLS
gypsum	--	GYP	travertine	--	TRV
limestone	L0	LST	tufa	--	TUA
INTERBEDDED					
limestone-sandst.-shale	B1	LSS	sandstone-shale	B5	SSH
limestone-sandstone	B2	LSA	sandstone-siltstone	B6	SSI
limestone-shale	B3	LSH	shale-siltstone	B7	SHS
limestone-siltstone	B4	LSI			

¹ Definitions for bedrock are found in the "Glossary of Landforms and Geologic Terms", NSSH - Part 629 (Soil Survey Staff, 1996c), and in the "Glossary of Geology" (Bates, et al., 1987).

ROCK CHARTS

The following rock charts (**Igneous, Metamorphic, and Sedimentary & Volcaniclastic**) summarize grain size, composition, or genetic differences between related rock types. **NOTE:** 1) Most, but not all, of the rocks in these tables are found in the NASIS (and PDP) choice lists (some are uncommon in the pedosphere). These additional rock types are included in these charts for completeness and to aid in the use of geologic literature. 2) Most, but not all of the rocks presented in these tables can be definitively identified in the field; some may require additional laboratory analyses; e.g., grain counts, thin section analyses, etc.

5/13/98

5-4

USDA - NRCS

IGNEOUS ROCKS CHART

CRYSTALLINE TEXTURE	KEY MINERAL COMPOSITION							
	Acidic (≈ Felsic) Potassium (K) Feldspar > 2/3 of Total Feldspar Content		Intermediate (---) Potassium (K) Feldspar & Plagioclase (Na, Ca) Feldspar in about equal proportions		Basic (≈ mafic) Plagioclase (Na, Ca) Feldspar > 2/3 of Total Feldspar Content			Ultrabasic (≈ ultramafic) Pyroxene and Olivine
	Quartz	No Quartz	Quartz	No Quartz	Sodic (Na) Plagioclase		Calcic (Ca) Plagioclase	peridotite (mostly olivine)
PEGMATITIC (very coarse, uneven-sized crystal grains)	granite pegmatite	syenite pegmatite	monzonite-pegmatite		Quartz	No Quartz	gabbro pegmatite	
PHANERITIC (crystals visible and of nearly equal size)	granite	syenite	quartz monzonite	monzonite	quartz-diorite granodiorite	diorite	gabbro	
PORPHYRITIC (relatively few visible crystals within a fine-grained matrix)	granite porphyry	syenite porphyry	quartz-monzonite porphyry	monzonite porphyry	quartz-diorite porphyry	diorite porphyry	diabase	} lava ³
	rhyolite porphyry	trachyte porphyry	quartz-latite porphyry	latite porphyry	dacite porphyry	andesite porphyry	porphyry basalt	
APHANITIC (crystals visible only with magnification) micro. ¹ crypto. ²	rhyolite	trachyte	quartz latite	latite	dacite	andesite	basalt	
GLASSY (amorphous: no crystalline structure)	obsidian (and its varieties: perlite, pitchstone, pumice, scoria) pyroclastics are shown on the Sedimentary and Volcaniclastic Rocks chart.				¹ Microcrystalline - crystals visible with ordinary magnification (hand lens, simple microscope). ² Cryptocrystalline - crystals only visible with SEM ³ Lava - generic name for extrusive flows of non-clastic, aphanitic rocks (rhyolite, andesite, basalt)			

METAMORPHIC ROCKS CHART

NONFOLIATED STRUCTURE			CRUDE ALIGNMENT	FOLIATED STRUCTURE (e.g. banded, etc.)				
CONTACT METAMORPHISM			MECHANICAL METAMORPHISM	REGIONAL METAMORPHISM			PLUTONIC METAMORPHISM	
<i>Low Grade</i>	<i>Medium Grade</i>	<i>High Grade</i>	<i>Very Low Grade</i>	<i>Low Grade</i>	<i>Medium Grade</i>	<i>High Grade</i>	<i>Extreme Grade</i>	
granofels	hornfels	marble	crush breccia mylonite	slate	phyllite greenstone	schist amphibolite	gneiss granulite	migmatite
metaquartzite	serpentinite	soapstone (<i>talc</i>)	← metaconglomerate → ← metavolcanics →					

* Not all rock types listed here can be definitively identified in the field (e.g. may require grain counts)

** Not all rock types shown here are available on Bedrock-Kind choice list. They are included here for completeness and as aids to using geologic literature.

SEDIMENTARY AND VOLCANICLASTIC ROCKS

CLASTIC				NONCLASTIC			
Dominant Grain Size				Chemical		Biochemical	Organic
Very Fine (Argillaceous) <0.002 mm	Fine (Argillaceous) 0.002-0.05 mm	Medium (Arenaceous) 0.05 - 2.0 mm	Coarse (Rudaceous) > 2.0 mm	Evaporates	Precipitates	Accretionates	Reduzates
	argillite shale	Sandstones (ss): arenite arkose (mainly feldspar) glaucconitic ss ("greensand") graywacke (dark, "dirty" ss) orthoquartzite (mainly quartz)	breccia (non-volcanic, angular frags) conglomerate (non-volcanic, rounded frags)	anhydrite (CaSO ₄) gypsum (CaSO ₄ • 2H ₂ O) halite (NaCl)	CARBONATE ROCKS Limestones (ls) (>50% calcite) chemical types: accretionary types		black shale (organics and fine sediments) bituminous ls bog iron ores coal
	mudstone claystone				caliche travertine tufa dolostone (>50% calcite + dolomite) phosphatic limestone	bio-stromal ls organic reef pelagic ls (chalk) bio-clastic types coquina oolitic ls lithographic ls	
VOLCANICLASTICS (includes Pyroclastics)				OTHER NONCLASTIC ROCKS			
	ignimbrite		agglomerate (rounded frags)	siliceous rocks (SiO ₂ dominated): chert (jasper, chalcedony, opal) diatomite rock phosphate iron bearing rocks (Fe-SiO ₂ dominated)			
	tuff		volcanic breccia (angular frags)				
	pumice (specific gravity <1.0; highly vesicular)						
	scoria (specific gravity >2.0; slightly to moderately vesicular)						

NORTH AMERICAN GEOLOGIC TIME SCALE ¹

Geologic Period	Geologic Epoch	Sub-Division	Oxygen Isotope Stage	Years (BP)
QUATERNARY	Holocene		(1)	0 to 10-12 ka*
	<i>Late Pleistocene</i>	Late Wisconsin	(2)	10-12 to 28 ka
		Mid Wisconsin	(3, 4)	28 to 71 ka
		Early Wisconsin	(5a - 5d)	71 to 115 ka
		<i>Late Sangamon</i> Sangamon	(5e)	115 to 128 ka
	Pleistocene	Late - Mid Pleistocene <i>(Illinoian)</i>	(6 - 8)	128 to 300 ka
		<i>Middle Pleistocene</i>	(9 - 15)	300 to 620 ka
		Early - Mid Pleistocene	(16 - 19)	620 to 770 ka
<i>Early Pleistocene</i>			770 ka to 1.64 Ma**	
TERTIARY	Pliocene			1.64 to 5.2 Ma
	Miocene			5.2 to 23.3 Ma
	Oligocene			23.3 to 35.4 Ma
	Eocene			35.4 to 56.5 Ma
	Paleocene			56.5 to 65.0 Ma
CRETACEOUS	<i>Late Cretaceous</i>			65.0 to 97.0 Ma
	<i>Early Cretaceous</i>			97.0 to 145.6 Ma
JURASSIC				145.6 to 208.8 Ma
TRIASSIC				208.8 to ≈ 243.0 Ma
PERMIAN				≈ 243.0 to 290.0 Ma
PENNSYLVANIAN				290.0 Ma to 322.8 Ma
MISSISSIPPIAN				322.8 to 362.5 Ma
DEVONIAN				362.5 to 408.5 Ma
SILURIAN				408.5 to 439.0 Ma
ORDOVICIAN				439.0 to 510.0 Ma
CAMBRIAN				510.0 to ≈ 570.0 Ma
PRECAMBRIAN				> ≈ 570.0 Ma

* ka = x 1,000

** Ma = x 1,000,000 (≈ = approximately)

¹ Modified from Morrison, 1991; Sibrava, et al., 1986; Harland, et al., 1990.

TILL TERMS

Genetic classification and relationships of till terms commonly used in soil survey. (Schoeneberger, 1994; adapted from Goldthwaite and Matsch, 1988.)

Location (Facies of tills grouped by position at deposition)	Till Types	
	Terrestrial	Waterlaid
Proglacial Till (at the front of, or in front of)	proglacial flow till	waterlaid flow till
Supraglacial Till (on top of, or within upper part of)	supraglacial flow till ^{1, 3} supraglacial melt-out till ¹ (ablation till - NP) ¹ (lowered till - NP) ² (sublimation till - NP) ²	-----
Subglacial Till (within the lower part of, or beneath)	lodgement till ¹ subglacial melt-out till subglacial flow till (= "squeeze till" ^{2, 3}) (basal till - NP) ¹ (deformation till - NP) ² (gravity flow till - NP) ²	waterlaid melt-out till waterlaid flow till iceberg till (= "ice-rafted")

¹ *Ablation till* and *basal till*, generic terms that only describe "relative position" of deposition, have been widely replaced by more specific terms that convey both relative position and process. *Ablation till* (any comparatively permeable debris deposited within or above stagnant ice) is replaced by *supraglacial melt-out till*, *supraglacial flow till*, etc. *Basal till* (any dense, non-sorted subglacial till) is replaced by *lodgement till*, *subglacial melt-out till*, *subglacial flow till*, etc.

² Additional (proposed) till terms that have not gained wide acceptance, and considered to be *Not Preferred*, should not be used.

³ Also called *gravity flow till* (not preferred).

VOLCANICLASTIC TERMS

Volcaniclastic Deposits (Unconsolidated)			
Size Scale: 0.062 mm ¹ 2 mm 64 mm ¹			
←----- tephra -----> (all ejecta)			
←----- ash ----->		←----- cinders -----> (specific gravity > 1.0 & < 2.0)	←----- bombs -----> (fluid-shaped coarse fragments)
←----- fine ash ----->	←----- coarse ash ----->	←----- lapilli -----> (specific gravity > 2.0)	←----- blocks -----> (angular-shaped coarse fragments)
←----- scoria ² -----> (slightly to moderately vesicular; specific gravity > 2.0)			
←----- pumiceous ash ³ ----->		←----- pumice -----> (highly vesicular; specific gravity < 1.0)	
Associated Lithified (Consolidated) Rock Types			
←----- fine tuff ----->	←----- coarse tuff ----->	←----- lapillistone -----> (sp. gr. > 2.0)	
←----- ignimbrite -----> (consolidated ash flows and nuee ardentes)		←----- agglomerate -----> (rounded, volcanic coarse fragments)	
←----- volcanic breccia -----> (angular, volcanic coarse fragments)			

¹ These size breaks are taken from geologic literature (Fisher, 1989) and based on the modified Wentworth scale. The 0.062 mm break is very close to the USDA's 0.05 mm break between *coarse silt* and *very fine sand* (Soil Survey Staff, 1993). The 64 mm break is close to the USDA's 76 mm break between *coarse gravel* and *cobbles*. (See "Relationships Among Particle Size Classes and Different Systems" in the "Profile / Pedon Description Section", under "Soil Texture".)

² A lower size limit of 2 mm is required in Soil Taxonomy, but is not required in geologic usage (Fisher, 1989).

³ The descriptor for pumice particles < 2 mm, as used in Soil Science. Geologic usage does not recognize any size restrictions for pumice.

REFERENCES

- Bates, R.L., and Jackson, J.A. (eds.). 1987. Glossary of geology, 3rd Ed. American Geological Institute, Alexandria, VA. 788 p.
- Fisher, R.V. 1989. Pyroclastic sediments and rocks. AGI Data Sheet 25.2.
In: Dutro, J.T., Dietrich, R.V., and Foose, R.M. 1989. AGI data sheets for geology in the field, laboratory, and office, 3rd edition. American Geological Institute, Washington, D.C.
- Goldthwaite, R.P. and Matsch, C.L. (eds.). 1988. Genetic classification of glacial deposits: final report of the commission on genesis and lithology of glacial quaternary deposits of the International Union for Quaternary Research (INQUA). A.A. Balkema, Rotterdam. 294 p.
- Harland, W.B., R.L. Armstrong, L.E. Craig, A.G. Smith, and D.G. Smith. 1990. A geologic time scale. Press Syndicate of University of Cambridge, Cambridge, UK. 1 sheet.
- Morrison, R.B. (ed.). 1991. Quaternary nonglacial geology: conterminous United States. Geological Society of America, Decade of North American Geology, Geology of North America, Vol. K-2. 672 p.
- Sibrava, V., D.Q. Bowen, and D.Q. Richmond (eds.). 1986. Quaternary glaciations in the Northern Hemisphere: final report of the International Geological Correlation Programme, Project 24. Quaternary Science Reviews, Vol. 5, Pergamon Press, Oxford. 514 p.
- Soil Survey Staff. 1995. Soil survey laboratory information manual. USDA - Natural Resources Conservation Service, Soil Survey Investigations Report No. 45, Version 1.0, National Soil Survey Center, Lincoln, NE. 305 pp.
- Soil Survey Staff. 1996 (being revised). Glossary of landforms and geologic materials. Part 629, National Soil Survey Handbook. USDA - Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.
- Tennissen, A.C. 1974. Nature of earth materials. Prentice-Hall, Inc., NJ.

LOCATION

Compiled by: P.J. Schoeneberger, W. Lynn, D.A. Wysocki, and E.C. Benham,
NRCS, Lincoln, NE.


PUBLIC LAND SURVEY

The Public Land Survey is the most widely used scheme in the U.S. for land surveying (legal location). Historically, many soil descriptions have been located using this system. Some states are not part of the Public Land Survey System and use the State Plane Coordinate System. The states include Connecticut, Delaware, Georgia, Kentucky, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, North Carolina, Ohio (parts), Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, and West Virginia.

The Public Land Survey System consists of a standard grid composed of regularly spaced squares which are uniquely numbered in reference to north-south Principle Meridians and to various, local, east-west Baselines. These squares are shown on U.S. Geological Survey topographic maps.

TOWNSHIPS AND RANGES - The primary grid network consists of squares (6 miles on a side) called Townships. Each Township can be uniquely identified using two coordinates: 1) **Township** (the north-south coordinate relative to a local, east-west Baseline); and 2) **Range** (the east-west coordinate relative to a local north-south Principle Meridian). (The local Baselines and Principle Meridians for the coterminous U.S. are shown on pp. 82-83, Thompson, 1987.) Commonly in soil survey, the local Baseline and the Principle Meridian are not recorded. The name of the appropriate USGS topographic 7.5-minute or 15-minute quadrangle is recorded instead; e.g., *Pleasant Dale, NE, 7.5 min. Quad.*


The **Township numbers** run in rows that parallel the local Baseline. Each Township row is sequentially numbered relative to the row's distance from, and whether it's north (N) or south (S) of the local Baseline; e.g., *T2N* (for the second township row north of the local Baseline). The **Range numbers** run in rows that parallel the local Principle Meridian. Range rows are sequentially numbered relative to the row's distance from, and whether it's east (E) or west (W) of the Principle Meridian e.g., *R2E* (for the second Range row east of the Principle Meridian in the area). The combined coordinates identify a unique square in the area; e.g., *T1S, R2E* (for Township 1 South and Range 2 East).



Modified from Mozola, 1989.

SECTIONS - Each Township square is further subdivided into smaller squares called **Sections**, which make up the secondary grid in this location system. Sections are 1 mile on a side (for a total of 36 Sections within each Township). The Section numbers begin in the northeast corner of a Township and progress sequentially in east-west rows, wrapping back and forth to fill in the Township; e.g., *Section 34, T1S, R2E* (for Section 34 of Township 1 South, Range 2 East).

CAUTION: Due to the curvature of the earth (trying to fit a flat grid to a non-flat surface), inaccessible areas (e.g., large swamps), or to joins to other survey schemes (e.g., pre-existing Metes and Bounds), you will occasionally find irregularities in the grid system. Adjustments to the grid layout result in non-standard sized, partial sections and/or breaks in the usual numbering sequence of sections. In some areas, **Lots** are appended to the northernmost tier of Sections in a Township to enable the adjoining Township to begin along the Baseline.



Modified from Mozola, 1989.

SUB-DIVISIONS - The tertiary (lower) levels of this system consist of subdividing Sections into smaller pieces that are halves or quarters of the Section. The fraction (1/2, 1/4) that the area of land represents of the Section is combined with the compass quadrant that the area occupies within the Section; e.g., *SW 1/4, Section 34, T1S, R2E* (for the southwest quarter of Section 34, Township 1 South, Range 2 East). Additional subdivisions, by quarters or halves, can be continued to describe progressively smaller areas. The information is presented consecutively, beginning with the smallest subdivision; e.g., *N 1/2, NW 1/4, SW 1/4, NW 1/4, of Section 34, T1S, R2E* (for the north half of the northwest quarter of the southwest quarter of the northwest quarter of Section 34, Township 1 South, Range 2 East).

NOTE: Point locations (e.g., soil pits) are measured, traditionally in English units, with reference to a specified section corner or quarter corner (1/4 post); e.g., *660 feet east and 1320 feet north of southwest corner post, Section 34, T1S, R2E.*

STATE PLANE COORDINATE SYSTEM

The State Plane Coordinate System is the second most widely used scheme in the U.S. for land surveying (legal location). Historically, many soil descriptions have been located using this system. The states that use this system are: Connecticut, Delaware, Georgia, Kentucky, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, North Carolina, Ohio (parts), Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, West Virginia. The other states use the Public Land Survey System.

The State Plane Coordinate System is based upon two principle lines in the state; a north-south line and an east-west line. Most USGS topographic quadrangle maps indicate the state grids by tick marks along the neatlines (outer-most border) on 7.5-minute topographic maps of states that use State Plane Coordinates.

Specific coordinates for a point are described by distance (commonly in meters) and primary compass direction [north (northing) / south (southing) and east (easting) / west (westing)] relative to the principle lines; e.g., *10,240 m easting, and 1,234 m northing.*

Contact the local State NRCS Office or the Regional MO Office for state-specific details.

UNIVERSAL TRANSVERSE MERCATOR (UTM) RECTANGULAR COORDINATE SYSTEM

The Universal Transverse Mercator (UTM) Rectangular Coordinate System is widely available and has been advocated as the universal map coordinate standard by the USGS (Morrison, 1987). It is a metric-based system whose primary unit of measure is the meter. The dominant UTM projection circles the globe and spans a wide range of latitudes [80° S through 84° N (the extreme polar areas require a different projection)]. The dominant projection is divided into 60 zones around the world. Zones begin at the International Date Line Meridian in the Pacific and progress eastward around the world. Each zone extends from pole to pole and spans 6 degrees of longitude. The logic of the UTM grid is similar to that of State Plane Coordinates. The UTM System uses 2 values to arrive at unique coordinates for any point on the earth's surface: 1) distance (and direction) away from the Equator called **Northing** (or **Southing**) to identify the hemisphere, and 2) distance away from the local zone's Meridian called an **Easting**.

Around the perimeter of 7.5-minute USGS topographic quadrangle maps are blue tic marks which intersect the map boundary at 1 km intervals. The

Northing measures the distance from the Equator northward (in the Southern Hemisphere the Southing measures the distance from the Equator southward); e.g., *4, 771,651 meters N*. The Easting measures the distance eastward from the local Meridian (the same Easting designation is used in the Southern Hemisphere); e.g., *305, 904 meters E*. A complete example: *305, 904 meters E; 4, 771,651 meters N; 16, N* (for the location of the capitol dome in Madison, Wisconsin, which is located within zone 16).

If the USGS topographic map has a complete kilometer grid (shown in blue), measure the distance (cm) from the point of interest to the closest north-south reference line (to the west of the point of interest). If the map scale is 1:24000, multiply the measured distance (cm) by 240 to calculate the actual ground distance in meters. If the scale is 1:20000, multiply by 200, etc. Add this partial distance to that of the chosen km reference line to obtain the Easting to be recorded.

If no kilometer grid is shown on the topographic map, locate the kilometer tic points along the east-west perimeters immediately south of the point of interest. Place a straight edge between the tic marks and draw a line segment south of the point of interest. Measure the distance (cm) from the point of interest to the east-west line segment. Multiply this distance by the appropriate map scale factor as mentioned above. Add this distance to that of the east-west baseline to obtain the Northing (distance from the Equator). The Northing must be identified as *N* for sites north of the Equator and *S* for sites south of the Equator.

Alternatively, a variety of clear UTM templates are commercially available which can be overlain upon the topographic map to facilitate determining distances and coordinates.

REFERENCES

Mozola, A.J. 1989. U.S. Public Land Survey. In: Dutro, J.T., Dietrich, R.V., and Foote, R.M. AGI Data Sheets, 3rd Ed., American Geologic Institute, United Book Press, Inc.


Thompson, M.M. 1987. Maps for America, 3rd Ed. U.S. Geological Survey, U.S. Dept. Interior, U.S. Gov. Print. Office, Washington, D.C.

MISCELLANEOUS

Compiled by: P.J. Schoeneberger, D.A. Wysocki, E.C. Benham, and H. LaGarry,
NRCS, Lincoln, NE.

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". **NOTE:** Within any given box, each quadrant contains the same total area covered, just different sized objects.



MEASUREMENT EQUIVALENTS & CONVERSIONS

METRIC TO ENGLISH

Known	Symbol	Multiplier	Product	Symbol
LENGTH				
micron (=10,000 Angstrom units)	μ	3.9370 $\times 10^{-5}$	inches	<i>in</i> or <i>"</i>
millimeters	<i>mm</i>	0.03937	inches	<i>in</i> or <i>"</i>
centimeters	<i>cm</i>	0.0328	feet	<i>ft</i> or <i>'</i>
centimeters	<i>cm</i>	0.03937	inches	<i>in</i> or <i>"</i>
meters	<i>m</i>	3.2808	feet	<i>ft</i> or <i>'</i>
meters	<i>m</i>	1.0936	yards	<i>yd</i>
kilometers	<i>km</i>	0.6214	miles (statute)	<i>mi</i>
AREA				
square centimeters	<i>cm</i> ²	0.1550	square inches	<i>in</i> ²
square meters	<i>m</i> ²	10.7639	square feet	<i>ft</i> ²
square meters	<i>m</i> ²	1.1960	square yards	<i>yd</i> ²
square kilometers	<i>km</i> ²	0.3861	square miles	<i>mi</i> ²
hectares	<i>ha</i>	2.471	acres	<i>ac</i>
VOLUME				
cubic centimeters	<i>cm</i> ³	0.06102	cubic inches	<i>in</i> ³
cubic meters	<i>m</i> ³	35.3146	cubic feet	<i>ft</i> ³
cubic meters	<i>m</i> ³	1.3079	cubic yards	<i>yd</i> ³
cubic meters	<i>m</i> ³	0.0008107	acre-feet (= 43,560 ft ³)	<i>acre-ft</i>
cubic kilometers	<i>km</i> ³	0.2399	cubic miles	<i>mi</i> ³
liters (=1000 cm ³)	<i>l</i>	1.0567	quarts (U.S.)	<i>qt</i>
liters	<i>l</i>	0.2642	gallons (U.S.)	<i>gal</i>
milliliter	<i>ml</i>	0.0338	ounces	<i>oz</i>
1 milliliter = 1 cm ³ = 1 gm (H ₂ O, at 25°C)				
MASS				
grams	<i>g</i>	0.03527	ounces (avdp.)	<i>oz</i>
kilograms	<i>kg</i>	2.2046	pounds (avdp.)	<i>lb</i>
megagrams (= metric tons)	<i>Mg</i>	1.1023	short tons (2000 lb)	
megagrams	<i>Mg</i>	0.9842	long tons (2240 lb)	

ENGLISH TO METRIC

Known	Symbol	Multiplier	Product	Symbol
LENGTH				
inches	<i>in</i> or <i>"</i>	2.54 x 10 ⁴	micron [= 10,000 Angstrom units (A)]	μ
inches	<i>in</i> or <i>"</i>	2.54	centimeters	<i>cm</i>
feet	<i>ft</i> or <i>'</i>	30.48	centimeters	<i>cm</i>
feet	<i>ft</i> or <i>'</i>	0.3048	meters	<i>m</i>
yards	<i>yd</i>	0.9144	meters	<i>m</i>
miles (statute)	<i>mi</i>	1.6093	kilometers	<i>km</i>
AREA				
square inches	<i>in</i> ²	6.4516	sq. centimeters	<i>cm</i> ²
square feet	<i>ft</i> ²	0.0929	sq. meters	<i>m</i> ²
square yards	<i>yd</i> ²	0.8361	sq. meters	<i>m</i> ²
square miles	<i>mi</i> ²	2.5900	sq. kilometers	<i>km</i> ²
acres	<i>ac</i>	0.405	hectares	<i>ha</i>
VOLUME				
acre-feet	<i>acre-ft</i>	1233.5019	cubic meters	<i>m</i> ³
acre-furrow-slice ≈ 2,000,000 lbs	<i>afs</i> (assumes b.d. = 1.3 g/cm ³)	= 6 in. thick layer that's 1 acre square		
cubic inches	<i>in</i> ³	16.3871	cubic centimeters	<i>cm</i> ³
cubic feet	<i>ft</i> ³	0.02832	cubic meters	<i>m</i> ³
cubic yards	<i>yd</i> ³	0.7646	cubic meters	<i>m</i> ³
cubic miles	<i>mi</i> ³	4.1684	cubic kilometers	<i>km</i> ³
gallons (U.S. liquid) (= 0.8327 Imperial gal)	<i>gal</i>	3.7854	liters	<i>l</i>
quarts (U.S. liquid)	<i>qt</i>	0.9463	liters (= 1000 cm ³)	<i>l</i>
ounces	<i>oz</i>	29.57	milliliters	<i>ml</i>
1 milliliter = 1 cm ³ = 1 gm (H ₂ O, at 25°C)				
MASS				
ounces (avdp.)	<i>oz</i>	28.3495	grams	<i>g</i>
ounces (avdp.) (1 troy oz. = 0.083 lb)				
pounds (avdp.)	<i>lb</i>	0.4536	kilograms	<i>kg</i>
short tons (2000 lb)		0.9072	megagrams (= metric tons)	<i>Mg</i>
long tons (2240 lb)		1.0160	megagrams	<i>Mg</i>

COMMON CONVERSION FACTORS

Known	Symbol	Multiplier	Product	Symbol
acres	<i>ac</i>	0.405	hectares	<i>ha</i>
acre-feet	<i>acre-ft</i>	1233.5019	cubic meters	<i>m³</i>
acre-furrow-slice ≈ 2,000,000 lbs	<i>afs</i> (assumes b.d. = 1.3 g/cm ³)	= 6 in. thick layer that's 1 acre square		
Angstrom units	<i>A</i>	1x 10 ⁻⁸	centimeters	<i>cm</i>
Angstrom units	<i>A</i>	1x 10 ⁻⁴	microns	<i>μ</i>
Atmospheres	<i>A</i>	1.0133 x 10 ⁶	dynes/cm ²	
Atmospheres	<i>atm</i>	760	mm of mercury (Hg)	
BTU (mean)	<i>BTU</i>	777.98	foot-pounds	
centimeters	<i>cm</i>	0.0328	feet	<i>ft</i> or ' <i></i>
centimeters	<i>cm</i>	0.03937	inches	<i>in</i> or " <i></i>
centimeters/second	<i>cm/s</i>	1.9685	feet/minute	<i>ft/min.</i>
centimeters/second	<i>cm/s</i>	0.0224	miles/hour	<i>mph</i>
chain (US)		66	feet	<i>ft</i>
chain (US)		4	rods	
centimeters	<i>cm³</i>	0.06102	cubic inches	<i>in³</i>
cubic centimeters	<i>cm³</i>	2.6417 x 10 ⁻⁴	gallons (U.S.)	<i>gal</i>
cubic centimeters	<i>cm³</i>	0.999972	milliliters	<i>ml</i>
cubic centimeters	<i>cm³</i>	0.0338	ounces (US)	<i>oz</i>
cubic feet	<i>ft³</i>	0.02832	cubic meters	<i>m³</i>
cubic feet (H ₂ O, 60°F)	<i>ft³</i>	62.37	pounds	<i>lbs</i>
cubic feet	<i>ft³</i>	0.03704	cubic yards	<i>yd³</i>
cubic inches	<i>in³</i>	16.3871	cubic centimeters	<i>cm³</i>
cubic kilometers	<i>km³</i>	0.2399	cubic miles	<i>mi³</i>
cubic meters	<i>m³</i>	35.3146	cubic feet	<i>ft³</i>
cubic meters	<i>m³</i>	1.3079	cubic yards	<i>yd³</i>
cubic meters	<i>m³</i>	0.0008107	acre-feet (= 43,560 ft ³)	<i>acre-ft</i>
cubic miles	<i>mi³</i>	4.1684	cubic kilometers	<i>km³</i>
cubic yards	<i>yd³</i>	0.7646	cubic meters	<i>m³</i>
degrees (angle)	°	0.0028	circumferences	
Faradays		96500	coulombs (abs)	
fathoms		6	feet	<i>ft</i>
feet	<i>ft</i> or ' <i></i>	30.4801	centimeters	<i>cm</i>
feet	<i>ft</i> or ' <i></i>	0.3048	meters	<i>m</i>
feet	<i>ft</i> or ' <i></i>	0.0152	chains (US)	
feet	<i>ft</i> or ' <i></i>	0.0606	rods (US)	
foot pounds		0.0012854	BTU (mean)	<i>BTU</i>
gallons (US)	<i>gal</i>	3.7854	liters	<i>l</i>
gallons (US)	<i>gal</i>	0.8327	Imperial gallons	

gallons (US)	<i>gal</i>	0.1337	cubic feet	<i>ft³</i>
gallons (US)	<i>gal</i>	128	ounces (US)	<i>oz</i>
grams	<i>g</i>	0.03527	ounces (avdp.)	<i>oz</i>
hectares	<i>ha</i>	2.471	acres	<i>ac</i>
horsepower		2545.08	BTU (mean)/hour	
inches	<i>in</i> or <i>"</i>	2.54×10^4	micron	μ
			[= 10,000 Angstrom units (A)]	
inches	<i>in</i> or <i>"</i>	2.5400	centimeters	<i>cm</i>
kilograms	<i>kg</i>	2.2046	pounds (avdp.)	<i>lb</i>
kilometers	<i>km</i>	0.6214	miles (statute)	<i>mi</i>
joules	<i>J</i>	1×10^7	ergs	
liters	<i>l</i>	0.2642	gallons (US)	<i>gal</i>
liters	<i>l</i>	33.8143	ounces	<i>oz</i>
liters (= 1000 cm ³)	<i>l</i>	1.0567	quarts (US)	<i>qt</i>
long tons (2240 lb)		1.0160	megagrams	<i>Mg</i>
megagrams	<i>Mg</i>	1.1023	short tons	
(= metric tons)			(2000 lb)	
megagrams	<i>Mg</i>	0.9842	long tons	
			(2240 lb)	
meters	<i>m</i>	3.2808	feet	<i>ft</i> or <i>'</i>
meters	<i>m</i>	39.37	inches	<i>in</i>
micron	μ	1×10^{-4}	centimeters	<i>cm</i>
micron	μ	3.9370	inches	<i>in</i> or <i>"</i>
(=10,000 Angstrom units)		$\times 10^{-5}$		
miles (statute)	<i>mi</i>	1.6093	kilometers	<i>km</i>
miles/hour	<i>mph</i>	44.7041	cent./second	<i>cm/s</i>
miles/hour	<i>mph</i>	1.4667	feet/second	<i>ft/s</i>
milliliter	<i>ml</i>	0.0338	ounces	<i>oz</i>
1 milliliter \approx 1 cm ³ = 1 gm (H ₂ O, at 25°C)				
milliliter	<i>ml</i>	1.000028	cubic centimeters	<i>cm³</i>
millimeters	<i>mm</i>	0.03937	inches	<i>in</i> or <i>"</i>
ounces	<i>oz</i>	29.5729	milliliters	<i>ml</i>
1 milliliter \approx 1 cm ³ = 1 gm (H ₂ O, at 25°C)				
ounces (avdp.)	<i>oz</i>	28.3495	grams	<i>g</i>
ounces (avdp.)				
1 troy oz. = 0.083 lb				
pints (US)	<i>pt</i>	473.179	cubic centimeters	<i>cm³</i>
			or <i>cc</i>	
pints (US)	<i>pt</i>	0.4732	liters	<i>l</i>
pounds (avdp.)	<i>lb</i>	0.4536	kilograms	<i>kg</i>

quarts (US liquid)	<i>qt</i>	0.9463	liters (= 1000 cm ³)	<i>l</i>
rods (US)		0.25	chains (US)	<i>ft</i>
rods (US)		16.5	feet (US)	<i>ft</i>
short tons (2000 lb)		0.9072	megagrams (= metric tons)	<i>Mg</i>
square centimeters	<i>cm²</i>	0.1550	square inches	<i>in²</i>
square feet	<i>ft²</i>	0.0929	square meters	<i>m²</i>
square inches	<i>in²</i>	6.4516	sq. centimeters	<i>cm²</i>
square kilometers	<i>km²</i>	0.3861	square miles	<i>mi²</i>
square meters	<i>m²</i>	10.7639	square feet	<i>ft²</i>
square meters	<i>m²</i>	1.1960	square yards	<i>yd²</i>
square miles	<i>mi²</i>	2.5900	square kilometers	<i>km²</i>
square yards	<i>yd²</i>	0.8361	square meters	<i>m²</i>
yards	<i>yd</i>	0.9144	meters	<i>m</i>

Guide to Map Scales and Minimum-Size Delineations ¹

Order of Soil Survey	Map Scale	Inches Per Mile	Minimum Size Delineation ²	
			Acres	Hectares
Order 1	1:500	126.7	0.0025	0.001
	1:1,000	63.4	0.100	0.004
	1:2,000	31.7	0.040	0.016
	1:5,000	12.7	0.25	0.10
	1:7,920	8.0	0.62	0.25
Order 2	1:10,000	6.34	1.00	0.41
	1:15,840	4.00	2.50	1.0
	1:20,000	3.17	4.00	1.6
	1:24,000 ³	2.64	5.7	2.3
Order 3	1:30,000	2.11	9.0	3.6
	1:31,680	2.00	10.0	4.1
Order 4	1:60,000	1.05	36	14.5
	1:62,500	1.01	39	15.8
	1:63,360	1.00	40	16.2
Order 5	1:80,000	0.79	64	25.8
	1:100,000	0.63	100	40
	1:125,000	0.51	156	63
	1:250,000	0.25	623	252
	1:500,000	0.127	2,500	1,000
Very General	1:750,000	0.084	5,600	2,270
	1:1,000,000	0.063	10,000	4,000
General	1:7,500,000	0.0084	560,000	227,000
	1:15,000,000	0.0042	2,240,000	907,000

¹ Modified from: Peterson, F.F. 1981. Landforms of the Basin and Range Province. Defined for soil survey. Nevada Agricultural Experiment Station. Technical Bulletin No. 28. Reno, NV. 52 p.

² Traditionally, the minimum size delineation is assumed to be a 1/4 inch square, or a circle with an area of 1/16 inches². Cartographically, this is about the smallest area in which a conventional soil map symbol can be legibly printed. Smaller areas can, but rarely are, delineated and the symbol "lined in" from outside the delineation.

³ Corresponds to USGS 7.5-minute topographic quadrangle maps.

COMMON SOIL MAP SYMBOLS (TRADITIONAL)

(From the National Soil Survey Handbook, Title 170, Part 601, 1990.) The following symbols are common on field sheets (original aerial photograph based soil maps) and in many soil surveys published prior to 1997. Current guidelines for map compilation symbols are in NSSH, Exhibit 627-5, 1997.

<u>FEATURE</u>	<u>SYMBOL</u>
LANDFORM FEATURES:	
SOIL DELINEATIONS:	
ESCARPMENTS:	
Bedrock	
Other than bedrock	
SHORT STEEP SLOPE	
GULLY	
DEPRESSION, <i>closed</i>	
SINKHOLE	
Prominent Hill or Peak	
EXCAVATIONS:	
Soil Sample Site	
Borrow pit	
Gravel pit	
Mine or quarry	
LANDFILL	

FEATURE	SYMBOL
MISC. SURFACE FEATURES:	
Blowout	☺
Clay spot	⊗
Gravelly spot	⋮
Lava flow	^
Marsh or swamp	☙
Rock outcrop (<i>includes sandstone and shale</i>)	∨
Saline spot	+
Sandy spot	⋮
Severely eroded spot	≡
Slide or slip (<i>tips point upslope</i>)	})
Sodic spot	∅
Spoil area	≡
Stony spot	○
Very stony spot	⊙
Wet spot	∩

FEATURE

SYMBOLS

ROAD EMBLEMS :

Interstate



Federal



State



County, farm or ranch



RAILROAD



POWER TRANSMISSION LINE
(normally not shown)



PIPELINE
(normally not shown)



FENCE
(normally not shown)




FEATURE

SYMBOLS


CULTURAL FEATURES (cont'd)

LEVEES:


Without road



With road



With railroad




Single side slope
(showing actual feature location)



DAMS

Medium or Small



Large



HYDROGRAPHIC FEATURES:

STREAMS:

Perennial, *double line*



Perennial, *single line*



Intermittent



Drainage end



**SMALL LAKES, PONDS
AND RESERVOIRS:**

Perennial water



Miscellaneous water



Flood pool line



Lake or pond
(perennial)



**MISCELLANEOUS
WATER FEATURES:**

Spring



Well, artesian



Well, irrigation



MISCELLANEOUS
CULTURAL FEATURES:

Farmstead, house (omit in urban areas)	■
Church	⊕
School	▣
Other Religion (label)	▲ Mt Carmel
Located object (label)	○ Ranger Station
Tank (label)	● Petroleum
Lookout Tower	⚓
Oil and/or Natural Gas Wells	⚓
Windmill	⚓
Lighthouse	⚓

FIELD SAMPLING

Compiled by: P.J. Schoeneberger and D.A. Wysocki, NRCS, Lincoln, NE.

INTRODUCTION

This section contains a variety of miscellaneous information pertinent to the sampling of soils in the field.

Additional details of soil sampling for the National Soil Survey Laboratory (NRCS, Lincoln, NE) are provided in Soil Survey Investigations Report No. 42 (Soil Survey Staff, 1996).

SOIL SAMPLING

The objective of the task determines the methodology and the location of the soil material collected for analysis. Sampling for Taxonomic Classification purposes involves different strategies than sampling for soil fertility, stratigraphy, hydric conditions, etc. There are several general types of samples and sampling strategies that are commonplace in soil survey.

SOIL SAMPLE KINDS -

Reference Samples (also loosely referred to as "grab" samples) - This is applied to any samples that are collected for very specific, limited analyses; e.g., only pH. Commonly, reference samples are not collected for all soil layers in a profile; e.g., only the top 10 cm; only the most root restrictive layer, etc.

Characterization Samples - These samples include sufficient physical and chemical soil analyses, from virtually all layers, to fully characterize a soil profile for Soil Taxonomic and general interpretive purposes. The specific analyses required vary with the type of material; e.g., a Mollisol requires some different analyses than does an Andisol. Nonetheless, a wide compliment of data (i.e., pH, particle size analysis, Cation Exchange Capacity, ECEC, Base Saturation, Organic Carbon content, etc.) are determined for all major soil layers.


SAMPLING STRATEGIES - [To be developed.]

FIELD EQUIPMENT CHECKLIST -

<p>Digging Tools (commonly choose 1 or 2; see graphic) Bucket Auger Sharp Shooter Montana Sharp Shooter (for rocky soils) Tile Spade (for well-cultivated or loose material) Spade (standard shovel) Push Probe (e.g., Backsaver®, Oakfield®, etc.) - include a clean-out tool Pulaski</p>
<p>Soil Description Knife Hand Lens (10X or combination lenses) Acid Bottle (1N - HCl) Water Bottle Color Book (e.g., Munsell®, EarthColors®, etc.) Picture Tapes ("pit tape" - metric preferred) Tape Measure (metric or English and metric) (3) Ultra-Fine Point Permanent Marker Pens Pocket pH Kit or Electronic "Wand" Pocket Soil Thermometer Camera Sample bags (for grab samples) Soil Description Sheet (232 or PEDON description form)</p>
<p>Site Description Field Note Book GPS Unit Abney Level Clinometer Compass Altimeter (pocket-sized)</p>
<p>Field References Field Book for Describing and Sampling Soils Aerial Photographs Topographic Maps (1:24,000, 7.5 min; 1:100,000) Geology Maps Soil Surveys (county or area) AGI Field Sheets</p>
<p>Personal Protective Gear Small First Aid Kit Leather Gloves Sunglasses Insect Repellent Sunscreen Hat Drinking Water</p>


EXAMPLES OF COMMON FIELD SAMPLING EQUIPMENT - (Use of trade or company names is for informational purposes only and does not constitute an endorsement.)

Digging Tools / Shovel Types




Soil Probes


Hydraulic Probes



Bucket Auger Types



External Thread Augers



REFERENCES

Soil Survey Staff. 1996. Soil survey laboratory methods manual. USDA - Natural Resources Conservation Service, Soil Survey Investigations Report No. 42, Version 3.0, National Soil Survey Center, Lincoln, NE. 693 pp.

INDEX

A

Across Slope • 3-23
Air Temperature • 1-1
Anthric Saturation • 1-12
Anthropogenic Feature • 1-4
Anthropogenic Features • 3-21
Area Covered (%) Example • 7-1
Argillans • 2-24
Aspect • See Slope Aspect

B

Bedrock • 1-18
 Hardness • 1-20
 Weathering Class • 1-20
 Depth To • 1-20
Bedrock-Fracture Interval Class • 1-19
Biological Concentrations • 2-17, 2-19
Bridging • 2-24
Brittleness • 2-49
Bypass Flow • 2-58

C

Calcium Carbonate Equivalent Test • 2-66
Carpedolith • 2-61
Cementing Agents • 2-48
Characterization Samples • 8-1
Chemical Response • 2-64
Clay Depletions • 2-13
Clay Films • 2-24
Climate • 1-1
Coats • 2-24
Color Location • 2-8
Color, Mechanical Condition • 2-8
Color, Redoximorphic Condition • 2-8
Common Conversion Factors • 7-4
Common Field Sampling Equipment (Examples) • 8-3
Common Horizons • 2-2
Common Soil Map Symbols (Traditional) • 7-8

Compositional Texture Modifiers • 2-33
Concentrations • 2-18
 Boundary • 2-23
 Color • 2-20
 Contrast • 2-20
 Hardness • 2-23
 Kind • 2-18
 Location • 2-22
 Moisture State • 2-20
 Quantity • 2-20
 Shape • 2-22
 Size • 2-20
Concentrations Discussion • 2-17
Concretions • 2-13, 2-17, 2-19
Consistence • 2-46
Contrast Of Soil Mottles • 2-12
Coprogeous Earth • 1-24, 2-33
County FIPS Code • 1-2, 1-3
Cracks • 2-58
 Depth • 2-60
 Kind • 2-59
 Relative Frequency • 2-60
Crust-Related Cracks • 2-59, 2-60
Cryptogamic Crust • 2-61
Crystals • 2-17, 2-19

D

Date • 1-1
Datum Name • 1-2
Decision Flowchart For Describing Soil Colors • 2-7
Delineation Size (Transect) • 1-3
Densic (Contact, Material) • 1-24
Depth (To Bedrock) • 1-20
Depth To Water Table • 1-13
Describer(s) Name • 1-1
Desert Pavement • 2-61
Diagnostic Horizons • 1-23, 2-4
Diagnostic Properties • 1-23
Diagnostic Properties - Mineral Soils • 1-24
Diagnostic Properties - Organic Soils • 1-24
Diagnostic Subsurface Horizons • 1-23
Diatomaceous Earth • 1-24, 2-33
Down Slope • 3-23
Drainage • 1-9

Drainage Class • 1-9
Durinode • 1-24, 2-19

E

Earth Cover - Kind • 1-14
Effervescence • 2-65
 Chemical Agent • 2-65
Electrical Conductivity • 2-66
Elevation • 1-4, 3-22
Endosaturation • 1-12
English To Metric • 7-3
Eolian Deposits • 1-16
Epipedons • 1-23
Episaturation • 1-12
Erosion • 1-20
 Degree Class • 1-21
 Kind • 1-20
Erosional Lag • 2-61
Evaporites • 1-19
Excavation Difficulty • 2-52
Excavation Difficulty Class • 2-52
Extra-Structural Cracks • 2-58

F

Faunal Burrows • 2-61
Fecal Pellets • 2-19
Ferriargillans • 2-15, 2-24
Field Equipment Checklist • 8-2
Field Notes • 2-67
Field Sampling • 8-1
Filaments • 2-22
Films • 2-15, 2-24
Finely Disseminated Materials • 2-17
FIPS Code • 1-3
Fissures • 2-58
Flat Plains (Geomorphic Components) • 1-8
Flooding • 1-10
 Duration • 1-11
 Frequency • 1-10
 Months • 1-11
Fluidity • 2-49
Fragment Roundness • *See Rock And Other Fragments - Roundness*

G

Geology • 5-1
References • 5-10
Geomorphic Component • 1-7, 3-24
Geomorphic Description • 1-4, 3-1, 3-11
References • 3-26
Geomorphic Description (Outline) • 3-10
Geomorphic Description System • 3-1
Geomorphic Information • 1-4
Gilgai Microfeatures • 2-39
Glacial Deposits • 1-16
Ground Surface • 2-4
Guide To Map Scales And Minimum-Size Delineations • 7-7

H

Hard Rock Fragments • 2-32
Hills (Geomorphic Components) • 1-7
Hillslope - Profile Position • 1-6, 3-23
Hillslope Position • See Hillslope - Profile Position
Horizon Boundary • 2-5
Distinctness • 2-5
Topography • 2-6
Horizon Depth • 2-4
Horizon Modifiers (Other) • 2-4
Numerical Prefixes • 2-4
Prime • 2-4
Horizon Nomenclature • 2-2, 4-1
Horizon Nomenclature Conversion Charts • 4-5
Horizon Subscripts • See Horizon Suffixes
Horizon Suffixes • 2-3, 4-3
Horizon Thickness • 2-5
Hortonian Flow • 1-21
Hydraulic Conductivity • 2-62
Hydrophobic Layer • 2-61
Hypocoats • 2-15, 2-23, 2-24

I

Ice Wedge Cast • 2-61
Igneous Rocks • 1-18
Index (Of) Surface Runoff Classes • 1-22
In-Place Deposits • 1-16
Interbedded Rocks • 1-19

Internal Flow • 1-21
Interstitial Pores • 2-56
Inter-Structural Voids • 2-56
Interval (Transect) • 1-4
Iron Depletions • 2-13
Irregular Pores • 2-56

K

Krotovinas • 2-61

L

Lamellae • 1-24, 2-22, 2-61
Lamina • 2-61
Landform • 1-4, 3-11
Landform Subset Lists • 3-10, 3-15
Landscape • 1-4, 3-11
Landuse • See Earth Cover - Kind
Limnic Materials • 1-24, 2-33
Lithic Contact • 1-24
Local Physiographic / Geographic Name • 3-8
Local Physiographic/Geographic Name • 1-4
Location • 1-2, 6-1
References • 6-5

M

Major Land Resource Area • 1-3
Mangans • 2-15, 2-24
Manner Of Failure • 2-49
Marl • 1-24, 2-33
Mass Movement Deposits • 1-16
Masses • 2-13, 2-17, 2-19
Master Horizons • 2-2, 4-1
Matrix Color • See Soil Matrix Color
Mean Sea Level • 1-4
Measurement Equivalents & Conversions • 7-2
Metamorphic Rocks • 1-18
Metric To English • 7-2
Microbiotic Crust • 2-61
Microfeature • 1-4
Microfeature Terms • 3-20
Microrelief • 1-8, 2-39, 3-26
Minimum Data Set • 2-67

Miscellaneous • 7-1
Miscellaneous Field Notes • 2-67
MLRA • 1-3
Month / Day / Year • 1-1
Mottles • 2-9
 Color • 2-11
 Contrast • 2-11
 Moisture State • 2-11
 Quantity • 2-9
 Shape • 2-11
 Size • 2-9
Mountains (Geomorphic Components) • 1-8
Multicolored Pattern • 2-8

N

Name • 1-1
Nodules • 2-13, 2-17, 2-19
North American Geologic Time Scale • 5-7

O

Observation Method • 2-1
 Kind • 2-1
 Relative Size • 2-1
Odor • 2-67
Organic Deposits • 1-17
Organics • 1-19
Overland Flow • 1-21
Oxidation / Reduction • 2-13

P

Para (Soft) Rock Fragments • 2-32
Paralithic Contact • 1-24
Parent Material • 1-16
Particle Size Distribution • 2-28
Ped & Void Surface Features • 2-24
 Amount • 2-26
 Color • 2-27
 Continuity • 2-26
 Distinctness • 2-27
 Kind • 2-24
 Location • 2-27
Pendant • 2-23

Penetration Resistance • 2-51
Penetration Resistance Class • 2-52
Perched Water Table • 1-13
Permeability • 2-63
Permeability (Discussion) • 2-62
Permeability Class • 2-63
Permeability Coefficient • 2-62
pH • 2-64
Physiographic Division • 1-4
Physiographic Divisions • 3-2
Physiographic Location • 1-4, 3-2
Physiographic Province • 1-4
Physiographic Provinces • 3-2
Physiographic Section • 1-4
Physiographic Sections • 3-2
Phytoliths • 2-19
Pipestems • 2-22
Plant Common Name • 1-15
Plant Opal • *See* Phytoliths
Plant Scientific Name • 1-15
Plant Symbol • 1-15
Plasticity • 2-50
Plasticity Class • 2-50
Plinthite • 1-24, 2-22
Pocket Penetrometer • 2-51
Ponding • 1-11
 Depth • 1-11
 Duration • 1-11
 Frequency • 1-11
Pores • 2-56
 Quantity • 2-53
 Quantity Class • 2-53
 Shape • 2-56
 Size • 2-53
 Size Classes • 2-55
 Vertical Continuity • 2-58
Pores Discussion • 2-56
Precipitates • 1-19
Preferential Flow • 2-58
Pressure Faces • 2-24
Primary Packing Voids • 2-56
Profile / Pedon Description • 2-1
 References • 2-68
Public Land Survey • 6-1

Q

Quadrangle Name • 1-2

R

Ranges (Location) • 6-1

Reaction (pH) • 2-64

Redox Concentrations • 2-13, 2-14

Redox Depletions • 2-13, 2-14

Redoximorphic Features • 2-14

 Boundary • 2-16

 Color • 2-16

 Contrast • 2-16

 Hardness • 2-16

 Kind • 2-14

 Location • 2-16

 Moisture State • 2-16

 Quantity • 2-15

 Shape • 2-16

 Size • 2-16

Redoximorphic Features (RMF) Discussion • 2-13

Reduced Conditions • 2-66

Reduced Matrix • 2-14

Reference Samples • 8-1

RMF Shapes • 2-22

Rock And Other Fragments • 2-35

 Kind • 2-35

 Roundness • 2-35

 Size Classes And Descriptive Terms • 2-37

 Volume Percent • 2-35

Rock Charts • 5-3

Rock Fragments • 2-32

 Quantity • 2-32

 Size • 2-32

Root Pseudomorphs • 2-22

Roots • 2-53

 Location • 2-55

 Quantity • 2-53

 Quantity Class • 2-53

 Size • 2-53

 Size Classes • 2-55

Roundness • 2-35

Runoff • 1-21

Rupture Resistance • 2-46
Blocks, Peds, And Clods • 2-47
Surface Crust And Plates • 2-48

S

Salinity • 2-66
Salinity Class • 2-66
Sand Coats • 2-24
Satiation • 1-12
Saturated Hydraulic Conductivity • 2-62, 2-63
Saturated Hydraulic Conductivity (Discussion) • 2-62
Saturation • 1-12
Scale • 1-2
Scientific Plant Name • 1-15
Seasonal High Water Table - Kind • 1-13
Secondary Carbonates • 1-24
Sections • 6-2
Sedimentary Rocks • 1-19
Series Name • 1-4
Shell Fragments • 2-19
Shot • 2-22
Silt Coats • 2-24
Site Description • 1-1
Skeletans • 2-24
Slickensides • 1-24, 2-24
Slope Aspect • 1-5, 3-22
Slope Complexity • 1-5, 3-22
Slope Gradient • 1-5, 3-22
Slope Shape • 1-6, 3-22
Smeariness • 2-49
Sodium Adsorption Ratio (SAR) • 2-66
Soft Powdery Lime • *See* Secondary Carbonates
Soft Rock Fragments • 2-32
Soil Color • 2-7
Soil Matrix Color • 2-7
Soil Moisture Status • *See* Soil Water State
Soil Permeability • 2-62
Soil Sample Kinds • 8-1
Soil Series Name • 1-4
Soil Structure • 2-38
 Grade • 2-40
 Size • 2-40
 Type • 2-38
Soil Surface • 2-5

Soil Survey Area Identification Number (SSID) • 1-2
Soil Taxonomy • 4-1
 References • 4-9
Soil Temperature • 1-1
Soil Temperature Depth • 1-2
Soil Texture • 2-28
Soil Water State • 1-12
Special Features • 2-61
Sphericity • 2-36
State Abbreviation • 1-2
State Physiographic Area • 1-4, 3-8
State Plane Coordinate System • 6-4
Stickiness • 2-50
Stickiness Class • 2-50
Stone Line • 2-61
Stop Number (Transect) • 1-3
Stress Features • 2-24
Stringers • 2-22
Structure • See Soil Structure
Structure Shape • See Soil Structure - Type
Subordinate Distinctions • See Horizon Suffixes
Surface Coats • 2-15
Surface Crusts • 2-61
Surface Fragments • 1-22
 Kind • 1-22
 Mean Distance Between Fragments • 1-22
Surface Morphometry • 1-4, 3-22
Surface Runoff • 1-21
Surface Stoniness • 1-22

T

Table Comparing Particle Size Systems • 2-31
Taxonomic Classification • 2-2
Terms Used In Lieu Of Texture • 2-34
Terraces (Geomorphologic Components) • 1-7
Texture • See Soil Texture
Texture Class • 2-28
Texture Modifiers • 2-29, 2-32
Texture Modifiers (Compositional) • 2-33
Threads • 2-22
Throughflow • 1-21
Till Terms • 5-8
Topographic Map • 1-2
Topographic Quadrangle • 1-2

Townships (Location) • 6-1
Transect Delineation Size • 1-3
Transect Direction • 1-3
Transect ID • 1-3
Transect Kind • 1-3
Transect Selection Method • 1-3
Transect Stop Number • 1-3
Transects • 1-3
Trans-Horizon Cracks • 2-59
Transitional Horizons • 2-2, 4-1
Transported Deposits • 1-17
Tubular Pores • 2-56

U

Unconfined Compressive Strength • 2-51
Undulation • 2-6
Universal Transverse Mercator (UTM) Rectangular Coordinate System • 6-4

V

Variigated • 2-8
Vegetation / Land Cover • 1-14
Vesicular Pores • 2-56
Volcanic Deposits • 1-17
Volcaniclastic Terms • 5-9
Vughs • 2-56

W

Water Laid Deposits • 1-17
Water Status • 1-9
Water Table • See Depth To Water Table, Seasonal High Water Table
Wedge Structure • 2-39