

106501

May 2008



**ENTERED**

LA-UR-08-2783

*Approved for public release;  
distribution is unlimited.*

## *Site-Representative Biota Concentration Guides at Los Alamos*

Mike McNaughton, Phil Fresquez, Bill Eisele



Los Alamos National Laboratory, MS M992, Los Alamos, NM 87545

31953



## Introduction

Biota dose assessment at Los Alamos National Laboratory, LANL, follows the guidance of the DOE Standard, DOE-STD-1153-2002, as described in the LANL Biota Dose Assessment Quality Assurance Project Plan, RRES-MAQ-BIOTA [http://int.lanl.gov/environment/all/docs/qa/maq/MAQ-BIOTA\\_R0.pdf](http://int.lanl.gov/environment/all/docs/qa/maq/MAQ-BIOTA_R0.pdf) and LANL procedure ENV-MAQ-514 [http://int.lanl.gov/environment/all/docs/qa/maq/RRES-MAQ-514\\_R1.pdf](http://int.lanl.gov/environment/all/docs/qa/maq/RRES-MAQ-514_R1.pdf).

According to DOE-STD-1153-2002 Section 2 and RRES-MAQ-BIOTA Section 7.1.1, general screening is performed using Biota Concentration Guides, BCGs, which are defined as "the limiting concentration of a radionuclide in soil, water, or sediment that would not lead to exceeding dose limits for the protection of biota in the environment." For the general screening, the maximum concentrations in water, sediment, and soil are compared with each radionuclide-specific BCG (DOE-STD-1153-2002 Table 2.1) and the sum of the fractions is calculated (DOE-STD-1153-2002 page M1-25.)

The DOE Standard encourages each site to establish site-specific or site-representative BCGs (DOE-STD-1153-2002 Module 1 pages 10, 24-26, 43, 45-47, 50, 59, 61-64, 79, 81-86; Module 2 pages 31, 41, 49, 50, 53; and Module 3 page 29). The purpose of this paper is to establish site-representative BCGs for cesium-137 and strontium-90 in water, sediment, and soil.

Sections 4.1 and 7.1.3.2 of DOE-STD-1153-2002 recommend the use of site-specific "lumped parameter values." The important lumped parameter is the bioaccumulation factor referred to in the DOE Standard as  $B_{iv}$ . It is also commonly called the wet-weight concentration ratio  $C_r$ ; see Module 3 Section 3.1 (page M3-19) of DOE-STD-1153-2002.

### Cesium-137, terrestrial

Cesium-137 in the environment is the subject of NCRP Report No. 154 (NCRP 2008), which examines the behavior of cesium in the environment. At the Savannah River Site (SRS) cesium bioaccumulates ( $C_r > 1$ ) because the soil is deficient in potassium (NCRP Report No. 76, NCRP 2008 Section 4.1, and Eisenbud and Gesell 1997.) At Los Alamos, however, the cesium concentration ratio is less than one, because of the relative abundance of potassium. For a comparison of potassium concentrations within the US, see <http://energy.cr.usgs.gov/radon/usak.gif> and [http://tin.er.usgs.gov/geochem/doc/lower48maps/k\\_icp40.jpg](http://tin.er.usgs.gov/geochem/doc/lower48maps/k_icp40.jpg).

At Los Alamos, cesium-137 concentrations have been measured in soil, vegetation, and animals in and near LANL (Bennett 1996, Fresquez 1996, 1997, 1998a, 1998b, and 2004, Hakonson 1973 and 1975, Miera 1978, White 1981, and the annual site environmental reports titled "Environmental Surveillance at Los Alamos.") The ratios are variously reported on the basis of wet-weight, dry weight, or ash weight. For the present paper, they are converted to wet-weight ratios using the conversion factors in Fresquez 2007.

The results consistently show that the wet-weight concentrations of cesium-137 in plants and animals are all much less than the concentrations in soil. In other words, the bioaccumulation factors are much less than 1.

Salazar 2006 examined the LANL plant data and concluded that the concentration ratio of wet-weight vegetation to soil is consistent with the range:  $C_r = 0.01$  to  $0.06$ . Comparable numbers have been reported at other locations that are not deficient in potassium, for example refer to NUREG/CR-5512 (1992) (available at [http://www.pnl.gov/bayesian/refs/KennedyJr1992A\\_NUREG-CR-5512%20PNL-7994.pdf](http://www.pnl.gov/bayesian/refs/KennedyJr1992A_NUREG-CR-5512%20PNL-7994.pdf) ) and Baes 1984 (ORNL-5786, available at <http://homer.ornl.gov/baes/documents/ornl5786.html> .)

Appendix A summarizes the deer and elk data (Fresquez 1998b) and also concludes that  $C_r < 0.06$  for cesium-137. If  $C_r \ll 1$ , the internal dose is much less than the external dose, in which case the exact value of  $C_r$  is not important. For terrestrial biota dose assessment of cesium-137 at LANL, it is assumed that  $C_r = B_{iv} = 0.06$ .

For a given concentration, the dose may be calculated using the coefficients in Tables 2.3 and 2.4 (page M3-15 to M3-18) of DOE-STD-1153-2002, or using RESRAD-BIOTA with the following changes to the defaults.

- $B_{iv}$  for soil to terrestrial animal and plant =  $0.06$ .

The limiting LANL site-representative BCG for cesium-137 soil is 2,000 pCi/g because this concentration would result in less than the DOE limit of 0.1 rad/day to terrestrial animals.

### **Strontium-90, terrestrial**

Whereas uptake of cesium-137 depends on the concentration of potassium, similarly the uptake of strontium-90 depends on the concentration of calcium (Eisenbud and Gesell 1997, NCRP Report No. 76, 1984, NCRP Report No. 110.) For nationwide calcium concentrations, see <http://tin.er.usgs.gov/geochem/doc/averages/ca/usa.html> and [http://tin.er.usgs.gov/geochem/doc/lower48maps/ca\\_icp40.jpg](http://tin.er.usgs.gov/geochem/doc/lower48maps/ca_icp40.jpg) . Calcium is relatively abundant in New Mexico soil and water so bioaccumulation is not expected.

The concentration ratio for strontium-90 has been measured at and near Los Alamos (Fresquez 1997, Salazar 2006.) For edible crop tissues, the ratios of wet-weight vegetation to soil are consistently less than 1 (Salazar 2006.) However, Fresquez 1997 reported a ratio of 4 for the non-edible tissues of pinto beans. It is possible that this measurement may have been influenced by rain splash (White 1981, Hackonson 1993) or by the low calcium content of the soil used in this study. In contrast, recent measurements of common lambsquarters and pigweed amaranth in low-calcium soil (LANL 2007) indicated a dry-weight ratio of 2, which corresponds to a wet-weight ratio less than 1.

Furthermore, a comparison of the strontium-90 wet-weight concentrations in deer and elk (Fresquez 1998) with the concentrations in soil (Fresquez 1996) (Appendix A) shows the concentration ratio is 3.3 for bone and 0.01 for muscle. To be conservative, the worst-

case bioaccumulation factor for strontium-90 is assumed to be 4 for both plants and animals.

Therefore, the following changes were made to the RESRAD-BIOTA defaults.

- $B_{iv}$  for soil to terrestrial animals and plants = 4.

Using these values in RESRAD-BIOTA indicates the limiting site-representative BCG for strontium-90 in soil is 300 pCi/g because this concentration would result in less than the DOE limit of 0.1 rad/day to terrestrial animals.

### **Cesium-137, aquatic**

There are no measurements of the concentration factor in aquatic animals at LANL because there are almost no aquatic animals to measure. Therefore the LANL site-representative BCGs rely on calculations. These calculations focus on Effluent Canyon, which has the highest radionuclide concentrations because it receives the effluent from the permitted outfall of the radioactive-liquid-waste treatment facility at TA-50.

The concentration ratio,  $C_r$ , for fish could be estimated from the potassium concentration,  $K$ , (micro-mol/L) and the sediment load,  $SL$ , (mg/L) using the following equation for non-piscivore fish from NCRP Report No. 154, p 244.

$$\log C_r = 4.332 - 0.718 \log(K) - 0.233 \log(SL).$$

According to Table S6-4 of the 2006 Environmental Surveillance Report, for Effluent Canyon,  $K = 200 \pm 40$  micro-mol/L and  $SL = 3 \pm 1$  mg/L, so  $C_r = 400 \pm 200$ . See also Fig. 3.9 of NCRP Report No. 76, which indicates an upper bound of  $C_r = 600$  for a potassium concentration of 8 mg/kg.

However, there are no fish at LANL, so consider the equation for invertebrates (NCRP Report No. 154, p 243.)

$$\log C_r = 3.628 - 0.583 \log(K)$$

So for invertebrates,  $C_r = 190 \pm 30$ .

For simplicity, assume  $C_r = 200$ . If the  $^{137}\text{Cs}$  concentration in the water is 20,000 pCi/L, the concentration in aquatic biota is  $200 \times 20,000 = 4,000,000$  pCi/L = 4,000 pCi/g, which results in a dose of 0.17 rad/day. This is 17% of the limit of 1 rad/day for aquatic biota.

Using the default parameters in RESRAD-BIOTA, there would be an additional dose of 0.2 rad/day from cesium-137 in sediment. In this case, the total dose would be  $0.17 + 0.2 = 0.37$  rad/day, which is well below the limit of 1 rad/day for aquatic biota.

Consider the dose to a riparian animal. The contaminated section of Effluent Canyon is small and has almost no food. There is more food available in the uncontaminated wetlands, 10 m upstream of the radioactive-liquid-waste outfall, and in the nearby human areas replete with irrigated picnic areas. Therefore, an area factor is appropriate.

The site-representative riparian animal is the southwestern willow flycatcher, which has a minimum individual home range of 0.4 ha and a minimum population area of  $40 \times 0.4 =$

16 ha (Ryti 2004, Gonzales 2006.) The area of the highly contaminated riparian habitat of Effluent Canyon is 0.05 ha (LANL 2006) so the population area factor is  $0.05/16 = 0.003$ . If this area factor is entered into RESRAD-BIOTA it is apparent that the dose to a riparian animal is less than 2% of the DOE limit.

RESRAD-BIOTA may also be used to demonstrate that the dose to terrestrial animals from drinking water containing 20,000 pCi/L is 3% of the DOE limit.

The following changes were made to the RESRAD-BIOTA defaults.

- $B_{iv}$  for water to aquatic and riparian animals = 200;
- calculate the sediment concentration from the default  $K_d$ ;
- riparian animal area factor = 0.003

Using these values in RESRAD-BIOTA indicates the limiting site-representative BCG for cesium-137 in water is 20,000 pCi/L. This is conservative and would result in less than half of the biota dose limits for aquatic, terrestrial, and riparian biota.

### **Strontium-90, aquatic**

Fig. 3.10 of NCRP Report No. 76, 1984, shows the bioaccumulation factor for fish as a function of calcium concentration (based on Vanderploeg 1975, ORNL-5002). According to the annual site environmental report for 2006 (LA-14341-ENV) the calcium concentration in filtered water at Effluent Canyon is 21 mg/L, and for this concentration, the bioaccumulation factor is 450 for bone and 4 for flesh. For reference man (ICRP 23) the mass of bone is less than 20% of the total body mass, and the proportion in fish is less than in humans; for example, Toppe et al.(2007) indicate that 10 to 15% of total fish biomass are bones from the head and vertebrae. To be conservative, assume 20% bone for fish; then the average bioaccumulation factor is 100, with an uncertainty of a factor of 2.

According to RESRAD-BIOTA, if the concentration of  $^{90}\text{Sr}$  in water is 30,000 pCi/L and the bioaccumulation factor is 100, the total dose to aquatic biota would be 0.2 rad/day, which is 20% of the DOE limit.

Using the area factor discussed previously, the dose to the representative riparian animal is 3 mrad/day, which is 3% of the limit.

The limiting dose from strontium-90 in water is to terrestrial animals that drink the water. Using the RESRAD-BIOTA defaults parameters, the dose to terrestrial animals from a  $^{90}\text{Sr}$  concentration of 30,000 pCi/L would be 55% of the DOE limit.

In summary, the following changes were made to the RESRAD-BIOTA defaults for strontium-90.

- $B_{iv}$  for water to aquatic animals = 100;
- $B_{iv}$  for water to riparian animals = 400;
- calculate the sediment concentration from the default  $K_d$ ;
- riparian animal area factor = 0.003

Using these values in RESRAD-BIOTA indicates the limiting site-representative BCG for strontium-90 in water is 30,000 pCi/L. This is conservative for aquatic, terrestrial, and riparian biota.

### **Sediment**

RESRAD-BIOTA includes default distribution coefficients,  $K_d$ , which are "for use in calculating BCGs for sediment or water for an aquatic system evaluation in the absence of co-located water and sediment," (DOE-STD-1153-2002 Module 3, page M3-52, Table 4.3.) In the absence of LANL site-specific data, these should be used for aquatic evaluations. For terrestrial evaluations of dry sediment in arroyos or stream banks, the soil BCGs should be used.

### **Conclusion**

The LANL limiting site-representative biota concentration guides, BCGs, are summarized in the following table.

Radionuclide	soil (pCi/g wet)	water (pCi/L)
Cesium-137	2,000	20,000
Strontium-90	300	30,000

## References

Baes CF, Sharp RD, Sjoreen, AL, Shor RW. *A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides through Agriculture*. ORNL-5786 (1984).

(Available at <http://homer.ornl.gov/baes/documents/ornl5786.html> )

Bennett K, Biggs J, Fresquez P. *Radionuclide Contaminant Analysis of Small Mammals, Plants, and Sediments within Mortandad Canyon, 1994*. LA-13104-MS (1996).

DOE-STD-1153-2002. *A graded approach for evaluating radiation doses to aquatic and terrestrial biota*. 2002.

Eisenbud M, Gesell T. *Environmental Radioactivity*. Academic Press 1997.

Fresquez PR, et al. *Radionuclides and radioactivity in soils within and around Los Alamos National Laboratory, 1974 through 1994, trends and dose comparisons*. LA-13149-MS (1996).

Fresquez, PR, D.A. Armstrong, M.A. Mullen, and L. Naranjo, Jr. *Radionuclide concentrations in pinto beans, sweet corn, and zucchini squash grown in Los Alamos Canyon at LANL*. LA-13304-MS (1997)

Fresquez 1998a. Fresquez, PR, D.A. Armstrong, M.A. Mullen, and L. Naranjo, Jr. *The uptake of radionuclides by beans, squash, and corn growing in contaminated alluvial soils at Los Alamos National Laboratory*, Journal of Environmental Science and Health B33 (1) 99-115 (1998).

Fresquez 1998b. Fresquez PR, Biggs JR, Bennett KD, Kraig DH, Mullen MA, Ferenbaugh JK. *Radionuclide Concentrations in Deer and Elk from Los Alamos National Laboratory*. LA-13553-MS (December 1998)

Fresquez 2004: Fresquez, P.R. and G.J. Gonzales. *Radionuclide Concentrations in Vegetation at the Los Alamos National Laboratory in 2002/2003*, Los Alamos National Laboratory report LA-14160-PR. Los Alamos, NM (2004).

Fresquez 2007: Fresquez PR, Ferenbaugh JK, Naranjo L, Jr. *Moisture conversion ratios for the foodstuffs and nonfoodstuffs biota environmental surveillance programs at LANL*. LA-UR-07-0280 (2007).

Gonzales G, Rytí R, Gallegos P, Gallegos A. *The Application of ECORSK.9 to the Mexican Spotted Owl, Western Bluebird, Deer Mouse, and Southwestern Willow Flycatcher in Mortandad Canyon at the Los Alamos National Laboratory*. LA-14305 (2006).

Hakonson TE, Nyhan JW, Johnson LJ, Bostick KV. *Ecological Investigation of Radioactive Materials in Waste Discharge Areas at Los Alamos*. LA-5282-MS, (1973).

Hakonson TE, K.V. Bostick KV. *Cesium-137 and plutonium in liquid waste discharge areas at Los Alamos*. Radioecology and Energy Resources, Proceedings of the Fourth National Symposium on Radioecology, 1975, edited by Colbert E. Cushing, Jr., and published by Dowden, Hutchinson, and Ross. (1975).

Hakonson, T.E. and L.J. Lane, 1993. *The Role of Physical Process in the Transport of Man-made Radionuclides in Arid Ecosystems*, R. M. Harrison (ed.) in Biogeochemical Pathways of Artificial Radionuclides, John Wiley & Sons.

ICRP 23. Reference Man. 1974.

LANL Biota Dose Assessment Quality Assurance Project Plan, RRES-MAQ-BIOTA  
[http://int.lanl.gov/environment/all/docs/qa/maq/MAQ-BIOTA\\_R0.pdf](http://int.lanl.gov/environment/all/docs/qa/maq/MAQ-BIOTA_R0.pdf)

LANL procedure ENV-MAQ-514  
[http://int.lanl.gov/environment/all/docs/qa/maq/RRES-MAQ-514\\_R1.pdf](http://int.lanl.gov/environment/all/docs/qa/maq/RRES-MAQ-514_R1.pdf) .

LANL 2006. *Mortandad Canyon Investigation Report*. LA-UR-06-6752 (2006)

LANL 2007. *Environmental Surveillance at Los Alamos During 2006*. LA-14341-ENV (2007)

Miera FR and T.E. Hakonson TE. *Radiation dose to rodents inhabiting a radioactive waste receiving area*. Health Physics 34, pp 603-609 (1978).

NCRP Report No. 76. *Radioecological Assessment: Predicting the Transport, Bioaccumulation, and uptake by Man of Radionuclides Released to the Environment*. NCRP 1984.

NCRP Report No. 110.

NCRP Report No. 154. *Cesium-137 in the Environment: Radioecology and Approaches to Assessment and Management*. NCRP 2008.

NUREG/CR-5512 (1992) (available at  
[http://www.pnl.gov/bayesian/refs/KennedyJr1992A\\_NUREG-CR-5512%20PNL-7994.pdf](http://www.pnl.gov/bayesian/refs/KennedyJr1992A_NUREG-CR-5512%20PNL-7994.pdf) )

Ryti RT, Markweise J, Miranda R, Soholt L. *Preliminary Remediation Goals for Terrestrial Wildlife*. Human and Ecological Risk Assessment, 10: 437-450 (2004)

Salazar S, McNaughton M, Fresquez PR. *Concentration Ratios for cesium and Strontium in Produce Near Los Alamos*. LA-14284 (2006).

Toppe J, Albrektsen S, Hope B, Aksnes A, *Chemical composition, mineral content and amino acid and lipid profiles in bones from various fish species*, Comparative Biochemistry and Physiology, Part B 146 (2007) 395-401.

Vanderploeg HA et al. *Bioaccumulation Factors for Radionuclides in Freshwater Biota*. ORNL-5002. 1975.

Whicker FW, Schultz V. *Radioecology: nuclear energy and the environment*. Published by CRC Press, 1982, ISBN 0-8493-5353-X

White GC, Hakonson TE, Ahlquist AJ. *Factors affecting radionuclide availability to vegetables grown at Los Alamos*. Journal of Environmental Quality volume 10 pages 294-299 (1981).

**Appendix A: Wet Weight Concentration Ratios for Elk and Deer based on Fresquez 1996 and Fresquez 1998b.**

**Notes**

1. Deer and elk have relatively large home ranges: about 100 ha for deer and 1,000 ha for elk. Therefore, it is assumed that the average LANL soil concentrations reported in Fresquez 1996 are representative for the deer and elk of Fresquez 1998b. There are some obvious exceptions. For example, two deer were road-kill near DP Canyon and had probably been foraging on contaminated land that is not represented by the soil data of Fresquez 1996. However, to be conservative, all data have been included in the averages.

2. The soil data extend from 1974 to 1994, whereas the deer and elk data extend from 1991 to 1998. Therefore, in the tables, the "corrected average" includes a correction factor of 1.26 for ten years of radioactive decay.

<b>Strontium-90</b>	pCi/g dry	dry/wet	pCi/g wet	pCi/g soil	ratio
Deer muscle LANL	0.049	0.25	0.0123	0.6	0.02
Deer muscle regional	0.014	0.25	0.0035	0.32	0.01
elk muscle LANL	0.025	0.26	0.0065	0.6	0.01
elk muscle regional	0.001	0.26	0.0003	0.32	0.00
average	0.022	0.26	0.0056	0.46	0.01
<b>SR90 corrected average</b>					<b>0.01</b>

<b>Strontium-90</b>	pCi/g dry	dry/wet	pCi/g wet	pCi/g soil	ratio
deer bone LANL	2.6	0.7	1.82	0.6	3.0
deer bone regional	0.9	0.7	0.64	0.32	2.0
elk bone LANL	1.6	0.79	1.26	0.6	2.1
elk bone regional	1.3	0.79	1.03	0.32	3.2
average	1.6	0.75	1.19	0.46	2.6
<b>SR90 corrected average</b>					<b>3.3</b>

<b>Cesium-137</b>	pCi/g dry	dry/wet	pCi/g wet	pCi/g soil	ratio
Deer muscle LANL	0.068	0.25	0.0170	0.6	0.03
Deer muscle regional	0.015	0.25	0.00375	0.43	0.01
elk muscle LANL	0.035	0.26	0.0091	0.6	0.02
elk muscle regional	0.095	0.26	0.0247	0.43	0.06
average	0.053	0.26	0.0136	0.515	0.03
<b>Cs137 corrected average</b>					<b>0.04</b>

<b>Cesium-137</b>	pCi/g dry	dry/wet	pCi/g wet	pCi/g soil	ratio
deer bone LANL	0.024	0.7	0.017	0.6	0.03
deer bone regional	0.010	0.7	0.007	0.43	0.02
elk bone LANL	0.032	0.79	0.025	0.6	0.04
elk bone regional	0.043	0.79	0.034	0.43	0.08
average	0.027	0.75	0.021	0.515	0.04
<b>Cs137 corrected average</b>					<b>0.05</b>

