

111612



**STRATIGRAPHIC AND STRUCTURAL REVISIONS
TO THE GEOLOGIC MAP OF THE WHITE ROCK
7.5-MINUTE QUADRANGLE, LOS ALAMOS AND
SANTA FE COUNTIES, NEW MEXICO**

BY

DAVID P. DETHIER¹ AND DANIEL J. KONING²

May, 2007

¹ Department of Geosciences, Clark Hall, Williams College, 947 Main Street, Williamstown, MA 01267; david.p.dethier@williams.edu. Mapped Pliocene, Quaternary, and outcrops of undivided Santa Fe group deposits in Dethier (1997).

² New Mexico Bureau of Geology and Mineral Resources, New Mexico Tech, 801 Leroy Place, Socorro, NM 87801-4796; dkoning@nmt.edu. Differentiated the undivided Santa Fe Group deposits of Dethier (1997) into several lithostratigraphic units and studied the Ancha Formation on this quadrangle.

34354



INTRODUCTORY REMARKS

This work constitutes a revision to the Geology of the White Rock quadrangle (Dethier, 1997) in the following ways. One, we subdivide the Santa Fe Group into seven lithostratigraphic units. Pliocene and Quaternary units remain the same except for changes in the description of the Ancha Formation and slight changes of mapped contacts near Buckman. We also include detailed stratigraphic sections of Santa Fe Group sediment (Appendix 1). Unit descriptions from Dethier (1997) are included here for completeness. Two, we show newly discovered faults in the Buckman area and White Rock gorge. For this report, we present an introduction to the geologic features and stratigraphy of the area followed by the unit descriptions. We then briefly discuss some of the newly discovered faults in the Buckman area and summarize the paleogeographic changes that occurred during the middle to late Miocene. We refer the reader to Dethier (1997) for more detailed discussion on Pliocene aggradation and canyon cuttings, Pliocene volcanism, and the Quaternary evolution of White Rock Canyon.

The prominent physiographic feature and namesake of the quadrangle is the White Rock Gorge, through which flows the Rio Grande. Immediately east of this gorge, at the mouth of Cañada Ancha, lies the location of the former town of Buckman, the namesake of the adjacent well field that supplies 15-50% of the water used by the burgeoning city of Santa Fe. (Figure 1). Buckman lies about 24 km northwest of downtown Santa Fe. The town of White Rock is located only 3 km to the southwest of Buckman, but perched 210 m above it on the east edge of the Pajarito Plateau

Both Buckman and White Rock lie in the south-central Española Basin. The Española Basin is one of many basins near the Rio Grande in New Mexico formed by tectonism associated with the Rio Grande rift (Kelley, 1956; Spiegel and Baldwin, 1963; Chapin, 1971). Like these other basins, the Española Basin is filled by siliciclastic sediment (primarily sand, with lesser mud and gravel) and volcanic rocks of the Santa Fe Group of Spiegel and Baldwin (1963), which ranges in age from upper Oligocene through upper Miocene (Smith, 2004; Koning et al., 2004a). In our study area, Santa Fe Group strata may be as thick as 3000 m (Biehler et al., 1991). Two formations have been assigned to the Santa Fe Group by Galusha and Blick (1971): the Chamita Formation underlain by the Tesuque Formation. The Chamita Formation was subdivided into five members by Koning and Aby (2005), four of which are applicable here: the Vallito, Hernandez, Cejita, and Cuarteles Members. Note that the Cejita and Cuarteles Members extend into both the Tesuque and Chamita Formations, as allowed by Article 25 of the North American Stratigraphic Code (North American Commission on Stratigraphic Nomenclature, 2005), with those members in the Chamita Formation being restricted to west of the Rio Grande (Koning and Aby, 2005; Koning et al., 2005a). The Tesuque Formation was originally subdivided into several member-rank units that included the Chama-El Rito, Ojo Caliente Sandstone, Pojoaque, Skull Ridge, and Nambe Members (Galusha and Blick, 1971). Later workers have subdivided the Tesuque Formation based on provenance and paleocurrents. Applicable to our study are lithosomes A and B of Cavazza (1986) and the Cejita Member of Manley (1976, 1977, and 1979). In the study

area, the Santa Fe Group is unconformably overlain by coarse-grained sand and gravel of the Puye Formation, which is Pliocene-age and includes volcanoclastic alluvial fan deposits derived from the Jemez Mountains intercalated with axial river deposits of the Pliocene-age Rio Grande (Griggs, 1964; Bailey et al., 1969; Waresback, 1986; Turbeville et al., 1989; Waresback and Turbeville, 1990). Overlying the Puye Formation, and locally interbedded with it (Brxton and Vaniman, 2005), are basalt flows of the Cerros del Rio volcanic field (primarily 2.3-2.8 Ma; WoldeGabriel et al., 1996; Sawyer et al., 2002). The two ash-flow members of the Bandelier Tuff, having interpreted ages of 1.2 and 1.6 Ma (Izett and Obradovich, 1994; Spell et al., 1996), cap the Pajarito Plateau and are mostly found west of the Rio Grande in the study area, with only minor remnants on the east side of the river along White Rock Canyon. An upcoming paper in the 2007 NMGS Guidebook by Koning et al. (in press) discusses the subsurface descriptions and stratigraphic relations of Miocene-age sediment in the Buckman and White Rock areas, as well as various interpretations relating to Miocene-age sedimentation and tectonics.

DESCRIPTION OF MAP UNITS

Grain sizes follow the Udden-Wentworth scale for clastic sediments (Udden, 1914; Wentworth, 1922) and are based on field estimates. Pebbles are subdivided as shown in Compton (1985). The term "clast(s)" refers to the grain size fraction greater than 2 mm in diameter. Clast percentages are determined using either clast counts at select outcrops or visual estimation with the aid of percentage charts. Descriptions of bedding thickness follow Ingram (1954). Colors of unconsolidated sediment are based on visual comparison of dry samples to the Munsell Soil Color Charts (Munsell Color, 1994). Surficial units are delineated on the map where they are estimated to be at least 1 m thick. Soil horizon designations and descriptive terms follow those of the Soil Survey Staff (1992) and Birkeland (1999). Stages of pedogenic calcium carbonate morphology follow those of Gile et al. (1966), Machette (1985), and Birkeland (1999).

Use of the term "amphibolite" may also include minor mafic-rich rocks such as gabbro, diorite, and pyroxene- or hornblende-gneiss. However, most of the clasts referred to as "amphibolite" appear to be true amphibolite in hand sample. The term "granitic clasts" includes foliated granite as well as feldspar and quartz clasts that were probably derived from the weathering of coarse-grained granite rock.

QUATERNARY DEPOSITS

Qa₃ **Alluvial deposit (Holocene)** – Cross-bedded to planar-bedded sand and pebbly sand and thin beds of silty sand exposed along the Rio Grande, along active channels of tributary arroyos, and on adjacent low (<4 m; <13 ft) surfaces. Beds generally <0.5 m (<1.5 ft) thick. Includes areas of late Pleistocene deposits in uplands west of the Rio Grande. Exposed thickness 2-4 m (6.5-13 ft). Actual thickness probably >10 m (>33 ft) along the Rio Grande and Cañada Ancha. Base

not exposed. Soils thin (<0.5 m; <1.5 ft) with A/C or Bw horizons on low terraces west of the Rio Grande; stage II carbonate in the eastern part of the area.

- Qa₂** **Alluvial deposit (Holocene and late Pleistocene)** – Well-sorted cobble to boulder gravel, cross-bedded sand, and thin-bedded sand and silty sand beneath terraces along the Rio Grande; cobble gravel along Los Alamos Canyon and other steep tributaries; sand and silty sand near Cañada Ancha; and gravelly sand and eolian silt on upland parts of the Cerros del Rio. Lies beneath surfaces 5-20 m (16-66 ft) above present drainages. Thickness probably 4-15 m (13-50 ft). Overlies and truncates rocks of the Santa Fe Group, landslide deposits, and older alluvium. Soils are 0.5-1.0 m (1.5-3 ft) thick and display stage II or stage III carbonate at dry sites and Bw or Bt horizons at wetter sites.
- Qa₁** **Alluvial and lacustrine(?) deposit (middle? Pleistocene)** – Well-sorted cobble to boulder gravel and rhythmically bedded fine sand, silt, and silty clay exposed beneath terrace remnants 25-45 m (80-150 ft) above the Rio Grande. Contact relations best exposed near the mouth of Cañada Ancha, where 15 m (50 ft) of gravelly deposits fill channels cut into the Santa Fe Group (Section E of Dethier, 1997). Laminated to thin-bedded sediment 15-20 m (50-66 ft) thick lies over the gravel. Thickness mainly 10-30 m (33-100 ft). Overlies and truncates Santa Fe Group and landslide deposits. One surface west of Chino Mesa is overlain by El Cajete tephra (~50-60 ka; Reneau et al., 1996). Soil at that locality is 0.8-1.3 m (2.5-4 ft) thick and contains weak stage IV carbonate; soils on other remnants are thinner, and carbonate development is stage III.
- Qf₂** **Alluvial-fan deposit (Holocene and late Pleistocene)** – Cross-bedded, poorly sorted cobble to boulder gravel deposited mainly along White Rock Canyon by high-gradient, ephemeral channels. Includes debris flows along channels that drain Bandelier Tuff. Most small fan deposits west of the Rio Grande are mapped as Qa₃ or Qc. Thickness 4-8 m (13-26 ft). Generally overlies landslide deposits. Interfingers with and merges laterally with alluvial deposits. Soils are thin to absent on active areas of fans and <0.8 m (<2.5 ft) thick with stage II or weak stage III carbonate elsewhere.
- Qf₁** **Alluvial-fan deposit (late to middle Pleistocene)** – Cross-bedded, poorly sorted cobble to boulder gravel and poorly sorted, matrix-rich debris-flow deposits that form remnants 5-15 m (16-50 ft) above adjacent Qf₂ deposits. Thickness 4-15 m (13-50 ft), but most contacts obscured by colluvium. Overlies landslide deposits and Santa Fe Group. Overlain by El Cajete tephra (~50-60 ka; Reneau et al., 1996) along Water Canyon. Soils poorly exposed but locally thicker than 1.0 m (3 ft), exposing stage III or weak stage IV carbonate.
- Qc** **Colluvial deposit (Holocene to middle Pleistocene)** – Rockfall, debris-flow, and poorly sorted alluvial deposits at the bases of cliffs, particularly along White Rock Canyon. Texture and clast lithology variable. Includes thin (<2 m; <6.5 ft) alluvial deposits west of the Rio Grande and, in places, El Cajete tephra (~50-60 ka;

Reneau et al., 1996). Thickness generally 4-10 m (13-33 ft) but locally along White Rock Canyon exceeds 25 m (82 ft). Overlies Bandelier Tuff west of White Rock; elsewhere overlies basaltic flows, phreatomagmatic deposits, and landslides. Soils thin (<0.5 m; <1.5 ft) and weakly developed at most locations, but lenses of El Cajete tephra, strong Bt (west of Rio Grande) or K horizons (White Rock Canyon and east), and buried soils (Birkeland, 1999) demonstrate that deposits are polygenetic and locally older than late Pleistocene.

Qse **Thin slopewash and eolian deposit (Holocene to late Pleistocene)** – Light yellowish brown to pale brown to brownish yellow, silt and very fine- to medium-grained sand mixed with varying amounts of coarser sand and gravel. This thin unit caps high-level surfaces. The underlying unit is also shown. Weakly to moderately consolidated. Qse is less than 4 m-thick.

Qlsa **Landslide deposit (Holocene to late Pliocene)** – Massive slumps and debris flows rich in basaltic-boulder gravel mainly exposed along White Rock Canyon, north of Chaquehui Canyon, and along Cañada Ancha. Most slides are inactive.

Qls Deposits consist of massive slump blocks with coherent internal stratigraphy near canyon rims and progressively deformed slumps and debris flows closer to the Rio Grande. Upper surfaces of many slumps (near Pajarito Springs and southwest of Otowi Bridge, for example) are sites of active fluvial and colluvial processes and are covered with >4 m (>13 ft) of fluvial, eolian, and colluvial deposits. Such flat to gently sloping areas are mapped as Qlsa. Dips in massive slump blocks range from 8° to 70° toward head scarps. Failures occurred along (1) steeply dipping planes rooted in the Santa Fe Group, (2) subhorizontal planes in clayey silt layers found at several levels of Pliocene fluvial and lacustrine deposits (Ta), and (3) steep walls of maars. Limited areas of autochthonous rocks are included in areas mapped as Qls. Slide material overlies rocks of the Santa Fe Group at most sites. At several locations undisturbed late Pliocene deposits lie stratigraphically above landslide deposits (Section H of Dethier, 1997). Morphology of most failures and inclusion of Bandelier Tuff in some suggest that slides were active in early to middle Pleistocene time but that many became stable in the middle to late Pleistocene. El Cajete tephra (~50-60 ka; Reneau et al., 1996) lies on landslide deposits south of the White Rock area. Soils are generally 0.8-1.4 m (2.5-4.5 ft) thick. Carbonate morphology (Birkeland, 1999) is stage IV at some sites, but stage III carbonate is present in most exposures. Cation ratios in rock varnish (Dethier et al., 1988) at two sites suggest that those landslides stabilized >250 ka.

Santa Fe Group deposits exposed in Quaternary landslides (not shown in correlation of map units)

Qls-ccuf **Exposures of fine-grained Cuarteles Member (Chamita Formation) involved with landslide-related subsidence and deformation** – See description of fine-grained Cuarteles Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.

- Qls-ccu** **Exposures of Cuarteles Member (Chamita Formation) involved with landslide-related subsidence and deformation** – – See description of Cuarteles Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-tcuf** **Exposures of fine-grained Cuarteles Member (Tesuque Formation) involved with landslide-related subsidence and deformation** – See description of fine-grained Cuarteles Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-tcu** **Exposures of Cuarteles Member(Tesuque Formation) involved with landslide-related subsidence and deformation** – – See description of Cuarteles Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-cv** **Exposures of Vallito Member (Chamita Formation) involved with landslide-related subsidence and deformation** – – See description of Vallito Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-ch** **Exposures of Hernandez Member (Chamita Formation involved with landslide-related subsidence and deformation** – – See description of Hernandez Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-p** **Exposures of Puye Formation involved with landslide-related subsidence and deformation** – – See description of Puye Formation. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-pt** **Exposures of axial river deposits of the Puye Formation (Totavi Lentil) involved with landslide-related subsidence and deformation** – – See description of Puye Formation. Bedding is tilted in various directions and somewhat deformed but generally intact.
- Qls-cm** **Exposures of phreatomagmatic deposits involved with landslide-related subsidence and deformation** – – See description of unit Tcm below. Bedding is tilted in various directions and somewhat deformed but generally intact.

PLIOCENE SEDIMENTARY STRATA

- Ta** **Ancha Formation (upper Pliocene)** – Brownish yellow to light yellowish brown sand and gravel under the Cerros del Rio volcanic flows (units Tcba and Tcb₂) and phreatomagmatic deposits (Tcm); unit is also interlayered with these volcanic flows. Overlies the Puye Formation (Tp) and local basalt flows near the Rio

Grande; to the east the Ancha Formation unconformably overlies Miocene-age Cuarteles Member deposits of the Tesuque Formation (Koning and Maldonado, 2001; Koning et al., 2002). At a description site near the eastern quadrangle boundary (UTM coordinates: 3,960,550 N, 398,125 S, \pm 20 m; zone 13, NAD 27), the sediment is a gravelly sand with very thin to thin, planar to lenticular beds. The gravel here is generally poorly sorted pebbles with 5-20% cobbles; clast composition is: 1-5% quartzite, trace amphibolite, locally trace basalt or andesite clasts, and 95-99% granite. Cobbles and coarse to very coarse pebbles are commonly rounded to subrounded, and very fine to medium pebbles are subrounded to subangular. Sand is generally coarse to very coarse, subangular to subrounded, poorly to moderately sorted, and arkosic (with 15% lithics that include 0.5% volcanic and olivine grains). At a description site near Caja del Rio Canyon (UTM coordinates: 3,962,350 N, 397,150 S, \pm 20 m; zone 13, NAD 27), the sediment has 15-25% thin to thick channel-fills of sandy pebbles to pebbly sand and the gravel consists of clast-supported, granitic pebbles; the non-channel sediment is comprised primarily of slightly silty very fine- to very coarse-grained sand in thin to thick, tabular beds, or otherwise massive. Sediment is weakly to moderately consolidated, and generally non-cemented except for the bases of some channels. Unit locally includes silty and clayey lacustrine sediment near the Rio Grande. Unit also includes thin-bedded silt and silty sand, and beds of phreatomagmatic deposits and debris flows exposed along Cañada Ancha and north of Water Canyon, west of the Rio Grande. Paleocurrent directions measured on channels and gravelly crossbeds range from 180° to 270° and average about 220°. Unit is correlated with the Ancha Formation in the Santa Fe embayment to the south (Koning et al., 2002; Koning and Maldonado, 2001) because these units are also interbedded with lapilli and phreatomagmatic deposits of the Cerros del Rio volcanic field, and thus occupy the same stratigraphic position. Also correlates with older alluvium of Griggs (1964). Contains lenses of dacitic tephra along Cañada Ancha. Manley (1976) reported an age of 2.7 Ma of one lense some 8 km (5 mi) east of Caja del Rio Canyon. Age elsewhere is likely Pliocene considering stratigraphic relations with Pliocene-age Cerros del Rio volcanic flows and phreatomagmatic deposits. Thickness of 1-30 meters (commonly less than 5 m, although generally not mapped where it is this thin).

Tp Puye Formation fanglomerate (Pliocene; Griggs, 1964) – Weakly lithified pebble to boulder gravel, boulder-rich debris flows, massive to planar-bedded sand, thin (<1 m; <3 ft) beds of dacitic tephra and pumiceous alluvium, and beds of fine sand and silt. Exposed west of the Rio Grande except for an isolated outcrop south of the mouth of Cañada Ancha (Section F of Dethier, 1997). Gravel beds generally 0.5-3.0 m (1.5-10 ft) thick. Debris flows range from 0.3 m (1 ft) to about 5.0 m (16 ft) thick. Clast and matrix lithology mainly dacite derived from the Tschicoma Formation of the Jemez Mountains, but Precambrian material composes >30% of some fluvial units. Thickness 5-30 m (16-100 ft). Fills channels cut in rocks of the undivided Santa Fe Group and, along Los Alamos Canyon, into cobble gravel of the Puye Formation, Totavi Lentil (Griggs, 1964). Exposed beneath quartzite-rich cobble gravel (Totavi Lentil) or phreatomagmatic

deposits at most locations. Paleocurrent directions measured on channels, gravelly crossbeds, and imbricated cobbles range from about 90° to 200° and average about 150°, slightly south of the trend of present canyons. Faulted locally near the mouth of Ancho Canyon; otherwise undeformed. Pumiceous Puye gravel 8 km (5 mi) north-northwest of Otowi Bridge gave a fission-track age of 2.9 Ma (Table 1 of Dethier, 1997). Turbeville et al. (1989) report that the upper part of the Puye Formation may be as young as 1.7±0.1 Ma northwest of the White Rock quadrangle.

Tpt Puye Formation, ancestral Rio Grande facies (Pliocene; Totavi Lentil of Griggs, 1964) – Slightly lithified pebble to cobble gravel rich in clasts of Precambrian rock, sand, and thin beds of silty sand west of the Rio Grande and at one outcrop south of the mouth of Cañada Ancha, east of the Rio Grande (Section F of Dethier, 1997). Coarse units are 0.5-3.0 m (1.5-10 ft) thick, cross-bedded, and locally planar bedded. Clasts generally >80% quartzite and other resistant lithologies from northern New Mexico, but clasts from the southern Sangre de Cristo Range are common locally. Thickness 5-45 m (16-150 ft). Maximum thickness and most extensive exposures in Sandia and Mortandad Canyons (Section D of Dethier, 1997). Fills channels in and locally interbedded with Puye fanglomerate except along Los Alamos Canyon, where it overlies Santa Fe Group. Lies beneath landslide deposits, Pliocene alluvium, or phreatomagmatic deposits at most exposures. Paleocurrent directions measured on channels, gravelly crossbeds, and imbricated cobbles range from 160° to 220° and average about 180°. Mainly undeformed. Near the mouth of Ancho Canyon (Section J of Dethier, 1997) lies under a basaltic flow (Tcb₂) reported by Bachman and Mehnert (1978) to have an age of about 2.6 Ma (Table 1 of Dethier, 1997).

PLEISTOCENE-PLIOCENE VOLCANIC DEPOSITS AND ROCKS

Qct El Cajete tephra (late? Pleistocene) – Pumiceous lapilli of rhyolitic composition. Forms surface layer in isolated outcrops as thick as 40 cm in southwest part of map area and <5 cm in northeast part of map area. Most exposures have been reworked by slope processes. Derived from the El Cajete vent in the Valles caldera; previously thought to be about 150,000 yrs old (Self et al., 1988) but may be as young as 50 ka (Reneau et al., 1996).

Qbtu Bandelier Tuff, upper part (early Pleistocene) – Slightly welded pyroclastic flows (Tshirege Member) and a thin (<1 m; <3 ft) pumiceous fall unit (Tsankawi Pumice Bed), both of rhyolitic composition. Dominates surfaces west of the Rio Grande and forms one prominent outcrop east of the Rio Grande (Section H of Dethier, 1997). Two to five pyroclastic flows separated by pumice concentrations or thin, sorted partings are present along deep canyons west of the Rio Grande. Thickness generally <60 m (<200 ft) but about 90 m (300 ft) east of the Rio Grande. As mapped, may include exposures of lower Bandelier Tuff, which it lies above. Paleoflow direction to the east. Derived from the Valles caldera west of the map area. Age 1.2 Ma (Izett and Obradovich, 1994).

Qbt1 Bandelier Tuff, lower part (early Pleistocene) – Slightly welded pyroclastic flows (Otowi Member) of pumiceous rhyolite and a compound pumiceous fall unit (Guaje Member) as thick as 6 m (20 ft), also of rhyolitic composition. Best exposed in deep canyons west of the Rio Grande, particularly in Los Alamos (Section A of Dethier, 1997) and Sandia Canyons. One or two thick flows overlie the Guaje Member, which is absent at many exposures. Maximum thickness about 50 m (165 ft). Lies beneath upper Bandelier Tuff. Fills canyons as deep as 50 m (165 ft) cut into tholeiitic olivine basalt (Tcb₃), basaltic andesite (Tcba), and phreatomagmatic deposits. Derived from the Valles caldera west of the map area. Age is 1.6 Ma (Izett and Obradovich, 1994).

Tcc Cinder cone deposit (Pliocene) – Oxidized cinders, agglomerate, and minor areas of phreatomagmatic deposits composed of olivine basaltic andesite and basalt. Contains 2-8% quartz xenocrysts. Exposed mainly along the Rio Grande and at the Caja del Rio Canyon. Granular surface deposits, dissected cinder cones, and slightly lithified exposures in canyon walls; best exposed near La Mesita (Section C of Dethier, 1997). Massive to planar bedded, locally rich in lava and accidental bombs <0.2 m (<0.5 ft) in diameter. Maximum thickness about 60 m (197 ft). Lies above phreatomagmatic deposits at many exposures. Age not closely bounded but probably late Pliocene.

Tcm Phreatomagmatic deposit (Pliocene) – Bedded to massive fall, surge, and flow deposits composed of basaltic tuff and cinders and accidental fragments of the Santa Fe Group. Thickest exposures along the Rio Grande and Cañada Ancha (Section G of Dethier, 1997). Fall beds, 0.3-3.0 m (1-10 ft) thick, are composed mainly of ash and lapilli containing sparse bombs of accidental fragments and basaltic fragments. Surge beds are planar and crossbedded, locally rippled, coarse silt to pebbly sand, generally 0.1-0.4 m (0.3-1.3 ft) thick. Flow deposits are mainly matrix-rich pebble to boulder gravel in discontinuous beds 1-4 m (3-13 ft) thick. Near maars the concentration of accidental fragments decreases upsection. Locally sheared, slumped, or brecciated. As much as 60 m (200 ft) thick near maars such as La Mesita (Section C of Dethier, 1997), "Buckman maar" (Section G of Dethier, 1997; Aubele, 1978), and "Montoso maar" (Aubele, 1978). Generally lies above Puye Formation, undivided Santa Fe Group, and interlayered basalt and phreatomagmatic deposits. Lies beneath flows of basaltic andesite, basalt, or cinder and agglomerate deposits. In Ancho Canyon (Section J of Dethier, 1997), lies on Tcb₂, which gave an age of 2.6 Ma (Table 1 of Dethier, 1997), but stratigraphic relations suggest deposits along Cañada Ancha may be somewhat older. Minimum age not well known.

Tcbm Basalt and interlayered phreatomagmatic deposits (Pliocene) – Thin (<10 m; <33 ft) basaltic flows interlayered with basaltic phreatomagmatic rocks, mainly surge and flow deposits, and mapped in the southern part of the map area near the Rio Grande. Multiple baked layers exposed in Chaquehui Canyon (Section K of Dethier, 1997) and along the north margin of Chino Mesa (Section I of Dethier, 1997). Maximum thickness about 50 m (165 ft). Lies on phreatomagmatic

deposits and Santa Fe Group. Lies beneath phreatomagmatic deposits and basalt flows. One flow in Chaquehui Canyon gave an age of 2.78 ± 0.04 Ma (Table 1 of Dethier, 1997).

Tci Basaltic intrusion (Pliocene) – Olivine, pyroxene basaltic andesite containing quartz xenocrysts. Forms fine-grained dikes and small, shallow intrusions, with sharp chilled margins, associated with maars and cinder cones along the east margin of White Rock Canyon. Most prominent intrusion is a small plug capped with agglomerate at the west edge of Chino Mesa. Aubele (1978) suggested that a sill lies beneath cinders and agglomerate along the west margin of Sagebrush Flats, and the andesitic exposure in Water Canyon (location 100) may be intrusive. Dikes generally vertical and <10 m (<33 ft) wide. Unit intrudes phreatomagmatic or cinder-cone deposits. Not dated but probably middle or late Pliocene.

Tca Andesite (early Pleistocene? and late Pliocene) – Massive, steep-sided flows, agglomerate, and domes of hypersthene andesite (Table 2) exposed in the southeast part of the map area. Strongly sheeted near flow surface; brecciated at bases. Thickness of flows >20 m (>66 ft). Total thickness unknown but likely exceeds 150 m (500 ft) in the southeast map area. Lies above basaltic andesite and hawaiite and lies beneath Quaternary Guaje Pumice Bed (Qbt1). Manley (1976) reported an age of about 2 Ma (Table 1 of Dethier, 1997) from an andesite flow 5 km (3 mi) east of the eastern edge of the map area. Age of youngest domes unknown.

Tcb₃ Basalt (late Pliocene) – Tholeiitic olivine basalt flows, pillow basalt, and palagonitic breccia exposed west of the Rio Grande north of Chaquehui Canyon. Flows thin (<10 m; <33 ft) with sharp to rubble-rich contacts. Subaerial thickness <30 m (<100 ft); maximum thickness ~ 80 m (~ 260 ft) in Mortandad Canyon (Section D of Dethier, 1997). Overlies lacustrine or sandy fluvial sediment (Ta) north of White Rock and older basaltic flows to the south (Section J of Dethier, 1997). Thin (<3 m; <10 ft) fluvial and lacustrine deposits separate the basalt from overlying lower Bandelier Tuff at some sites (Sandia Canyon, for instance). Flow directions measured on foreset deposits of pillow basalt in Los Alamos, Sandia, Mortandad, Buey, and an unnamed canyon range from about 70° to 150° (average about 110°). Topset-foreset contacts in deltas of basaltic debris at $6,200 \pm 25$ ft elevation suggest that a lake dammed near Chaquehui Canyon persisted in White Rock Canyon during eruption of much of the basalt. Basalt originated from vents buried by Bandelier Tuff west and northwest of the map area. A sample from the basalt exposed near the intersection of Pueblo and Los Alamos Canyons (Section A) in the northwest map area gave an Ar/Ar age of 2.33 ± 0.08 Ma (Table 1 of Dethier, 1997), and flows capping Ancho Canyon and underlying White Rock gave ages between 2.5 and 2.4 Ma.

Tcba Basaltic andesite and related flows (Pliocene) – Massive flows of olivine-hypersthene basaltic andesite, andesite, and hawaiite, containing as much as 7% quartz, exposed along the Rio Grande. Forms thick (>30 m; >100 ft locally) flows

with brecciated bases and cooling joints 4 m (13 ft) in diameter. As much as 170 m (560 ft) thick along the Rio Grande at the south end of the map area. Fills canyons as deep as 40 m (130 ft) southwest of Otowi Bridge (Section B of Dethier, 1997), where flows lie on top of phreatomagmatic deposits (Tcm) and the Puye Formation (Tp and Tpt) and lie beneath fluvial and lacustrine sediment (Ta). Overlies basalt (Tcb₂), phreatomagmatic deposits and agglomerate, and the Puye Formation south of Water Canyon west of the Rio Grande (Sections J and K of Dethier, 1997) and south of Sagebrush Flats east of the Rio Grande. Outcrop pattern and channel orientation suggest that lava flowed mainly southwest and south from vents near La Mesita, on Sagebrush Flats between Water and Ancho Canyons, and probably in the vicinity of Chino Mesa (Aubele, 1978). The Otowi flow of Galusha and Blick (1971) gave an age of 2.57±0.02 Ma (location 145a), and a massive flow probably derived from La Mesita (location 145c) gave an age of 2.55±0.02 Ma. The flow between Water and Ancho Canyons gave an Ar/Ar age of 2.3±0.3 Ma. Age bracketed by the 2.6±0.4 Ma age of stratigraphically lower Tcb₂ near Ancho Canyon (Table 1 of Dethier, 1997) and the 2.33 Ma age of overlying Tcb₃ along Los Alamos Canyon (Table 1 of Dethier, 1997).

Tcb₂ Basalt flows (Pliocene) – Thin (<10 m; <33 ft) flows of olivine basalt (hawaiite, see Table 2 of Dethier, 1997) containing <5% quartz xenocrysts and exposed mainly beneath large areas of Sagebrush Flats (Sections F and G of Dethier, 1997) and along the east boundary of White Rock quadrangle. Brecciated and sparse baked zones between flows. Maximum thickness ~70 m (~230 ft) near Chino Mesa. Overlies phreatomagmatic deposits (Tcm) and Pliocene alluvial deposits (Ta) along Cañada Ancha, at the White Rock Overlook, and in Pajarito Canyon. A latite-andesite flow (Baldrige, 1979) included with Tcb₂ lies above the Totavi Lentil (Tpt) in Ancho Canyon. Andesite and basaltic-andesite flows (Tca, Tcba) and Quaternary alluvium lie above hawaiite in the Sagebrush Flats area, whereas Tcb₂ flows lie beneath basaltic andesite (Tcba) or olivine basalt (Tcb₃) at most exposures west of the Rio Grande. Probable vent areas are marked by cinder and agglomerate deposits from Sagebrush Flats south to Chino Mesa; younger basaltic and andesitic flows probably cover other vents. A flow exposed on the south rim of Caja del Rio Canyon (location 122) gave an age of 2.49±0.03 Ma (Table 1 of Dethier, 1997). An age of about 2.6 Ma (Table 1 of Dethier, 1997) was obtained by Bachman and Mehnert (1978) in Ancho Canyon.

Tcb₁ Basaltic flows (Pliocene?) – Olivine basalt flows and more evolved rocks exposed within 20 m (66 ft) of the present level of the Rio Grande and Cañada Ancha. Flows are thin to massive and include a lava lake (Caja del Rio Canyon) and a number of poorly exposed outcrops along the Rio Grande from Water Canyon to Chaquehui Canyon. Thickness 20-95 m (66-310 ft). The latter flows lie above deformed phreatomagmatic deposits (Tcm) and beneath cinders and agglomerate (Tcc). Sources of the flows are buried, probably diverse (Table 2 of Dethier et al., 1997), and may lie in a stratigraphic position similar to subsurface volcanic rocks reported by Griggs (1964), some as old as middle Miocene

(WoldeGabriel et al., 1996). A flow near the mouth of Water Canyon gave an age of 2.50 ± 0.04 (Table 1 of Dethier, 1997).

Tcbu Basalt, undivided (Pliocene) – Olivine basalt, hawaiite, and basaltic andesite containing 2-8% quartz xenocrysts and exposed in the southern part of the map area (Section I of Dethier, 1997). Thin to massive flows. Thickness from 20 m to >150 m (from 66 ft to >490 ft) in the southern map area. Lies above phreatomagmatic deposits (Tcm). Generally lies beneath Tcba and may be equivalent, in part, to basalt mapped as Tcb₂. Montoso maar (Aubele, 1978) was the probable source for much of this unit. Age unknown but probably middle to late Pliocene.

MIDDLE-LATE MIOCENE BASIN-FILL DEPOSITS (SANTA FE GROUP)

The Santa Fe Group in the White Rock quadrangle is composed of fluvial deposits associated with basin-floor and alluvial-slope (also known as piedmont slope) drainages. The fluvial deposits consist of sandy-gravelly channel-fills intercalated with floodplain deposits of clay, silt, very fine- to fine-grained sand, and silty fine sand. Among these channel-fills and floodplain deposits, we differentiate several lithostratigraphic units based on standard sedimentologic properties, particularly composition and texture but to a lesser extent bedding characteristics, color, and paleocurrent directions. These are generally member-rank units that belong to the Chamita and Tesuque Formations, as defined by Koning and Aby (2005) and Koning et al. (2005a) from initial definitions of Galusha and Blick (1971) and Spiegel and Baldwin (1963). Note that the Vallito Member of the Chamita Formation extends east of the Rio Grande in the study area. We also extend the Vallito Member down-section south of Española to include sandy fluvial strata coeval with eolian strata of the Ojo Caliente Sandstone of the Tesuque Formation. Consistent with the nomenclature of Koning and Aby (2005), the Cejita and Cuarteles Members extend into both the Chamita Formation (west of the Rio Grande) and the Tesuque Formation (east of the Rio Grande), as allowed by the North American Stratigraphic Code (North American Commission on Stratigraphic Nomenclature, 2005). Note that some details of the sand descriptions may change with further work using thin sections and grain counts.

Age control for Santa Fe Group

The ages of Santa Fe Group strata in the study area appear to range from ca. 13.5-8.5 Ma based on the following data. A sharp gamma ray high at 1346-1348 ft in the Buckman-9 well is interpreted as an ash. This ash is located 13 ft below the Cejita Member – Pojoaque Member contact. Near Española, ashes at this stratigraphic position have been assigned to the Pojoaque white ash zone (e.g., unit 5a of the Cuarteles section in Koning and Manley, 2003, and Koning et al., 2005a), which is interpreted to have an age range of 14.0-13.2 Ma (Koning et al., 2005a; Koning, 2002; Barghoorn, 1981; Izett and Obradovich, 2001). Since this particular bed is likely at the top of the Pojoaque white ash zone, it probably has an age of ca. 13.2 Ma (based on interpretations of Koning et al.,

2005a, using magnetic-polarity stratigraphy data from Barghoorn, 1981, and the revised geomagnetic polarity time scale of Cande and Kent, 1995).

Direct age control is provided by a coarse white ash-lapilli bed at Buckman. Denoted on the geologic map as the “coarse white ash marker bed” (see also map inset of Fig. 2), this tephra bed is found in the western Buckman well field. We interpret that it extends to the top of the Lower Buckman section (Figures 2-3). $^{40}\text{Ar}/^{39}\text{Ar}$ analyses on biotite grains from this tephra bed yield an isochron age of 10.9 ± 0.2 Ma (William McIntosh and Steve Cather, unpublished data for lab #6240).

The basalt flow encountered at 586-671 ft in the R-10 well (see cross-section) is interpreted to be 8.5-9.0 Ma because it lies at about the same stratigraphic level as basalt flows in the nearby R-9 well that were dated at $8.45-8.63 \pm 0.24$ Ma by $^{40}\text{Ar}/^{39}\text{Ar}$ methods (Broxton et al., 2001). This flow also appears to project to a basalt in the R-22 well that returned an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 8.97 ± 0.11 Ma (G. WoldeGabriel, personal communication, 2003).

Tcvwp Chamita Formation, Vallito Member interbedded with western piedmont deposits (upper Miocene) – Only found in outcrop at the mouth of Ancha Canyon. Light brown to reddish yellow (7.5YR 6/4-6) silt, very fine- to fine-grained sandstone, and minor clay intercalated with ~25% coarse-grained channel fills. The channel-fills are >10 m wide, 1 to 2 m thick, and consist of pebbly sand with lesser sandy pebble conglomerate. The sand is commonly planar-laminated. Pebbly beds are thin to medium and planar to lenticular. Pebbles are subrounded to rounded, poorly to moderately sorted, and clast-supported. Gravel is composed of felsic to intermediate volcanic rocks with 0.5-1% quartzite; the volcanic rocks have a large amount of monolithic dacite presumed to be derived from the Jemez Mountains. Channel-fill sand is fine- to very coarse-grained and moderately to poorly sorted. Deposits are non- to weakly cemented. Greater than 30 m-thick.

Ttcuf Tesuque Formation, fine-grained Cuarteles Member east of Rio Grande (Upper Miocene) – Pink to light brown to reddish yellow to light yellowish brown, silty very fine- to fine-grained sandstone, siltstone, and mudstone. Coarse channel-fills of pebbly sandstone and sandy-pebble-conglomerate comprise about 3-25% of the unit. Fine-grained sediment outside of coarse-grained channel-fills is in very thin to thick, tabular beds that may be internally laminated. 1-5% muscovite flakes locally are present. Sparse cross-laminations up to 40 cm-tall may possibly represent eolian dunes. Sand is generally very fine- to medium-grained, an arkose, well sorted, and subangular to subrounded. There is 0.5 to 1%, very thin to thin, brown to light brown (7.5YR 5-6/4) claystone beds. Coarse-grained channel-fills are scattered, lenticular to ribbon-shaped, and up to 10 to 100 cm-thick; locally they these channel-fills are stacked as to form thicker complexes. The internal bedding of the channels is very thin to thin, and planar to lenticular to cross-stratified. The pebbles are moderately to poorly sorted, subangular to subrounded, and granitic (with trace to 1% yellowish Paleozoic

siltstone and limestone, quartzite, and gneiss). Channel-fill sand is fine- to very coarse-grained, subangular to subrounded, poorly to well sorted, and an arkose. Individual channels are up to 35 cm-deep and have westward-orientated paleoflow indicators. Channels-fills may fine-upward from pebbly to sandy sediment; sorting may become better upwards as well. Very thin to medium, lenticular, isolated channels also are present within the fine-grained sediment (generally 3-10% of the volume). Isolated channels tend to be strongly to moderately cemented, whereas laterally extensive and thick channel complexes are generally not as cemented. The finer, non-channelized sediment of the unit is moderately to well consolidated and weakly cemented by calcium carbonate. Unit possesses 1-3% weakly developed paleosols with reddish Bw horizons 20- to 30-cm thick. Interpreted to represent a generally low-energy alluvial environment where the distal alluvial slope transitioned to the flat basin floor. This unit grades eastward and upward into coarser-grained Cuarteles Member. In the subsurface, this unit interfingers westward with the Cejita Member (Tesuque Formation) and the Vallito Member (Chamita Formation). Unit progrades westward over the Vallito Member. Up to 50 m-thick.

Ttcuf Chamita Formation, fine-grained Cuarteles Member west of the Rio Grande (Upper Miocene) – See description above. Here, the Cuarteles Member interfingers westward with Vallito Member and the Hernandez Member of the Chamita Formation.

Ttcu Tesuque Formation, Cuarteles Member (Upper Miocene) – Sandy pebble-conglomerate and pebbly sandstone comprising coarse channel-fill complexes, with subordinate slightly clayey-silty sand containing <5% pebbles that was deposited by lower-energy flow outside of confined channels – the latter is referred to below as the “finer sediment” of this unit (and as “extra-channel sediment” in Appendix 1). Interbedded within this unit are sparse beds of black coarse ash and white coarse ash-lapilli, which are described below. Channel-fill complexes may be up to ~ 2 m thick and commonly very pale brown to light yellowish brown to pink in color. Bedding within the channel-fill complexes (from most to least common) is lenticular, planar, U-shaped, and cross-stratified (up to 20 cm-thick foresets), and generally the beds are very thin to medium (with internal planar-laminations); bedding becomes more planar westward where this unit grades into Ttcuf. Margins of individual channels are up to 90 cm tall but commonly 20-40 cm tall. Channel complexes may fine-upwards from gravel- to sand-dominated sediment. Pebble conglomerate is clast-supported and has 0 to 5% cobbles (average size of largest clasts are 11 x 8 cm). Pebbles are poorly to moderately sorted and mostly subangular (some subrounded). Clasts are composed of granite with trace to 1% yellowish Paleozoic limestone and siltstone, trace quartzite, trace chert, 0.5-2% gneiss, and trace amphibolite. Channel sand ranges from fine- to very coarse-grained but mostly is medium- to very coarse-grained, angular to subangular (mostly subangular), poorly to moderately sorted, and an arkose. 1-15% of individual channel complexes are strongly to moderately

cemented by calcium carbonate, the rest is weakly to non-cemented. The non-cemented to weakly cemented channel sediment may locally have up to 5-8% (visual estimate) clay in the gravel and sand interstices, and is generally weakly to moderately consolidated; where clay is present, channel-fills have a reddish yellow color.

The finer sediment is reddish yellow to light brown to pink and composed of slightly clayey-silty to clayey-silty (visual estimate of 1-10% clay + silt) sand with <10% scattered pebbles. Very minor planar mudstone laminations are present. This sediment is generally massive but may locally be very thinly to thickly, tabular-bedded (bedding is poor) or planar-laminated. The sand is very fine- to very coarse-grained but mostly very fine- to medium-grained, poorly to moderately sorted, subangular to subrounded, and an arkose. This sediment is weakly to well consolidated. Within this sediment are 1-15% very thin to thin, lenticular beds of medium- to very coarse-grained, arkosic sandstone and granitic pebble-conglomerate. Locally, there are sparse beds of pale brown (10YR 6/3), silty very fine- to medium-grained sand that is well-sorted and arkosic. These deposits locally grade laterally into coarse deposits similar to those seen in channels, but buttress margins with older channel deposits have not been observed. The finer sediment may represent small aggradational lobes deposited on the alluvial slope at the mouths of channels. They are commonly scoured and inset by the coarse channel complexes.

Weakly developed paleosols are characteristic of this unit and commonly are preserved at or near the top of the finer sediment; less commonly, soils have also been observed on top of the coarse channel deposits. These soils are characterized by a reddish yellow (5YR 6/6), 10-30 cm-thick Bw or Bt horizons with a sharp upper contact and a gradational lower contact (over 2-6 cm). Locally, these reddish horizons are underlain by whitish to pinkish Bk horizons 10-50 cm-thick possessing Stage I carbonate morphology. On the south end of Buckman Mesa, the abundance of soils increases up-section from 1% to 5% of total sediment volume.

Unit overlies and grades westward into unit Tccuf, so it is of equivalent age or younger. 300-400 m-thick.

Ttce **Cejita Formation, Tesuque Formation (Upper Miocene)** – Only exposed on the northeast slope of Buckman Mesa, where it grades upward into fine-grained Cuarteles Member of the Tesuque Formation. Here, the Cejita Member consists of sandy and gravelly channel-fills that are extensively, very thinly to thinly cross-stratified (up to ~ 1m-thick foresets) within 1-2 m-thick channel-fills of pebbly very fine- to very coarse-grained sand and sandy pebble-cobble conglomerate. Clast lithologic types are dominated by Paleozoic sandstone, limestone, and siltstone with an estimated 10-50% granite and 5-8% quartzite. Locally, granites are the dominant lithologic type (probably due to input from

alluvial-slope tributaries from the east). There may be 10-90% pink-gray dacites and rhyolites together with light gray dacites-andesites(?), probably representing mixing and interfingering with the Vallito Member to the west. Clast imbrication is approximately due south (+/- 25°). Subsurface data also indicates that the Cejita Member grades westward into the Vallito Member (Chamita Formation) and eastward into the Cuarteles Member (Tesuque Formation). Age of base is ca, 13.2 Ma (Koning et al., 2005a). Exposed thickness of 40-45 m; subsurface thickness of 125-135 m.

Tch Chamita Formation, Hernandez Member (Upper Miocene) – Light gray channel-fills with subordinate finer-grained floodplain deposits. Amalgamated channel-fills form packages as much as 18 m-thick. These channel-fills locally fine-upwards into horizontal-bedded floodplain deposits of clay, silt, and clayey-silty very fine- to fine-grained sand. The coarse-grained channel-fills are marked by a variety of bed forms, ranging from planar to lenticular to cross-stratified. Gravels include very fine to very coarse pebbles and cobbles that are subrounded to rounded, very poorly sorted, and commonly clast-supported. Lithologic types include a high amount of gray to dark gray to greenish gray to brown dacites to andesites, with minor amounts of rhyolite, welded tuff, and less than 15% quartzite in our study area (Table 1). Locally, there is less than 10% Paleozoic sedimentary clasts and minor granitic detritus. The sand fraction has low amounts of the frosted quartz and rounded, red-brown chert and volcanic(?) sand grains observed in the Vallito Member (generally less than 15% of the sand fraction). The sand mostly contains subangular to subrounded, relatively clear quartz and plagioclase, with less than 15% orange-stained quartz + potassium feldspar, 1-15% mafics, <2% chert, <5% green quartz grains, and 3-50% volcanic grains similar in composition to the gravel fraction. In outcrops, reddish brown to brown to pale yellow floodplain deposits are minor to very minor compared to the coarse-channel-fills, but locally floodplain deposits are abundant in wells R-10 and R-16.

The Hernandez Member interfingers eastward with the Vallito Member. Locally, such as at the base of the west slope of White Rock Canyon directly across from the Buckman well field, the Hernandez Member directly interfingers eastward with the Cuarteles Member. The Hernandez Member is 27 m-thick in the Upper Buckman stratigraphic section, and attains unknown, but probably much higher, thickness values to the west.

Tcv Chamita Formation, Vallito Member (Upper Miocene) – Broad channel-fills (typically >10s of meters wide) consisting of sand-dominated sediment in horizontal to cross-stratified, laminated to very thin to medium beds. Stacked channel-fills form amalgamated packages as much as 12 m-thick. The pebble fraction rarely exceeds 40 mm in diameter; clasts are moderately sorted, subrounded, and consists largely of volcanic clasts having similar composition as

the volcanic sand grains together with minor amounts of quartzite, granite, and Paleozoic sedimentary clasts (mostly sandstone, but also local limestone and siltstone). Pebbles are generally insufficient to produce clast-supported gravel beds. Basalt lithologic types are unique to the Vallito Member only in the subsurface east of the Rio Grande, but are locally present in exposures west of the Rio Grande. Sand is very pale brown to pink to light gray, fine- to coarse-grained, and consists of subrounded (with subordinate rounded and subangular), locally frosted quartz grains. There is typically trace to 3% red to brown, rounded chert and possible volcanic grains, 1-8% mafics, and 5-20% orange-stained quartz grains together with minor potassium feldspar. Up to ~25% volcanic grains are also present, ranging from welded tuff, other tuff, rhyolite, basalt, and dacite (dacite being the most common). Other lithic grains include <5% quartzite and Paleozoic sedimentary grains. Floodplain deposits consist of very fine- to fine-grained sand and silty sand, silt, and clay in various proportions. Locally, the Vallito Member contains intervals of extensively cross-stratified sand with ~20 cm-thick foresets; these are possibly eolian deposits. Non- to weakly cemented, with only minor strong cementation. In the Buckman area, the lower contact of the Vallito Member is placed at the lowest occurrence of thick sand beds characterized by fine to coarse, subrounded to rounded, locally frosted quartz with minor rounded, red-brown chert or volcanic grains and orange-stained quartz.

The Vallito Member interfingers eastward with the Cuarteles Member, and locally with the Cejita Member of the Tesuque Formation. The Vallito Member interfingers westward with the Hernandez Member in a broad zone that includes much mixing between the two members (as much as 7 km wide). East of the Rio Grande, the Vallito Member is gradationally overlain by the Cuarteles Member of the Tesuque Formation. West of the Rio Grande, the two are in an interfingering and mixed relation with no complete progradation of the Cuarteles Member over the Vallito Member. The base of this member is coeval with the lower Cejita Member of the Tesuque Formation (see cross-section), and thus is ca. 13.2 Ma. In R-10, this member extends up to the basalt flow that likely correlates to 8.5-9.0 Ma flows to the west. Unit is over 900 m-thick, including where it is mixed with Hernandez Member detritus (cross-section).

TEPHRA BEDS

(refer to Explanation of Map Symbols for identifying respective tephra locations on the map)

Black-gray coarse ashes (upper Miocene) – Gray to light gray (10YR 5-7/1 and 10YR 7/2), coarse basaltic to andesitic ashes. Lower 15-25 cm locally is well to moderately cemented by calcium carbonate and composed of fine- to very coarse-sand-size ash with trace very fine-pebble-size lapilli. Lapilli becomes more common to the west but still is minor. Coarse ash is subrounded

to subangular, moderately to poorly sorted, and internally massive. Upper part of a given ash bed is generally weakly cemented, weakly consolidated, and composed of reworked ash that is finer grained than below (very fine- to medium-sand-size), slightly clayey to clayey, moderately sorted, subangular to subrounded, massive to planar-laminated, and mixed with varying amounts of arkosic sand. 10-160 cm thick.

White lapilli and coarse ashes (upper Miocene) – White coarse ash and fine pumice (the latter is up to 5 mm in diameter); includes intermediate to felsic volcanic lithics, and arkosic sand. Contains up to 5% biotite. A relatively continuous bed near the base of this zone is called the “marker coarse white ash.” 10-120 cm thick.

UNITS DEPICTED ONLY IN CROSS-SECTION

Mixed and interfingering Vallito and Hernandez Members of the Chamita Formation (Upper Miocene; shown as Tcv-Tch on cross-section): See descriptions for the Vallito Member and Hernandez Member.

Basin floor deposits of the Pojoaque Member of the Tesuque Formation (middle Miocene): Mostly of floodplain deposits of silt, very fine- to fine-grained sand, silty very fine- to fine-grained sand, and clay, based on well data for the OSE-USGS Buckman monitoring well and inspection of outcrops south of Highway 502 near Pojoaque Pueblo (Koning and Maldonado, 2001; Koning, 2002). There are subordinate, intercalated channel-fills composed of fine- to coarse-grained sand that are up to 4.6 m-thick. Beds are very thin to thick, broadly lenticular (10-30 m lateral length) to tabular (Koning, 2002). The sediment is pinkish gray to light gray to light brown to pale brown. In general, this sediment has a noticeable darker or grayer hue than that of the Vallito and Cejita Members up-section. The sand possesses 0 to 5% green-colored quartz grains together with very minor Paleozoic detritus (usually trace-3%, locally as much as 15%). Most of the sand, however, consists of quartz with 5-20% feldspar. Near the top, there may be as much as 10-15% quartzite, as seen in the Buckman-9 well. In some arkosic intervals (e.g., 2000-2090 ft in the OSE-USGS Buckman monitoring well), the feldspar percentage may be as high as 60%.

Faults near Buckman

There are numerous, relatively small, north-trending normal faults on the White Rock quadrangle. Of these, the two biggest faults are the east-down West Buckman fault and the west-down East Buckman fault east of Buckman (labeled in Figure 2). We interpret that two faults offset strata along the cross-section line based on our mapping and inter-well correlations; there may possibly be more faults not shown on the cross-section. To the east is a west-down fault located en echelon to the southern tip of the East Buckman

fault. This fault appears to have a throw magnitude of 20-25 m (70-80 ft) at the cross-section line. Interestingly, the East Buckman fault does not appear to displace strata in the upper part of the Cuarteles Member on the south slope of Buckman Mesa. Consequently, offset along the East Buckman fault appears to have slowed and then effectively stopped sometime after the emplacement of the coarse white ash marker bed (10.9 Ma) but before deposition of the upper part of the Cuarteles Member (probably 9-10.5 Ma). This fault appears to align with a west-down fault inferred in the northern part of the White Rock quadrangle.

To the west is an apparent east-down fault near Buckman well #8 (located near the Rio Grande immediately south of the mouth of Canada Ancha) having a throw magnitude of ~30 m (100 ft). However, the last movement along this fault appears to have been west-down with a significant component of lateral slip (Scott Minor of the U.S.G.S., personal communication, March-2007). This last movement has offset and deformed a terrace deposit (Qa1) which is overlain by well-laminated lacustrine sediment rich in silt and clay. Charcoal collected from the basal lacustrine sediments returned a possibly finite C-14 age of 43.2 ka (Reneau and Dethier, 1996). Interestingly, this fault does not appear to have offset the overlying lacustrine deposits. It is possible that the earthquake accompanying the last slip event on this fault triggered the landslide that dammed the Rio Grande and created the lake associated with the lacustrine sedimentation of the upper Qa1 terrace deposit.

Temporal changes in paleogeography and depositional environments during the Miocene

The Buckman and White Rock areas occupied a basin-floor to distal alluvial slope environment during the middle to late Miocene. During the latter stages of Pojoaque Member deposition (ca. 13.5-13.2 Ma), the southward-flowing streams on the basin-floor were characterized by subordinate, sandy-floored channels within extensive floodplains underlain by clay, silt, and very fine- to fine-grained sand. The Cejita Member heralded a more vigorous fluvial system from the north-northeast that left more abundant channel-fill deposits of sand and gravel. Likewise, reworking of Ojo Caliente Member eolian sand into the axial river resulted in more abundant sandy-bedload deposits (Vallito Member) compared to prior to ~13.2 Ma. This post-13.2 Ma, relatively coarse-grained deposition generally continued through the remainder of the Miocene, with some grossly finer-grained intervals preserved under the Buckman well field. During and after the waning of eolian deposition to the north (~11-12.8 Ma; Koning et al., 2005a and 2005b; Dethier et al., 1986; Aldrich and Dethier, 1990), an ancestral Rio Chama became established (Hernandez Member) and flowed south-southeast onto the floor of the Espanola Basin from the northwest, where it then flowed parallel alongside of the ancestral Rio Grande (Vallito and/or Cejita Member) before merging in the general study area. Westward-flowing alluvial-slope deposits (Cuarteles Member) progressively extended westwards during the late Miocene.

ACKNOWLEDGMENTS

San Ildefonso Pueblo graciously allowed access to their tribal lands for this project. Jack Frost provided borehole geophysical data for Buckman-09 and the SEO observation well. Funding for this project was provided by the AQUIFERMAP and STATEMAP programs as well as the Office of the New Mexico State Engineer. Descriptions for the OSE-USGS Buckman monitoring well were provided by Scott Aby. David Broxton and David Vaniman of the Los Alamos National Laboratories provided much assistance in describing and interpreting cuttings and down-hole geophysical logs for wells west of the Rio Grande. David Broxton also helped construct the west end of the cross-section.

REFERENCES

- Aldrich, M.J., and Dethier, D.P., 1990, Stratigraphic and tectonic evolution of the northern Española basin, Rio Grande rift, New Mexico: Geological Society of America Bulletin, v. 102, p. 1695-1705.
- Aubele, J.C., 1978, Geology of the Cerros del Rio volcanic field, Santa Fe, Sandoval, and Los Alamos Counties, New Mexico [unpublished M.S. thesis]: University of New Mexico, Albuquerque, New Mexico, 136 pp.
- Bachman, G.O., and Mehnert, H.H., 1978, New K-Ar dates and the late Pliocene to Holocene geomorphic history of the central Rio Grande region, New Mexico: Geological Society of America Bulletin, v. 89, no. 2, p. 283-292.
- Bailey, R.A., Smith, R.L., and Ross C.S., 1969, Stratigraphic nomenclature of volcanic rocks in the Jemez Mountains, New Mexico: U.S. Geological Survey Bulletin 1274-P., 19 p.
- Baldrige, W.S., 1979, Petrology and petrogenesis of Plio-Pleistocene basaltic rocks from the central Rio Grande rift, New Mexico, and their relation to rift structure, *in* Riecker, R.E., (ed.), Rio Grande rift – tectonics and magmatism: American Geophysical Union, Washington, D.C., p. 323-353.
- Barghoorn, S., 1981, Magnetic-polarity stratigraphy of the Miocene type Tesuque Formation, Santa Fe Group, in the Española Valley, New Mexico: Geological Society of America Bulletin, v. 92, p. 1027-1041.
- Biehler, S., Ferguson, J., Baldrige, W.S., Jiracek, G.R., Aldern, J.L., Martinez, M., Fernandez, R., Romo, J., Gilpin, B., Braile, L.W., Hersey, D.R., Luyendyk, B.P., and Aiken, C.L., 1991, A geophysical model of the Española Basin, Rio Grande rift, New Mexico: Geophysics, v. 56, no. 3, p. 340-353.
- Birkeland, P.W., 1999, Soils and geomorphology: New York, Oxford University Press, 430 p.
- Broxton, D and Vaniman D., 2005, Geologic framework of a groundwater system on the margin of a rift basin, Pajarito Plateau, North-Central New Mexico: Vadose Zone Journal, v. 4, pp. 522-550.

- Broxton D., Gilkeson, R., Longmire, P., Marin, J., Warren, R., Vaniman, D., Crowder, A., Newman, B., Lowry, B., Rogers, D., Stone, W., McLin, S., WoldeGabriel, G., Daymon, D., and Wycoff, D., 2001, Characterization well R-9 completion report: Los Alamos National Laboratory Report LA-13742-MS, 85 p.
- Cande, S.C., Kent, D.V., 1995, Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic: *Journal of Geophysical Research*, v. 100, n. B4, p. 6093-6095.
- Cavazza, W., 1986, Miocene sediment dispersal in the central Española Basin, Rio Grande rift, New Mexico, USA: *Sedimentary Geology*, v. 51, p. 119-135.
- Chapin, C.E., 1971, The Rio Grande rift; Part 1, Modifications and additions: *New Mexico Geological Society, 22nd Field Conference, Guidebook*, p. 191-201.
- Compton, R.R., 1985, *Geology in the field*: New York, John Wiley & Sons, Inc., 398 p.
- Dethier, D.P., 1997, *Geology of the White Rock quadrangle, Los Alamos and Santa Fe Counties, New Mexico*: New Mexico Bureau of Mines and Mineral Resources in cooperation with Los Alamos National Laboratory, University of California, Geologic Map 73, scale 1:24,000.
- Dethier, D.P., Aldrich, M.J., Jr., Shafiqullah, M., 1986, New K-Ar ages for Miocene volcanic rocks from the northeastern Jemez Mountains and Tejana Mesa, New Mexico: *Isotopes*, no. 47, p. 12-14.
- Dethier, D.P., Harrington, C.D., and Aldrich, M.J. Jr., 1988, Late Cenozoic rates of erosion in the western Española Basin, New Mexico – evidence from geologic dating of erosion surfaces: *Geological Society of America Bulletin*, v. 100, no. 6, p. 928-937.
- Galusha, T., and Blick, J.C., 1971, Stratigraphy of the Santa Fe Group, New Mexico: *Bulletin of the American Museum of Natural History*, v. 144, 127 p.
- Gile, L.H., Peterson, F.F., and Grossman, R.B., 1966, Morphological and genetic sequences of carbonate accumulation in desert soils: *Soil Science*, v. 101, p. 347-360.
- Griggs, R.L., 1964, *Geology and ground-water resources of the Los Alamos area, New Mexico*: U.S. Geological Survey Water Supply Paper 1753, 107 p., Map scale 1:24,000.
- Ingram, R.L., 1954, Terminology for the thickness of stratification and parting units in sedimentary rocks: *Geological Society of America Bulletin*, v. 65, p. 937-938, table 2.
- Izett, G.A., and Obradovich, J.D., 1994, $^{40}\text{Ar}/^{39}\text{Ar}$ age constraints for the Jaramillo Normal Subchron and the Matuyama-Brunhes geomagnetic boundary: *Journal of Geophysical Research*, v. 99, no. B2, p. 2925-2934.
- Izett, G.A., and Obradovich, J.D., 2001, $^{40}\text{Ar}/^{39}\text{Ar}$ ages of Miocene tuffs in basin-fill deposits (Santa Fe Group, New Mexico, and Troublesome Formation, Colorado) of the Rio Grande rift system: *The Mountain Geologist*, v. 38, no. 2, p. 77-86.
- Kelley, V.C., 1956, The Rio Grande depression from Taos to Santa Fe: *New Mexico Geological Society, 7th field conference, Guidebook*, p. 109-114.
- Koning, D.J., 2002, revised July-2005, *Geologic map of the Española 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico*: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM 54, scale 1:24,000.
- Koning, D.J., 2003, revised 2005, *Geologic map of the Chimayo 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico*: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM-71, scale 1:24,000.

- Koning, D.J., and Maldonado, F., 2001, Geologic map of the Horcado Ranch quadrangle, Santa Fe County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Open-file Geologic Map OF-GM-54, scale 1:24,000.
- Koning, D.J., and Manley, K., 2003, revised-2005, Geologic map of the San Juan Pueblo 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM-70, scale 1:24,000.
- Koning, D.J., Aby, S.B., and Dunbar, N., 2004a, Middle-upper Miocene stratigraphy of the Velarde graben, north-central New Mexico: Tectonic and paleogeographic implications: New Mexico Geological Society, 55th Field Conference, Guidebook, p. 359-373.
- Koning, D., J., and Aby, Scott B., 2005, Proposed Members of the Chamita Formation, north-central New Mexico: New Mexico Geological Society, 56th Field Conference Guidebook, Geology of the Chama Basin, p. 258-278.
- Koning, D.J., Connell, S.D., Pazzaglia, F.J., and McIntosh, W.C., 2002, Redefinition of the Ancha Formation and Pliocene-Pleistocene deposition in the Santa Fe embayment, north-central New Mexico: New Mexico Geology, v. 24, no. 3, p. 75-857.
- Koning, D.J., Skotnicki, S., Moore, J., and Kelley, S., 2005b, Geologic map of the Chili 7.5-minute quadrangle, Rio Arriba county, New Mexico: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM-81, scale 1:24,000.
- Koning, Daniel J., Connell, Sean D., Morgan, Gary S., Peters, Lisa, and McIntosh, William C., 2005, Stratigraphy and depositional trends in the Santa Fe Group near Espanola, north-central New Mexico -- tectonic and climatic implications: New Mexico Geological Society, 56th Field Conference Guidebook, Geology of the Chama Basin, p. 237-257.
- Machette, M.N., 1985, Calcic soils of the southwestern United States, *in* Weide, D.L. (ed.), Soils and Quaternary geology of the southwestern United States: Geological Society of America, Spioecial Paper 203, p. 1-21.
- Manley, K., 1976, The Late Cenozoic History of the Española Basin, New Mexico [Ph.D. thesis]: University of Colorado, 1-171 p.
- Manley, K., 1977, Geologic map of the Cejita Member (new name) of the Tesuque Formation, Española Basin, New Mexico: U.S. Geological Survey Miscellaneous Field Studies Map MF-877, scale 1:24000.
- Manley, K., 1979, Tertiary and Quaternary stratigraphy of the Northeast Plateau, Española Basin, New Mexico: New Mexico Geological Society Guidebook, 30th Field Conference, Guidebook, p. 231-236.
- Munsell Color, 1994 edition, Munsell soil color charts: New Windsor, N.Y., Kollmorgen Corp., Macbeth Division.
- North American Commission on Stratigraphic Nomenclature, 2005, The North American Stratigraphic Code: The American Association of Petroleum Geologists Bulletin, vol. 89, no. 11, p. 1547-1591.
- Reneau, S.L., and Dethier, D., 1996, Pliocene and Quaternary history of the Rio Grande, White Rock Canyon and vicinity, New Mexico: New Mexico Geological Society Guidebook, 47th Field Conference, Jemez Mountains Region, p. 317-324.

- Reneau, S.L., Gardner, J.N., and Forman, S.L., 1996, New evidence for the age of the youngest eruptions in the Valles caldera, New Mexico: *Geology*, v. 24, no. 1, p. 7-10.
- Sawyer, D.A., Shroba, R.B., Minor, S.A., and Thompson, R.A., 2002, Geologic map of the Tetilla Peak quadrangle, Santa Fe and Sandoval counties, New Mexico: U.S. Geological Survey Miscellaneous Field Studies Map MF-2352, Map scale 1:24,000.
- Self, S.A., Kircher, D.E., and Wolff, J.A., 1988, The El Cajete Series, Valles caldera, New Mexico: *Journal of Geophysical Research*, v. 93, no. B6, p. 6113-6127.
- Smith, G.A., Middle to late Cenozoic development of the Rio Grande Rift and adjacent regions in northern New Mexico, *in* Mack, G.H., and Giles, K.A., eds., *The Geology of New Mexico, A Geologic History*: New Mexico Geological Society, p. 331-358.
- Soil Survey Staff, 1992, *Keys to Soil Taxonomy*: U.S. Department of Agriculture, SMSS Technical Monograph no. 19, 5th edition, 541 p.
- Spell, T. L., McDougall, I., and Doulgeris, A.P., 1996, Cerro Toledo Rhyolite, Jemez volcanic field, New Mexico: $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of eruptions between two caldera-forming events: *Geological Society of America Bulletin*, v. 108, p. 1549–1566.
- Spiegel, Z., and Baldwin, B., 1963, *Geology and Water Resources of the Santa Fe Area, New Mexico*: Washington, D.C., Geological Survey Water-Supply Paper 1525, 258 p.
- Turbeville, B.N., Waresback, D.B., and Self, S., 1989, Lava dome growth and explosive volcanism in the Jemez Mountains, New Mexico – evidence from the Pliocene-Pleistocene Puye alluvial fan: *Journal of Volcanology and Geothermal Research*, v. 36, p. 267-291.
- Waresback, D.B., 1986, *The Puye Formation, New Mexico – analysis of a continental rift-filling volcanoclastic alluvial fan sequence* [M.S. thesis]: University of Texas, Arlington, 225 pp.
- Waresback, D.B., and Turbeville, B.N., 1990, Evolution of a Pliocene-Pleistocene volcanogenic alluvial fan – the Puye Formation, Jemez Mountains, New Mexico: *Geological Society of America Bulletin*, v. 102, p. 298-314.
- Udden, J.A., 1914, The mechanical composition of clastic sediments: *Bulletin of the Geological Society of America*, v. 25, p. 655-744.
- Wentworth, C.K., 1922, A scale of grade and class terms for clastic sediments: *Journal of Geology*, v. 30, p. 377-392.
- WoldeGabriel, G., Laughlin, A.W., Dethier, D.P., and Heizler, M., 1996, Temporal and geochemical trends of lavas in White Rock Canyon and the Pajarito Plateau, Jemez volcanic field, New Mexico, USA: *New Mexico Geological Society, 47th Field Conference Guidebook, the Jemez Mountains region*, p. 251-261.

FIGURE CAPTIONS

FIGURE 1: Location of White Rock quadrangle (shown by orange shading) relative to the cities of Santa Fe and White Rock, New Mexico. .

FIGURE 2: Geologic map of the Buckman and White Rock areas within the White Rock quadrangle. The A-A' cross-section line and its east end are shown (at longitude 106° 7.5'), but the west end of the cross-section is located 1.53 km west of the boundary of this figure. The locations of the stratigraphic sections of Figure 3 are also depicted. Explanation for unit shading, patterns, and associated labels is given in the cross-section.

FIGURE 3: Stratigraphic fence diagram showing correlations between the R-16 well, Overlook section, lower Buckman section, and the upper Buckman section (left to right, west to east). Thick black lines are possible tie lines between tephra beds. Background shading illustrates stratigraphic relations between the sections. JMAF = Jemez Mountains alluvial fan sediment. Detailed descriptive data for these sections are found in Appendix 1 of the report.


TABLE 1. Clast count data for Vallito and Hernandez Members, Chamita Formation

COMMENTS TO MAP USERS

A geologic map displays information on the distribution, nature, orientation, and age relationships of rock and deposits and the occurrence of structural features. Geologic and fault contacts are irregular surfaces that form boundaries between different types or ages of units. Data depicted on this geologic quadrangle map are based on reconnaissance field geologic mapping, compilation of published and unpublished work, and photogeologic interpretation. Locations of contacts are not surveyed, but are plotted by interpretation of the position of a given contact onto a topographic base map; therefore, the accuracy of contact locations depends on the scale of mapping and the interpretation of the geologist(s). Any enlargement of this map could cause misunderstanding in the detail of mapping and may result in erroneous interpretations. Site-specific conditions should be verified by detailed surface mapping or subsurface exploration. Topographic and cultural changes associated with recent development may not be shown.

The map has not been reviewed according to New Mexico Bureau of Mines and Mineral Resources standards. Revision of the map is likely because of the on-going nature of work in the region. The contents of the report and map should not be considered final and complete until reviewed and published by the New Mexico Bureau of Mines and Mineral Resources. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the State of New Mexico, or the U.S. Government. Cross-sections are constructed based upon the interpretations of the authors made from geologic mapping, and available geophysical (regional gravity and aeromagnetic surveys), and subsurface (drillhole) data.

Cross-sections should be used as an aid to understanding the general geologic framework of the map area, and not be the sole source of information for use in locating or designing wells, buildings, roads, or other man-made structures.



DESCRIPTION OF MAP UNITS

- Qa1 Alluvial deposit (Holocene)** - Cross-bedded to plane bedded sand and pebbly sand and thin beds of silty sand exposed along the Rio Grande, along active channels of tributary arroyos, and on adjacent low (<4 m, <13 ft) surfaces.
- Qa2 Alluvial deposit (Holocene and late Pleistocene)** - Well-sorted cobble to boulder gravel, cross-bedded sand, and thin-bedded sand and silty sand beneath terraces along the Rio Grande; cobble gravel along Los Alamos Canyon and other steep tributaries; sand and silty sand near Cañada Ancha; and gravelly sand and colluvial soil on upland parts of the Cerros del Rio.
- Qa3 Alluvial deposit (Holocene and late Pleistocene)** - Well-sorted cobble to boulder gravel and rhythmically bedded fine sand, silt, and silty clay exposed beneath terrace remnants 25-45 to (80-150 ft) above the Rio Grande.
- Qa4 Alluvial fan deposit (Holocene and late Pleistocene)** - Cross-bedded, poorly sorted cobble to boulder gravel deposited mainly along White Rock Canyon by high-gradient, ephemeral channels.
- Qa5 Alluvial fan deposit (late to middle Pleistocene)** - Cross-bedded, poorly sorted cobble to boulder gravel and poorly sorted, matrix-rich debris-flow deposits that form remnants 5-15 m (16-50 ft) above adjacent QZ deposits.
- Qa6 Cultural deposit (Holocene to middle Pleistocene)** - Rockfall, debris-flow, and poorly sorted alluvial deposits at the bases of cliffs, particularly along White Rock Canyon. Texture and clay lithology variable.
- Qa7 Thin slopewash and colluvial deposit (Holocene to late Pleistocene)** - Light yellowish brown to pale brown to brownish yellow, silt and fine- to medium-grained sand mixed with varying amounts of coarser sand and gravel. This unit thin caps high-level surfaces. The underlying unit is also shown. Weedy to moderately consolidated.
- Qa8 Landslide deposit (Holocene to late Pleistocene)** - Massive slumps and debris flows rich in basaltic-boulder gravel mainly exposed along White Rock Canyon.
- Qa9 north of Chapequi Canyon, and along Cañada Ancha.** Most slides are inactive. Deposits consist of massive slump blocks with coherent internal stratigraphy near canyon rims and progressively deformed slumps and debris flows closer to the Rio Grande.
- Santa Fe Group deposits exposed in Quaternary landscapes (not shown in correlation of map units)**
 - Qb5-cutf Exposures of fine-grained Cuartetes Member (Chamita Formation) involved with landslide-related subsidence and deformation** - See description of Cuartetes Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-cu Exposures of Cuartetes Member (Chamita Formation) involved with landslide-related subsidence and deformation** - See description of Cuartetes Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-tecf Exposures of fine-grained Cuartetes Member (Tesque Formation) involved with landslide-related subsidence and deformation** - See description of Cuartetes Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-teu Exposures of Cuartetes Member (Tesque Formation) involved with landslide-related subsidence and deformation** - See description of Cuartetes Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-ev Exposures of Vallito Member (Chamita Formation) involved with landslide-related subsidence and deformation** - See description of Vallito Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-ch Exposures of Hernandez Member (Chamita Formation) involved with landslide-related subsidence and deformation** - See description of Hernandez Member below. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-pf Exposures of Puye Formation involved with landslide-related subsidence and deformation** - See description of Puye Formation. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-pt Exposures of axial river deposits of the Puye Formation (Totavi Lenti) involved with landslide-related subsidence and deformation** - See description of Puye Formation. Bedding is tilted in various directions and somewhat deformed but generally intact.
 - Qb5-cm Exposures of phreatomagmatic deposits involved with landslide-related subsidence and deformation** - See description of unit Tcm below. Bedding is tilted in various directions and somewhat deformed but generally intact.

PLEISTOCENE-PLIOCENE VOLCANIC STRATA

- Ta Ancha Formation (upper Pliocene)** - Brownish yellow to light yellowish brown sand and gravel under the Cerros del Rio volcanic flows (units Tba and Tcb) and phreatomagmatic deposits (Tcm); unit is also interlayered with these volcanic flows. Overlies the Puye Formation (Tp) and local basal flows near the Rio Grande; to the east the Ancha Formation unconformably overlies Miocene-age Cuartetes Member deposits of the Tesque Formation (Koning and Maldonado, 2001; Koning et al., 2002).
- Tp Puye Formation (lower Pliocene)** - Weakly lithified pebbly to boulder gravel, boulder-rich debris flows, massive to planar-bedded sand, thin (<1 m, <3 ft) beds of dacitic tephra and pumiceous alluvium, and beds of fine sand and silt. Exposed west of the Rio Grande except for an isolated outcrop south of the mouth of Cañada Ancha (Section F of Dethier, 1997).
- PELSTOCENE-PLIOCENE VOLCANIC DEPOSITS AND ROCKS**
 - Qet El Cajete tephra (late Pleistocene)** - Pumiceous lapilli of rhyolitic composition. Forms surface layer in isolated outcrops as thick as 40 cm in southwest part of map area and <5 cm in northeast part of map area. Most exposures have been reworked by slope processes. Derived from the El Cajete vent in the Valles calderas; previously thought to be about 150,000 yrs old (Self et al., 1988) but may be as young as 50 ka (Rena et al., 1996).
 - Qbtu Banderil Tuff, upper part (early Pleistocene)** - Slightly welded pyroclastic flows (Tshireg Member) and a thin (<1 m, <3 ft) pumiceous fall unit (Tskanki Pumice Bed), both of rhyolitic composition. Dominates surface west of the Rio Grande and forms one prominent outcrop east of the Rio Grande (Section H of Dethier, 1997). Follows direction to the east. Derived from the Valles caldera west of the map area. Age 1.2 Ma (Zett and Obradovich, 1994).
 - Qbtl Banderil Tuff, lower part (early Pleistocene)** - Slightly welded pyroclastic flows (Owens Member) of pumiceous rhyolite and a component pumiceous ash unit (Owens Member) as thick as 60 m (200 ft), also of rhyolitic composition. Derived from the Valles caldera west of the map area. Age is 1.6 Ma (Zett and Obradovich, 1994).
 - Tcc Cinder cone deposit (Pliocene)** - Oxidized cinders, agglomerate, and minor areas of phreatomagmatic deposits composed of olivine basaltic andesite and basalt. Bedded to massive fall, surge, and flow deposits composed of basaltic tuff and cinders and accidental fragments of the Santa Fe Group. Thickest exposures along the Rio Grande and Cañada Ancha (Section G of Dethier, 1997).
 - Tcm Phreatomagmatic deposit (Pliocene)** - Bedded to massive fall, surge, and flow deposits composed of basaltic tuff and cinders and accidental fragments of the Santa Fe Group. Thickest exposures along the Rio Grande and Cañada Ancha (Section G of Dethier, 1997).

EXPLANATION OF MAP SYMBOLS

- Geologic contact** - Solid where exposed, dashed where approximately located; dotted where concealed, queried where uncertain.
- Deleted geologic contact from previous mapping** - Dashed line with a cross-hatch pattern.
- Normal fault** - Solid where exposed, dashed where approximately located; dotted where concealed, queried where uncertain. Ball and bar are on the downthrown block. Number by arrow denotes the location and value of fault dip measurement. \bar{R} - Fault that is probably a landslide block.
- True strike and dip of bedding** - Arrow with a number indicating the dip.
- Apparent strike and dip of bedding** - Arrow with a number indicating the dip.
- Beds that dip less than 2 degrees and whose strike cannot be precisely determined** - Arrow with a number indicating the dip.
- Bi-directional paleocurrent measurement** - Solid ball on vector is located at measurement location. Primarily measured using charcoal trends (abbreviated as "C") or the trend of a single gravel margin (abbreviated as "G"). Although bi-directional, only one arrow head may be depicted that shows the author's preferred direction; this preference is based on surrounding mono-directional paleocurrent data and source area considerations. A double-headed arrow is shown where there is no preference. Different lengths of vector are proportional to the number of data measurements at a site (5-dm measurements in feet (ft)). The number of data measurements is shown by the number preceding the vector. Paleocurrent azimuth values are also listed next to the vector.
- Mono-directional paleocurrent measurement** - Solid ball on vector is located at measurement location. Primarily measured using clay imbrication or trough cross-stratification (abbreviated as "C") or the trend of a single gravel margin (abbreviated as "G"). Different lengths of the vector are proportional to the number of data measurements at a site (5-dm measurements in feet (ft)). The number of data measurements is shown by the number preceding the vector. Paleocurrent azimuth values are also listed next to the vector.
- Back-gravel coarse ash** - See description of map units.
- White lapilli and coarse ash** - See description of map units.
- Top of landslide block in slide complex** - Dashed line with a cross-hatch pattern.
- Cross-section and point** - Arrow with a number indicating the dip.
- Supply of groundwater monitoring well; name of well listed alongside (B stands for Buckman)** - Arrow with a number indicating the dip.

Preliminary geologic map of the White Rock quadrangle, Santa Fe, Sandoval and Los Alamos Counties, New Mexico.

May 2007
by David P. Dethier and Daniel J. Koning
Department of Geosciences, Williams College, 947 Main Street, Williamstown, MA 01267
NMBGMR, New Mexico Tech, 801 Leroy Place, Socorro, NM 87801-4796

COMMENTS TO MAP USERS

A geologic map displays information on the distribution, nature, orientation, and age relationships of rock and deposits and the occurrence of structural features. Geologic and fault contacts are irregular surfaces that form boundaries between different types or ages of units. Data depicted on this geologic quadrangle map may be based on any of the following: reconnaissance field geologic mapping, compilation of published and unpublished work, and photogeologic interpretation. Locations of contacts are not surveyed, but are plotted by interpretation of the position of a given contact onto a topographic base map; therefore, the accuracy of contact locations depends on the scale of mapping and the interpretation of the geologists. Any enlargement of this map could be misleading in the detail of mapping and may result in erroneous interpretations. Site-specific conditions should be verified by detailed surface mapping or subsurface exploration. Topographic and cultural changes associated with recent development may not be shown.

Cross sections are constructed based upon the interpretation of the author made from geologic mapping and available geophysical and subsurface (driftless) data. Cross sections should be used as an aid to understanding the general geologic framework of the map area, and not be the sole source of information for use in locating or designing wells, buildings, roads, or other man-made structures.

The map has not been reviewed according to New Mexico Bureau of Geology and Mineral Resources standards. The contents of the report and map should not be considered final and complete until reviewed and published by the New Mexico Bureau of Geology and Mineral Resources. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the State of New Mexico, or the U.S. Government.



QUADRANGLE LOCATION



This draft geologic map is preliminary and will undergo revision. It was produced from either scans of hand-drafted originals or from digitally drafted original maps and figures using a wide variety of software, and is currently in cartographic production. It is being distributed in this draft form as part of the bureau's Open-File map series (OFGM), due to high demand for current geologic map data in these areas where STATEMAP quadrangles are located, and it is the bureau's policy to disseminate geologic data to the public as soon as possible.

After this map has undergone scientific peer review, editing, and final cartographic production adhering to bureau map standards, it will be released in our Geologic Map (GM) series. This final version will receive a new GM number and will supersede this preliminary open-file geologic map.

DRAFT

New Mexico Bureau of Geology and Mineral Resources
New Mexico Tech
801 Leroy Place
Socorro, New Mexico
87801-4796
[505] 835-5490
http://geoinfo.nmt.edu

This and other STATEMAP quadrangles are (or soon will be) available for free download in both PDF and ArcGIS formats at:
http://geoinfo.nmt.edu/publications/maps/geologic/ofgm/home.html

- Tcm Basalt and interlayered phreatomagmatic deposits (Pliocene)** - Thin (<10 m, <33 ft) basaltic flows interlayered with basaltic phreatomagmatic rocks, mainly surge and flow deposits, and mapped in the southern part of the map area near the Rio Grande.
- Tei Basaltic intrusion (Pliocene)** - Olivine, pyroxene basaltic andesite containing quartz xenocrysts. Forms fine-grained dikes and small, shallow intrusions, with sharp chilled margins, associated with inners and cinder cones along the east margin of White Rock Canyon.
- Tea Andesite (early Pleistocene) and late Pliocene** - Massive, steep-dip silts, agglomerate, and minor areas of hypersthene andesite (Table 2) exposed in the southeast part of the map area. Strongly sheared near flow surface; brecciated at bases.
- Teb1 Basalt (late Pliocene)** - Theolitic olivine basalt flows, pillow basalt, and galenitic breccia exposed west of the Rio Grande north of Chapequi Canyon.
- Teb2 Basaltic andesite and related flows (Pliocene)** - Massive flows of olivine hypersthene basaltic andesite, andesite, and basaltic, containing as much as 7% quartz, exposed along the Rio Grande.
- Teb3 Basaltic flows (Pliocene)** - Thin (<10 m, <33 ft) flows of olivine basalt (hawaiite, see Table 2 of Dethier, 1997) containing <5% quartz xenocrysts and exposed mainly beneath large areas of Saguahash Plains (Sections F and G of Dethier, 1997) and along the east boundary of White Rock quadrangle.
- Teb4 Basaltic flows (Pliocene)** - Olivine basalt flows and more evolved rocks exposed within 20 m (66 ft) of the present level of the Rio Grande and Cañada Ancha. Flows are thin to massive and include a lava lake (Caja del Rio Canyon) and a number of poorly exposed outcrops along the Rio Grande from Water Canyon to Chapequi Canyon.
- Tebu Basalt, undivided (Pliocene)** - Olivine basalt, hawaiite, and basaltic andesite containing 2-8% quartz xenocrysts and exposed in the southern part of the map area (Section I of Dethier, 1997).

MIDDLE-LATE MIOCENE BASIN-FILL DEPOSITS (SANTA FE GROUP)

- Twep Chamita Formation, Vallito Member interbedded with western piedmont deposits (upper Miocene)** - Only found in outcrop at the mouth of Ancha Canyon. Light brown to reddish yellow (1.5YR 6/4-6) silt, very fine- to fine-grained sandstone, and minor silt intercalated with <25% coarse-grained channel fills.
- Tecuf Tesque Formation, fine-grained Cuartetes member east of Rio Grande (Upper Miocene)** - Pink to light brown to reddish yellow to light yellowish brown, silty very fine- to fine-grained sandstone, siltstone, and mudstone. Coarse channel-fills of pebbly sandstone and sandy pebble-conglomerate comprise about 3-25% of the unit. Fine-grained sediment outside of coarse-grained channel-fills is very thin to thick, tabular beds that may be internally laminated.
- Tecuf Chamita Formation, fine-grained Cuartetes Member west of the Rio Grande (Upper Miocene)** - See description above. Here, the Cuartetes Member intertongues westward with Vallito Member and the Hernandez Member of the Chamita Formation.
- Tecu Tesque Formation, Cuartetes Member (Upper Miocene)** - Sandy pebble conglomerate and pebbly sandstone comprising coarse channel fill complexes, with subordinate slightly clayey-silty sand containing <5% pebbles that was deposited by lower-energy flow outside of confined channels - the latter is referred to below as the "finer sediment" of this unit. Interbedded within this unit are sparse beds of block coarse ash and white coarse ash-lapilli.
- Tee1 Cejita Formation, Tesque Formation (Upper Miocene)** - Sandy and gravelly channel-fills that are extensively, very thin to thinly cross-stratified (up to 1-1 m-thick foresets) within 1-2 m-thick channel fills of pebbly very fine- to very coarse-grained sand and sandy pebble-cobble conglomerate. Clay lithology types are dominated by subordinate sandstone, limestone, and siltstone with an estimated 10-50% granite and 5-8% quartzite.
- Teb2 Chamita Formation, Hernandez Member (Upper Miocene)** - Light gray channel-fills with foliaceous fine-grained floodplain deposits. Amalgamated channel-fills per packages as much as 18 m thick. These channel-fills locally fine-upwards into horizontal-bedded floodplain deposits of clay, silt, and clayey-silty very fine- to fine-grained sand.
- Tev Chamita Formation, Vallito Member (Upper Miocene)** - Broad channel-fills (typically >10 m of meters wide) consisting of sand-dominated sediment in horizontal to cross-stratified, laminated to very thin to medium beds. Slacked channel-fills form amalgamated packages as much as 12 m thick.

The Vallito Member interfingers eastward with the Cuartetes Member, and locally with the Cejita Member of the Tesque Formation. The Vallito Member interfingers westward with the Hernandez Member in a broad zone that includes much mixing between the two members (as much as 7 km wide). East of the Rio Grande, the Vallito Member is gradually overlain by the Cuartetes Member of the Tesque Formation.

TEPHRA BEDS

- Black-gray coarse ash (upper Miocene)** - Gray to light gray (10YR 5/7-1 and 10YR 7/2), coarse basaltic to andesitic ashes. Lower 15-25 cm locally is well to moderately cemented by calcium carbonate and composed of fine- to very coarse sand-size ash with trace very fine-pebble-size lapilli. Lapilli becomes more common to the west but still is minor. Coarse ash is truncated to subangular, moderately to poorly sorted, and internally massive. Upper part of a given ash bed is generally weakly cemented, weakly consolidated, and composed of rounded ash that is finer grained than below (very fine- to medium sand-size), slightly clayey, moderately sorted, subangular to subrounded, massive to planar-laminated, and mixed with varying amounts of arkosic sand, 10-160 cm thick.
- White lapilli and coarse ash (upper Miocene)** - White coarse ash and fine pumice (the latter is up to 5 mm in diameter), includes intermediate to felsic volcanic lithics, and arkosic sand. Contains up to 5% biotite. A relatively continuous bed near the base of this zone is called the "marker coarse white ash", 10-120 cm thick.

UNITS DEPICTED ONLY IN CROSS-SECTION

Mixed and interfingering Vallito and Hernandez Members of the Chamita Formation (Upper Miocene; shown as Tcv-Tch on cross-section): See descriptions for the Vallito Member and Hernandez Member.

Basin floor deposits of the Pojoaque Member of the Tesque Formation (middle Miocene)

Mostly of floodplain deposits of silt, very fine- to fine-grained sand, silty very fine- to fine-grained sand, and clay, based on well data for the OSE-USGS Buckman monitoring well and inspection of outcrops south of Highway 502 near Pojoaque Pueblo (Koning and Maldonado, 2001; Koning, 2002).

REFERENCES

Arbuckle, J.C., 1978. Geology of the Cerros del Rio volcanic field, Santa Fe, Sandoval, and Los Alamos Counties, New Mexico (unpublished M.S. thesis). University of New Mexico, Albuquerque, New Mexico, 130 pp.

Bullington, C.O., and Malin, H.H., 1975. New K-Ar dates and the late Pliocene-Holocene geomorphic history of the central Rio Grande region, New Mexico. Geological Society of America Bulletin, v. 86, p. 20-28.

Bullington, W.S., 1979. Probing and progress of the Pleistocene basaltic rocks from the central Rio Grande rift, New Mexico, and their relation to rift structure. In Becker, R.B., ed., Rio Grande rift: tectonics and geomorphology. American Geophysical Union, Washington, D.C., p. 23-50.

Burgoyne, S., 1961. Geologic interpretation of the Miocene type Tesque Formation, Santa Fe Group, in the Española Valley, New Mexico. Geological Society of America Bulletin, v. 72, p. 1027-1041.

Burgoyne, S., 1967. Sedimentology. New York, Oxford University Press, 487 p.

Brown, D., Gibson, R., Langford, P., Martin, J., Warren, R., Vannote, D., Crowder, A., Newman, B., Lovers, B., Rogers, D., Stone, W., McLean, S., WoldeGabriel, G., Dymov, O., and Whittow, D., 2001. Geologic map of the Los Alamos National Laboratory Region, Los Alamos National Laboratory Report LA-1774-MS, 86 p.

Carr, S.C., Koen, D.V., 1995. Revised calibration of the potassium-argon system for the Late Cretaceous and Cenozoic. Journal of Geophysical Research, v. 100, p. B4, p. 999-1009.

Carpenter, R.B., 1985. Geology in the field. New York, John Wiley & Sons, Inc., 389 p.

Dethier, D.P., 1997. Geology of the White Rock quadrangle, Santa Fe and Sandoval Counties, New Mexico. New Mexico Bureau of Mines and Mineral Resources in cooperation with Los Alamos National Laboratory, University of California, Geologic Map 73, scale 1:24,000.

Dethier, D.P., Hartung, C.B., and Koning, D.J., 1988. Late Cenozoic cones of erosion in the eastern Española Basin, New Mexico - evidence from geologic dating of erosion surfaces. Geological Society of America Bulletin, v. 100, no. 9, p. 929-937.

Gaillardet, J., and Bockheim, J.G., 1971. Stratigraphy of the Santa Fe Group, New Mexico. Bulletin of the American Museum of Natural History, v. 144, 127 p.

Gil, L.H., Peterson, F.F., and Grouman, R.B., 1966. Morphological and genetic sequences of carbonates accumulation in desert soils. Soil Science, v. 101, p. 347-360.



FIGURE 1

CROSS-SECTION A-A' OF WHITE ROCK QUADRANGLE

by Daniel Koning and David Broxton 2007



FIGURE 2

EXPLANATION OF UNITS IN CROSS-SECTION AND FIGURE 2.

- QUATERNARY UNITS**
 - Qa Quaternary alluvium
 - Qa1 Landslide deposits (may include colluvium west of Rio Grande)
 - Qa2 Banderil Tuff
- PLIOCENE**
 - Tba Cerros del Rio Volcanic Field: Basalt and basaltic andesite
 - Tcb Cerros del Rio Volcanic Field: Phreatomagmatic deposits
 - Tcp Puye Formation: Total Lenti (total river deposits)
 - Tct Puye Formation: Alluvial fan deposits
- MIOCENE**
 - Tch Chamita Formation
 - Tch1 Hernandez Member
 - Tch2 Cejita Member
 - Tcv Vallito Member (so mostly coarse deposits beneath Buckman well field; i.e. mostly fine deposits under Buckman well field)
 - Tcu Mixed and intercalated Vallito and Hernandez Mbrs
 - Tcu1 Cuartetes Member west of Rio Grande
 - Tcu2 Cuartetes Member east of Rio Grande
 - Tcu3 fine-grained, distal Cuartetes Member west of Rio Grande
 - Tcu4 fine-grained, distal Cuartetes Member east of Rio Grande
 - Tcm Miocene basalt

- Griggs, R.L., 1964. Geology and ground-water resources of the Los Alamos area, New Mexico. U.S. Geological Survey Water Supply Paper 1573, 107 p., Map scale 1:24,000.
- Hagner, R.L., 1954. Terminology for the direction of stratification and parting units in sedimentary rocks. Geological Society of America Bulletin, v. 65, p. 977-978.
- Lee, G.A., and Obradovich, D.J., 1967. Age constraints for the Ancha and Cañada Ancha volcanic fields and the Ancha and Cañada Ancha phreatomagmatic deposits. Journal of Geophysical Research, v. 72, p. 2623-2634.
- Leet, C.A., and Obradovich, D.J., 2007. Age of Miocene tuffs in basin fill deposits (Santa Fe Group, New Mexico, and Troadstone Formation, Colorado) of the Rio Grande rift system. The Mountain Geologist, v. 38, no. 2, p. 77-81.
- Koning, D.J., 2002. Revised 2002. Geologic map of the Española 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico. New Mexico Bureau of Geology and Mineral Resources, Open-File Geologic Map (OF-GM 54), scale 1:24,000.
- Koning, D.J., 2003. Revised 2003. Geologic map of the Chapequi 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico. New Mexico Bureau of Geology and Mineral Resources, Open-File Geologic Map (OF-GM 55), scale 1:24,000.
- Koning, D.J., and Maldonado, F., 2001. Geologic map of the Huelmo Reach quadrangle, Santa Fe County, New Mexico. New Mexico Bureau of Mines and Mineral Resources, Open-File Geologic Map (OF-GM 53), scale 1:24,000.
- Koning, D.J., and Mader, K., 2001. Revised 2001. Geologic map of the San Juan Pueblo 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico. New Mexico Bureau of Geology and Mineral Resources, Open-File Geologic Map (OF-GM 56), scale 1:24,000.
- Koning, D.J., and Scott, B., 2001. Proposed Members of the Chamita Formation, north-central New Mexico. New Mexico Geological Society, 56th Field Conference Guidebook, Geology of the Chamita Basin, p. 217-220.
- Koning, D.J., Corbett, S.D., Pratziger, E.J., and Malin, H.H., 2002. Redefinition of the Ancha Formation and Pliocene-Pleistocene deposition in the Santa Fe embayment, north-central New Mexico. New Mexico Geological Society, 57th Field Conference Guidebook, Geology of the Chamita Basin, p. 217-220.
- Malin, H.H., 1985. Calcic soils of the southwestern United States. In Wade, D.L., ed., Soils and Quaternary geology of the southwestern United States. Geological Society of America Special Paper 203, 1-13.
- Mander, K., 1976. The Late Cenozoic History of the Española Basin, New Mexico (Ph.D. thesis). University of Colorado, 1-171 pp.
- Mason, C.A., 1964. Cities, Mines and climate implications. New Mexico, N.Y., Kollman, C.P., Malin, H.H., ed., Geology of the Chamita Basin, p. 217-220.
- Rena, S.L., Gardner, J.N., and Forman, S.L., 1996. New evidence for the age of the youngest tephra in the Valles caldera, New Mexico. Geological Society of America Bulletin, v. 108, no. 1, p. 7-10.
- Self, S.A., Richter, D.D., and Wright, J.A., 1988. The El Cajete vent in the Valles caldera, New Mexico. Journal of Geophysical Research, v. 93, no. B6, p. 6111-6127.
- Soil Survey Staff, 1992. Keys to Soil Taxonomy. U.S. Department of Agriculture, 38MS Technical Monograph, 19, 388 edition, 541 p.
- Spiegel, Z., and Malin, H.H., 1985. Geology and Water Resources of the Santa Fe Area, New Mexico. Washington, D.C., Geological Survey Water Supply Paper 1523, 258 pp.
- Tarver, B.N., Wareback, D.B., and Sol, S., 1989. Lava dome growth and explosive volcanism in the Jemez Mountains, New Mexico - evidence from the Plio-Pleistocene. Journal of Volcanology and Geothermal Research, v. 36, p. 207-294.
- Valdez, J.A., 1974. The mechanical composition of clastic sediment. Bulletin of the American Association of Petroleum Geologists, v. 55, p. 65-74.
- Wright, C.K., 1922. A study of fault and cause for the Chamita Basin. Journal of Geology, v. 30, p. 377-392.
- WoldeGabriel, G., Langhi, A.W., Dethier, D.P., and Heiler, M., 1998. Temporal and geochronological trends in the White Rock Canyon and the Pajarito Plateau, Jemez volcanic field, New Mexico, USA. New Mexico Geological Society, 47th Field Conference Guidebook, the Jemez Mountains region, p. 241-261.

FIGURE CAPTIONS

FIGURE 1: Location of White Rock quadrangle (shown by orange shading) relative to the cities of Santa Fe and White Rock, New Mexico.

FIGURE 2: Geologic map of the Buckman and White Rock areas within the White Rock quadrangle. The A-A' cross-section line and its east end are shown at longitude 106° 7.5', but the west end of the cross-section is located 1.53 km west of the boundary of this map. The locations of the stratigraphic sections of Figure 3 are also depicted. Explanation for unit labeling, patterns, and associated labels is given in the cross-section.

FIGURE 3: Stratigraphic three diagram showing correlations between the R-16 well, Overlook section, lower Buckman section, and the upper Buckman section (left to right, west to east). Thick black lines are possible line between tephra beds. Background shading illustrates stratigraphic relations between the sections. JMAF = Jemez Mountains-derived volcanoclastic alluvial fan deposits. Detailed descriptive data for these sections are found in Appendix 1 of the report.




FIGURE 3

EXPLANATION OF SYMBOLS

- Blank only sand
- Sandy pebbles
- White coarse ash-fine lapilli
- Cross-stratification
- Trough cross-stratification
- Revolving
- Silt and clay
- Horizontal bedded sand

SEDIMENTARY ROCKS




FIGURE 3

EXPLANATION OF UNITS IN CROSS-SECTION AND FIGURE 2.

- QUATERNARY UNITS**
 - Qa Quaternary alluvium
 - Qa1 Landslide deposits (may include colluvium west of Rio Grande)
 - Qa2 Banderil Tuff
- PLIOCENE**
 - Tba Cerros del Rio Volcanic Field: Basalt and basaltic andesite
 - Tcb Cerros del Rio Volcanic Field: Phreatomagmatic deposits
 - Tcp Puye Formation: Total Lenti (total river deposits)
 - Tct Puye Formation: Alluvial fan deposits
- MIOCENE**
 - Tch Chamita Formation
 - Tch1 Hernandez Member
 - Tch2 Cejita Member
 - Tcv Vallito Member (so mostly coarse deposits beneath Buckman well field; i.e. mostly fine deposits under Buckman well field)
 - Tcu Mixed and intercalated Vallito and Hernandez Mbrs
 - Tcu1 Cuartetes Member west of Rio Grande
 - Tcu2 Cuartetes Member east of Rio Grande
 - Tcu3 fine-grained, distal Cuartetes Member west of Rio Grande
 - Tcu4 fine-grained, distal Cuartetes Member east of Rio Grande
 - Tcm Miocene basalt

EXPLANATION OF UNITS IN CROSS-SECTION AND FIGURE 2.

- QUATERNARY UNITS**
 - Qa Quaternary alluvium
 - Qa1 Landslide deposits (may include colluvium west of Rio Grande)
 - Qa2 Banderil Tuff
- PLIOCENE**
 - Tba Cerros del Rio Volcanic Field: Basalt and basaltic andesite
 - Tcb Cerros del Rio Volcanic Field: Phreatomagmatic deposits
 - Tcp Puye Formation: Total Lenti (total river deposits)
 - Tct Puye Formation: Alluvial fan deposits
- MIOCENE**
 - Tch Chamita Formation
 - Tch1 Hernandez Member
 - Tch2 Cejita Member
 - Tcv Vallito Member (so mostly coarse deposits beneath Buckman well field; i.e. mostly fine deposits under Buckman well field)
 - Tcu Mixed and intercalated Vallito and Hernandez Mbrs
 - Tcu1 Cuartetes Member west of Rio Grande
 - Tcu2 Cuartetes Member east of Rio Grande
 - Tcu3 fine-grained, distal Cuartetes Member west of Rio Grande
 - Tcu4 fine-grained, distal Cuartetes Member east of Rio Grande
 - Tcm Miocene basalt